



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2022 – 01:20 pm BST

PDB ID : 7Z4N
Title : Plasmodium falciparum pyruvate kinase complexed with pyruvate
Authors : Dillenberger, M.; Rahlfs, S.; Becker, K.; Fritz-Wolf, K.
Deposited on : 2022-03-04
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.29
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	616	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	0	4	0
			3782	2381	651	719	31			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	512	LEU	-	expression tag	UNP C6KTA4
A	513	GLU	-	expression tag	UNP C6KTA4
A	514	HIS	-	expression tag	UNP C6KTA4
A	515	HIS	-	expression tag	UNP C6KTA4
A	516	HIS	-	expression tag	UNP C6KTA4
A	517	HIS	-	expression tag	UNP C6KTA4
A	518	HIS	-	expression tag	UNP C6KTA4
A	519	HIS	-	expression tag	UNP C6KTA4


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is PYRUVIC ACID (three-letter code: PYR) (formula: C₃H₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
7	A	1	12	6	1	4	1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	161	Total	O	0	0
			161	161		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	79.89Å 112.40Å 131.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.20 – 1.80 38.53 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.6 (36.20-1.80) 97.7 (38.53-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.81Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.205 , 0.245 0.202 , 0.242	Depositor DCC
R_{free} test set	3239 reflections (6.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4048	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, MG, PYR, MES, K, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3839	0.76	2/5183 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	505	MET	CG-SD-CE	5.70	109.32	100.20
1	A	209	ASP	CB-CG-OD2	-5.59	113.27	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3782	0	3943	74	2
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	6	0	3	0	0
5	A	49	0	68	4	0
6	A	36	0	46	10	0
7	A	12	0	13	1	0
8	A	161	0	0	6	0
All	All	4048	0	4073	76	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:LYS:NZ	1:A:140:CYS:SG	2.24	1.10
1:A:267:LYS:NZ	8:A:701:HOH:O	1.93	1.01
1:A:442:ASP:OD1	1:A:458:LYS:NZ	2.08	0.86
1:A:127:LYS:HB3	1:A:178:ILE:HD13	1.59	0.84
1:A:109:ARG:HH22	6:A:615:GOL:H31	1.45	0.81
1:A:16:GLN:O	8:A:702:HOH:O	1.99	0.81
1:A:192:ASN:H	6:A:615:GOL:H11	1.47	0.79
1:A:240:ARG:NH1	6:A:616:GOL:O1	2.16	0.79
1:A:113:LEU:HD13	1:A:118:VAL:HG13	1.67	0.76
1:A:466:ASP:OD1	8:A:703:HOH:O	2.12	0.67
5:A:607:PEG:O1	5:A:607:PEG:O4	2.02	0.67
1:A:15:MET:SD	8:A:720:HOH:O	2.52	0.66
1:A:240:ARG:HD2	6:A:616:GOL:H2	1.77	0.66
1:A:171:GLU:HB3	1:A:178:ILE:HB	1.80	0.64
1:A:294:ALA:O	1:A:298:MET:HG3	1.96	0.64
1:A:138:GLU:H	1:A:138:GLU:CD	2.02	0.64
1:A:50:LYS:HE2	1:A:80:GLU:OE1	1.98	0.63
6:A:614:GOL:H31	8:A:762:HOH:O	1.99	0.63
1:A:245:PRO:O	1:A:248:ARG:HG2	1.98	0.63
1:A:240:ARG:HB2	6:A:616:GOL:H31	1.83	0.61
1:A:109:ARG:NH2	6:A:615:GOL:H31	2.15	0.60
1:A:137:ASP:OD2	1:A:139:THR:HG22	2.02	0.60
1:A:443:SER:HB2	5:A:604:PEG:H21	1.83	0.59
1:A:416:LEU:CD2	1:A:438:LEU:HD12	2.32	0.59
1:A:240:ARG:CD	6:A:616:GOL:H2	2.33	0.58
1:A:489:ILE:HD12	1:A:507:VAL:HG22	1.86	0.57
1:A:401:ALA:HB2	1:A:491:ILE:HD11	1.86	0.57
1:A:461:SER:HB2	1:A:463:GLN:NE2	2.19	0.57
1:A:150:GLN:N	1:A:150:GLN:OE1	2.37	0.56
1:A:170:LEU:HD11	1:A:180:GLU:HB2	1.88	0.56
1:A:257:GLU:HB3	1:A:278:ALA:HB3	1.88	0.55
1:A:146:LYS:NZ	1:A:147:LYS:HB2	2.22	0.55
1:A:113:LEU:HD21	1:A:141:ILE:HD13	1.90	0.54
1:A:35:ARG:HD3	7:A:617:MES:H62	1.89	0.54
1:A:175:ASP:OD1	1:A:175:ASP:N	2.40	0.54
1:A:127:LYS:HD2	1:A:127:LYS:C	2.28	0.53
1:A:114:LYS:HA	1:A:135:LEU:HD23	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:CYS:HB3	1:A:179:THR:HB	1.92	0.51
1:A:491:ILE:CD1	1:A:505:MET:HG3	2.41	0.50
1:A:31:ASN:N	1:A:31:ASN:OD1	2.45	0.50
1:A:149:PRO:HB3	1:A:173:HIS:O	2.12	0.49
1:A:283:GLY:O	1:A:286:ILE:O	2.30	0.49
1:A:394:GLN:HG3	1:A:502:THR:HB	1.94	0.48
1:A:101:LEU:HD13	1:A:217:ALA:HB2	1.95	0.48
1:A:171:GLU:CB	1:A:178:ILE:HB	2.42	0.47
1:A:477:LYS:HD3	1:A:510:ILE:HG21	1.96	0.46
1:A:377:LYS:HD2	1:A:406:GLU:CG	2.46	0.46
1:A:223:ASN:HD21	5:A:604:PEG:H42	1.80	0.46
1:A:240:ARG:HD2	6:A:616:GOL:C2	2.45	0.46
1:A:30:ASN:OD1	1:A:32:VAL:HG22	2.16	0.45
1:A:421:TYR:CZ	1:A:425:LEU:HD11	2.52	0.45
1:A:127:LYS:CB	1:A:178:ILE:HD13	2.40	0.45
1:A:134:PHE:HE2	1:A:136:GLY:HA2	1.81	0.45
1:A:354:LYS:N	1:A:354:LYS:HD3	2.31	0.45
1:A:158:ILE:HG12	1:A:195:LEU:CD2	2.46	0.45
1:A:148:LEU:O	1:A:152:VAL:HG22	2.16	0.45
1:A:233:ALA:HB3	5:A:610:PEG:H11	1.99	0.45
1:A:112:PHE:CZ	1:A:116:LYS:HG2	2.53	0.44
1:A:119:HIS:N	1:A:119:HIS:CD2	2.85	0.44
1:A:112:PHE:HZ	1:A:116:LYS:HG2	1.83	0.43
1:A:401:ALA:HB2	1:A:491:ILE:CD1	2.48	0.43
1:A:35:ARG:O	6:A:614:GOL:H32	2.18	0.43
1:A:112:PHE:CG	1:A:113:LEU:N	2.86	0.42
1:A:409:GLN:NE2	8:A:718:HOH:O	2.52	0.42
1:A:317:GLU:O	1:A:320:THR:HG23	2.19	0.42
1:A:161:ALA:HB3	1:A:164:SER:HG	1.85	0.42
1:A:492:HIS:CE1	1:A:504:LEU:HD13	2.55	0.41
1:A:128:LEU:HA	1:A:141:ILE:O	2.20	0.41
1:A:121:LYS:O	1:A:124:SER:HB3	2.20	0.41
1:A:169:VAL:HA	1:A:179:THR:HG22	2.01	0.41
1:A:218:ILE:HB	1:A:219:PRO:HD3	2.03	0.41
1:A:416:LEU:HD21	1:A:438:LEU:HD12	2.03	0.41
1:A:101:LEU:HD23	1:A:101:LEU:C	2.40	0.41
1:A:148:LEU:HB3	1:A:149:PRO:HD3	2.03	0.41
1:A:491:ILE:HD11	1:A:505:MET:HG3	2.02	0.40
1:A:205:ILE:HD12	1:A:238:LEU:HD23	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:GLU:OE1	1:A:463:GLN:NE2[7_545]	1.87	0.33
1:A:403:GLU:OE2	1:A:403:GLU:OE2[3_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	495/519 (95%)	481 (97%)	13 (3%)	1 (0%)	47 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	429/446 (96%)	423 (99%)	6 (1%)	67 59

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	SER
1	A	127	LYS
1	A	190	ARG
1	A	199	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	248	ARG
1	A	304	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	HIS
1	A	463	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	612	-	5,5,5	1.43	1 (20%)	5,5,5	0.97	0
5	PEG	A	608	-	6,6,6	0.76	0	5,5,5	0.59	0
5	PEG	A	611	-	6,6,6	1.05	0	5,5,5	0.64	0
6	GOL	A	614	-	5,5,5	1.03	0	5,5,5	1.08	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PYR	A	603	2	5,5,5	1.34	0	3,6,6	2.11	2 (66%)
6	GOL	A	606	-	5,5,5	0.75	0	5,5,5	1.31	0
7	MES	A	617	-	12,12,12	2.19	1 (8%)	14,16,16	1.74	3 (21%)
6	GOL	A	613	-	5,5,5	0.75	0	5,5,5	0.92	0
6	GOL	A	615	-	5,5,5	1.88	1 (20%)	5,5,5	1.11	0
5	PEG	A	604	-	6,6,6	0.76	0	5,5,5	0.69	0
5	PEG	A	605	-	6,6,6	0.77	0	5,5,5	0.34	0
5	PEG	A	607	-	6,6,6	0.76	0	5,5,5	1.03	0
6	GOL	A	616	-	5,5,5	0.79	0	5,5,5	1.07	0
5	PEG	A	609	-	6,6,6	0.69	0	5,5,5	0.32	0
5	PEG	A	610	-	6,6,6	0.87	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	612	-	-	1/4/4/4	-
5	PEG	A	608	-	-	4/4/4/4	-
5	PEG	A	611	-	-	0/4/4/4	-
6	GOL	A	614	-	-	1/4/4/4	-
4	PYR	A	603	2	-	0/4/4/4	-
6	GOL	A	606	-	-	0/4/4/4	-
7	MES	A	617	-	-	6/6/14/14	0/1/1/1
6	GOL	A	613	-	-	4/4/4/4	-
6	GOL	A	615	-	-	4/4/4/4	-
5	PEG	A	604	-	-	2/4/4/4	-
5	PEG	A	605	-	-	1/4/4/4	-
5	PEG	A	607	-	-	1/4/4/4	-
6	GOL	A	616	-	-	2/4/4/4	-
5	PEG	A	609	-	-	2/4/4/4	-
5	PEG	A	610	-	-	3/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	617	MES	C8-S	-7.27	1.67	1.77
6	A	615	GOL	O2-C2	-3.09	1.34	1.43
6	A	612	GOL	C1-C2	2.58	1.62	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	617	MES	C5-N4-C3	4.03	117.89	108.83
7	A	617	MES	O3S-S-C8	3.41	111.28	105.77
4	A	603	PYR	O2-C1-C2	2.98	122.11	113.97
7	A	617	MES	O2S-S-C8	2.36	109.76	106.92
6	A	614	GOL	C3-C2-C1	-2.22	103.09	111.70
4	A	603	PYR	O2-C1-O1	-2.11	118.77	123.61

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	613	GOL	C1-C2-C3-O3
6	A	613	GOL	O2-C2-C3-O3
6	A	615	GOL	O1-C1-C2-C3
6	A	616	GOL	O1-C1-C2-C3
5	A	609	PEG	O1-C1-C2-O2
5	A	610	PEG	O1-C1-C2-O2
7	A	617	MES	C7-C8-S-O3S
5	A	608	PEG	O1-C1-C2-O2
6	A	613	GOL	O1-C1-C2-C3
6	A	615	GOL	O1-C1-C2-O2
6	A	616	GOL	O1-C1-C2-O2
5	A	608	PEG	O2-C3-C4-O4
5	A	605	PEG	O2-C3-C4-O4
7	A	617	MES	C8-C7-N4-C3
7	A	617	MES	C8-C7-N4-C5
6	A	615	GOL	O2-C2-C3-O3
5	A	609	PEG	C4-C3-O2-C2
5	A	604	PEG	C1-C2-O2-C3
5	A	610	PEG	C1-C2-O2-C3
7	A	617	MES	C7-C8-S-O1S
7	A	617	MES	C7-C8-S-O2S
5	A	608	PEG	C1-C2-O2-C3
6	A	612	GOL	O1-C1-C2-O2
6	A	613	GOL	O1-C1-C2-O2
5	A	604	PEG	O1-C1-C2-O2
5	A	607	PEG	C4-C3-O2-C2
6	A	614	GOL	C1-C2-C3-O3
5	A	608	PEG	C4-C3-O2-C2
6	A	615	GOL	C1-C2-C3-O3
5	A	610	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	617	MES	N4-C7-C8-S

There are no ring outliers.

7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	614	GOL	2	0
7	A	617	MES	1	0
6	A	615	GOL	3	0
5	A	604	PEG	2	0
5	A	607	PEG	1	0
6	A	616	GOL	5	0
5	A	610	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	495/519 (95%)	0.76	73 (14%) 2 1	22, 35, 76, 95	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	134	PHE	8.3
1	A	177	VAL	7.0
1	A	11	ALA	7.0
1	A	128	LEU	6.5
1	A	170	LEU	6.1
1	A	112	PHE	5.8
1	A	141	ILE	5.6
1	A	115	ASN	5.5
1	A	140	CYS	5.2
1	A	169	VAL	5.2
1	A	152	VAL	5.1
1	A	512	LEU	5.0
1	A	502	THR	4.9
1	A	114	LYS	4.8
1	A	113	LEU	4.7
1	A	172	THR	4.4
1	A	119	HIS	4.4
1	A	130	THR	4.3
1	A	156	ASN	4.2
1	A	125	LYS	4.2
1	A	176	HIS	4.2
1	A	153	LYS	4.1
1	A	111	GLY	4.0
1	A	123	GLY	4.0
1	A	135	LEU	4.0
1	A	126	LEU	3.9
1	A	139	THR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	171	GLU	3.8
1	A	178	ILE	3.7
1	A	129	VAL	3.7
1	A	142	ALA	3.7
1	A	389	THR	3.6
1	A	116	LYS	3.5
1	A	388	GLU	3.3
1	A	148	LEU	3.3
1	A	181	VAL	3.3
1	A	155	GLY	3.2
1	A	182	LEU	3.2
1	A	118	VAL	3.2
1	A	179	THR	3.2
1	A	132	TYR	3.1
1	A	157	ILE	3.0
1	A	180	GLU	2.9
1	A	127	LYS	2.9
1	A	120	LEU	2.9
1	A	124	SER	2.9
1	A	189	GLU	2.9
1	A	146	LYS	2.9
1	A	121	LYS	2.8
1	A	467	ILE	2.8
1	A	137	ASP	2.8
1	A	13	ALA	2.8
1	A	151	SER	2.7
1	A	461	SER	2.6
1	A	168	LYS	2.6
1	A	173	HIS	2.6
1	A	463	GLN	2.5
1	A	117	GLU	2.5
1	A	310	ILE	2.4
1	A	154	PRO	2.4
1	A	343	CYS	2.4
1	A	122	GLU	2.3
1	A	253	ILE	2.3
1	A	150	GLN	2.2
1	A	149	PRO	2.2
1	A	186	VAL	2.2
1	A	199	LYS	2.2
1	A	80	GLU	2.2
1	A	390	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	311	THR	2.1
1	A	462	PHE	2.1
1	A	312	ALA	2.0
1	A	460	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	GOL	A	612	6/6	0.66	0.28	51,61,67,67	0
5	PEG	A	610	7/7	0.78	0.34	56,58,74,76	0
5	PEG	A	604	7/7	0.81	0.22	47,65,70,84	0
5	PEG	A	607	7/7	0.84	0.20	40,51,64,72	0
6	GOL	A	613	6/6	0.85	0.29	41,60,64,72	0
6	GOL	A	616	6/6	0.86	0.23	35,51,60,74	0
5	PEG	A	611	7/7	0.87	0.12	35,39,44,48	0
5	PEG	A	608	7/7	0.88	0.11	42,45,62,67	0
6	GOL	A	606	6/6	0.89	0.16	41,42,51,53	0
5	PEG	A	605	7/7	0.90	0.21	47,57,65,72	0
6	GOL	A	614	6/6	0.91	0.15	33,51,54,55	0
6	GOL	A	615	6/6	0.92	0.17	38,43,48,60	0
5	PEG	A	609	7/7	0.93	0.10	40,44,51,58	0
4	PYR	A	603	6/6	0.95	0.12	32,35,38,45	0
2	MG	A	601	1/1	0.95	0.05	36,36,36,36	0
7	MES	A	617	12/12	0.96	0.13	38,51,62,67	0
3	K	A	602	1/1	0.97	0.08	34,34,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around MG A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K A 602:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.