



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2021 – 02:35 PM EDT

PDB ID : 2Z5Y
Title : Crystal Structure of Human Monoamine Oxidase A (G110A) with Harmine
Authors : Son, S.Y.; Ma, J.; Yoshimura, M.; Tsukihara, T.
Deposited on : 2007-07-20
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	

2 Entry composition ⓘ

There are 5 unique types of molecules in this entry. The entry contains 4397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amine oxidase [flavin-containing] A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	513	4101	2635	699	746	21	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	GLY	engineered mutation	UNP P21397

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is 7-METHOXY-1-METHYL-9H-BETA-CARBOLINE (three-letter code: HRM)

HRM

The chemical structure of HRM (Hexahydroindole(1,2-b)pyridine) is shown. It consists of a benzene ring fused to a five-membered ring, which is further fused to a six-membered ring containing a nitrogen atom. The structure is substituted with 12 groups, labeled CAA through CAC, in green text. The substituents are: CAA (top left), OAJ (bottom left), CAG (top left), CAM (top left), CAG (top left), CAP (top left), CAB (top left), CAL (top left), CAH (top left), CAN (top left), CAO (top left), and CAC (top left).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			16	13	2	1		

- # DCX
-
- Chemical structure diagram of DCX (Dichlorophosphate) showing a central phosphorus atom (P1) bonded to two chlorine atoms (C2, C3) and two oxygen atoms (O4, O5). The structure is shown in a perspective view with a zigzag chain of atoms labeled C1 through C14.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			14	12	1	1		



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			14	12	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	199	Total	O	0	0
			199	199		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amine oxidase [flavin-containing] A



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	135.55Å 217.36Å 54.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.24 – 2.17 44.24 – 2.17	Depositor EDS
% Data completeness (in resolution range)	72.6 (44.24-2.17) 72.6 (44.24-2.17)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.3.0021	Depositor
R, R_{free}	0.193 , 0.244 0.191 , 0.242	Depositor DCC
R_{free} test set	1608 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4397	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HRM, FAD, DCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	3/4203 (0.1%)	0.92	8/5698 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	228	GLU	CG-CD	5.97	1.60	1.51
1	A	166	TRP	CB-CG	5.62	1.60	1.50
1	A	124	TYR	CE2-CZ	5.40	1.45	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	ARG	NE-CZ-NH1	-9.54	115.53	120.30
1	A	45	ARG	NE-CZ-NH2	9.21	124.90	120.30
1	A	81	SER	CB-CA-C	-7.55	95.75	110.10
1	A	454	ARG	NE-CZ-NH2	-5.81	117.39	120.30
1	A	229	ARG	NE-CZ-NH1	5.79	123.19	120.30
1	A	229	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	A	81	SER	N-CA-CB	-5.44	102.34	110.50
1	A	343	ASP	CB-CG-OD2	5.06	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4101	0	4097	67	0
2	A	53	0	29	2	0
3	A	16	0	12	3	0
4	A	28	0	54	1	0
5	A	199	0	0	2	0
All	All	4397	0	4192	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ASN:HD22	1:A:129:ARG:HH21	1.33	0.77
1:A:260:ASN:OD1	1:A:262:GLU:OE1	2.03	0.75
1:A:240:LEU:O	1:A:242:HIS:HD2	1.72	0.73
1:A:81:SER:OG	1:A:219:PHE:CE2	2.43	0.71
1:A:454:ARG:HD2	1:A:472:TRP:CH2	2.28	0.69
1:A:14:PHE:O	1:A:266:CYS:HA	1.97	0.65
1:A:424:ARG:HB3	1:A:437:THR:HB	1.79	0.63
1:A:352:PHE:HE2	3:A:700:HRM:HAE	1.67	0.60
1:A:397:TRP:CD1	2:A:600:FAD:HM71	2.38	0.59
1:A:352:PHE:HE2	3:A:700:HRM:CAE	2.15	0.58
1:A:507:PHE:O	1:A:511:VAL:HG22	2.02	0.58
1:A:406:CYS:HB3	1:A:407:TYR:HA	1.84	0.58
1:A:454:ARG:HD2	1:A:472:TRP:CZ2	2.40	0.57
1:A:156:THR:HG22	1:A:413:PRO:HG2	1.88	0.56
1:A:168:LYS:O	1:A:172:ARG:HD3	2.05	0.56
1:A:190:SER:HB2	1:A:413:PRO:HA	1.88	0.56
1:A:378:ALA:HB2	1:A:387:LEU:HD13	1.87	0.56
1:A:294:LEU:HB2	1:A:419:TYR:HB3	1.88	0.54
1:A:257:GLU:OE2	1:A:261:HIS:HD2	1.92	0.53
1:A:406:CYS:CB	1:A:407:TYR:HA	2.41	0.51
1:A:193:TRP:CG	1:A:411:PHE:HB2	2.45	0.51
1:A:491:TRP:HB2	4:A:1:DCX:O4	2.11	0.51
1:A:57:ASN:HD22	1:A:59:HIS:H	1.59	0.50
1:A:26:LEU:HB3	1:A:230:ILE:HG21	1.96	0.48
1:A:260:ASN:CG	1:A:262:GLU:OE1	2.52	0.48
1:A:56:ARG:NH2	1:A:228:GLU:OE1	2.47	0.47
1:A:165:CYS:HB2	1:A:171:ARG:HG3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:VAL:HG21	1:A:215:GLN:O	2.14	0.46
1:A:260:ASN:HD22	1:A:260:ASN:C	2.17	0.46
1:A:457:ARG:NH1	5:A:769:HOH:O	2.36	0.46
1:A:369:ARG:NH2	1:A:394:GLU:OE1	2.36	0.46
1:A:378:ALA:HA	1:A:383:SER:O	2.15	0.46
1:A:510:SER:O	1:A:513:ALA:HB3	2.14	0.46
1:A:117:ASN:HB3	1:A:120:ALA:HB3	1.98	0.46
1:A:357:LYS:O	1:A:361:LEU:HD12	2.15	0.46
1:A:101:VAL:HG13	1:A:327:GLU:HB3	1.98	0.46
1:A:246:HIS:HB2	1:A:257:GLU:HB3	1.98	0.46
1:A:335:ILE:HD11	3:A:700:HRM:OAJ	2.15	0.45
1:A:518:LEU:HD22	1:A:523:LEU:HB2	1.97	0.45
1:A:376:LEU:O	1:A:380:VAL:HG23	2.16	0.45
1:A:132:ASP:OD1	1:A:204:THR:HG23	2.17	0.45
1:A:260:ASN:ND2	1:A:262:GLU:OE1	2.50	0.45
1:A:70:VAL:HG11	1:A:219:PHE:CZ	2.51	0.45
1:A:16:VAL:HB	1:A:39:VAL:HG13	1.98	0.45
1:A:16:VAL:HG21	1:A:32:LEU:HD13	1.99	0.45
1:A:441:TRP:O	1:A:444:TYR:HB2	2.17	0.45
1:A:81:SER:HG	1:A:219:PHE:HE2	1.56	0.44
1:A:126:ASN:HD22	1:A:129:ARG:NH2	2.10	0.44
1:A:185:GLU:HA	1:A:185:GLU:OE1	2.18	0.43
1:A:57:ASN:HD22	1:A:59:HIS:N	2.16	0.43
1:A:329:GLU:HG3	5:A:863:HOH:O	2.17	0.43
1:A:72:PRO:HB3	1:A:483:ALA:HA	2.00	0.43
1:A:128:TRP:CZ2	1:A:173:PHE:CD2	3.07	0.43
2:A:600:FAD:N1	2:A:600:FAD:H2'	2.34	0.43
1:A:69:TYR:HA	1:A:217:ARG:O	2.18	0.43
1:A:191:ALA:O	1:A:195:LEU:HG	2.19	0.42
1:A:284:ARG:HA	1:A:285:PRO:C	2.39	0.42
1:A:254:ILE:O	1:A:265:GLU:HA	2.19	0.42
1:A:150:ASP:HA	1:A:417:THR:HG21	2.01	0.42
1:A:331:ALA:HA	1:A:332:PRO:HD2	1.93	0.42
1:A:391:HIS:HE1	1:A:393:GLU:OE1	2.02	0.42
1:A:245:THR:OG1	1:A:246:HIS:HD2	2.03	0.41
1:A:391:HIS:CE1	1:A:393:GLU:HG2	2.55	0.41
1:A:218:LYS:HE3	1:A:218:LYS:HB2	1.87	0.41
1:A:183:THR:HG22	1:A:354:LEU:HD22	2.03	0.41
1:A:234:LEU:HD23	1:A:234:LEU:HA	1.85	0.40
1:A:78:LEU:HD23	1:A:78:LEU:HA	1.93	0.40
1:A:236:ASP:OD1	1:A:237:GLN:NE2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	511/513 (100%)	489 (96%)	21 (4%)	1 (0%)	47 52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	439/439 (100%)	417 (95%)	22 (5%)	24 27

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	MET
1	A	57	ASN
1	A	81	SER
1	A	101	VAL
1	A	164	ILE
1	A	169	THR
1	A	172	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	184	SER
1	A	251	SER
1	A	260	ASN
1	A	267	LYS
1	A	334	SER
1	A	361	LEU
1	A	375	GLU
1	A	387	LEU
1	A	423	ILE
1	A	466	VAL
1	A	474	GLN
1	A	485	GLU
1	A	514	LEU
1	A	516	PHE
1	A	517	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	126	ASN
1	A	242	HIS
1	A	246	HIS
1	A	260	ASN
1	A	261	HIS
1	A	292	ASN
1	A	296	GLN
1	A	391	HIS
1	A	418	GLN
1	A	461	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	600	-	51,58,58	1.96	12 (23%)	60,89,89	2.08	16 (26%)
4	DCX	A	1	-	12,13,13	0.52	0	13,15,15	0.47	0
4	DCX	A	2	-	12,13,13	0.48	0	13,15,15	0.86	1 (7%)
3	HRM	A	700	-	18,18,18	1.77	6 (33%)	21,26,26	1.60	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	600	-	-	3/30/50/50	0/6/6/6
4	DCX	A	1	-	-	9/11/11/11	-
4	DCX	A	2	-	-	4/11/11/11	-
3	HRM	A	700	-	-	0/2/2/2	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	FAD	C4X-N5	6.11	1.42	1.33
2	A	600	FAD	O4B-C1B	5.87	1.49	1.41
2	A	600	FAD	C10-N1	4.19	1.38	1.33
2	A	600	FAD	C4-N3	4.13	1.40	1.33
2	A	600	FAD	C5'-C4'	3.66	1.57	1.51
2	A	600	FAD	C4-C4X	-3.05	1.36	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	FAD	C2B-C1B	-2.86	1.49	1.53
3	A	700	HRM	CAN-CAO	-2.75	1.37	1.45
3	A	700	HRM	CAL-CAP	-2.51	1.39	1.43
2	A	600	FAD	C7M-C7	2.50	1.56	1.51
2	A	600	FAD	C4'-C3'	-2.31	1.49	1.53
3	A	700	HRM	CAL-NAH	2.31	1.34	1.32
2	A	600	FAD	C2B-C3B	2.29	1.59	1.53
2	A	600	FAD	C1'-N10	-2.27	1.45	1.48
3	A	700	HRM	CAB-CAL	2.24	1.52	1.50
2	A	600	FAD	C9A-N10	2.23	1.41	1.38
3	A	700	HRM	CAG-CAK	2.21	1.41	1.37
3	A	700	HRM	CAE-CAC	2.18	1.39	1.36

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	FAD	C4-N3-C2	7.33	121.33	115.14
2	A	600	FAD	C4-C4X-N5	4.83	124.12	118.60
2	A	600	FAD	N3A-C2A-N1A	-4.69	121.35	128.68
2	A	600	FAD	C4-C4X-C10	-4.60	116.91	119.95
2	A	600	FAD	O4B-C1B-C2B	-3.74	101.47	106.93
2	A	600	FAD	C5'-C4'-C3'	-3.55	105.35	112.20
3	A	700	HRM	CAE-CAC-NAH	-3.49	120.37	123.81
3	A	700	HRM	CAA-OAJ-CAK	-3.48	109.95	117.51
2	A	600	FAD	C10-C4X-N5	-3.42	118.89	121.26
3	A	700	HRM	CAC-NAH-CAL	3.33	122.37	118.33
2	A	600	FAD	C1'-N10-C10	2.95	121.05	118.41
2	A	600	FAD	C6-C5X-C9A	2.87	122.81	119.05
2	A	600	FAD	C9A-N10-C10	-2.82	118.22	121.91
2	A	600	FAD	C9A-C5X-N5	-2.78	118.01	122.36
3	A	700	HRM	CAF-CAN-CAM	2.74	121.81	118.17
2	A	600	FAD	C5X-C9A-N10	2.60	119.60	117.72
2	A	600	FAD	C2A-N1A-C6A	2.60	123.19	118.75
4	A	2	DCX	C2-P1-C3	-2.42	103.15	106.47
2	A	600	FAD	O2A-PA-O1A	2.19	123.09	112.24
2	A	600	FAD	C5B-C4B-C3B	-2.14	107.14	115.18
2	A	600	FAD	C4X-C4-N3	-2.10	120.57	123.43

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	600	FAD	C5B-O5B-PA-O1A
4	A	1	DCX	C6-C4-P1-O4
4	A	1	DCX	C6-C4-P1-C2
4	A	2	DCX	C10-C11-C12-C13
4	A	1	DCX	P1-C4-C6-C7
4	A	1	DCX	C11-C12-C13-C14
4	A	2	DCX	C7-C8-C9-C10
4	A	2	DCX	C6-C7-C8-C9
4	A	1	DCX	C6-C4-P1-C3
4	A	1	DCX	C10-C11-C12-C13
4	A	1	DCX	C4-C6-C7-C8
4	A	1	DCX	C7-C8-C9-C10
4	A	1	DCX	C9-C10-C11-C12
2	A	600	FAD	O4B-C4B-C5B-O5B
2	A	600	FAD	C5B-O5B-PA-O3P
4	A	2	DCX	C4-C6-C7-C8

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	FAD	2	0
4	A	1	DCX	1	0
3	A	700	HRM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	513/513 (100%)	-0.18	10 (1%) 66 67	21, 34, 53, 100	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	518	LEU	5.2
1	A	521	TYR	4.7
1	A	516	PHE	3.9
1	A	12	HIS	3.4
1	A	515	GLY	3.0
1	A	513	ALA	3.0
1	A	512	THR	2.7
1	A	523	LEU	2.6
1	A	166	TRP	2.5
1	A	252	ASP	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	DCX	A	1	14/14	0.87	0.14	74,76,85,86	0
4	DCX	A	2	14/14	0.90	0.16	64,69,83,84	0
2	FAD	A	600	53/53	0.98	0.13	14,21,27,29	0
3	HRM	A	700	16/16	0.98	0.12	29,31,32,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.