



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 10:14 AM BST

PDB ID : 2Z73
Title : Crystal structure of squid rhodopsin
Authors : Murakami, M.; Kouyama, T.
Deposited on : 2007-08-13
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>4%</div> <div> <div></div> <div>50%</div> <div>27%</div> <div>•</div> <div>22%</div> </div> </div>
1	B	448	<div> <div>52%</div> <div>22%</div> <div>•</div> <div>23%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PLM	B	1002	-	-	-	X

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 5835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	0	0
			2780	1838	443	473	26			
1	B	347	Total	C	N	O	S	0	0	0
			2762	1828	440	468	26			

- Molecule 2 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: $C_{14}H_{28}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	14	6		
2	B	1	Total	C	O	0	0
			20	14	6		

- Molecule 3 is RETINAL (three-letter code: RET) (formula: $C_{20}H_{28}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C		0	0
			20	20			
3	B	1	Total	C		0	0
			20	20			

- Molecule 4 is PALMITIC ACID (three-letter code: PLM) (formula: C₁₆H₃₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			17	16	1		
4	A	1	Total	C	O	0	0
			17	16	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			17	16	1		
4	B	1	Total	C	O	0	0
			17	16	1		

- Molecule 5 is DOCOSANE (three-letter code: TWT) (formula: $C_{22}H_{46}$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C	0	0
			22	22		
5	B	1	Total	C	0	0
			22	22		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	O	P	0	0
			39	30	8	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	28	Total 28	O 28	0	0
8	B	29	Total 29	O 29	0	0

TYR
PRO
GLN
GLY
TYR
PRO
PRO
GLN
GLY
TYR
PRO
PRO
GLN
GLY
TYR
PRO
PRO
GLN
GLY
TYR
PRO
PRO
PRO
PRO
GLN
GLY
ALA
ALA
PRO
GLN
GLY
ALA
PRO
PRO
ALA
ALA
PRO
PRO
GLN
GLY
VAL
ASP
ASN
GLN
ALA
TYR
GLN
ALA

4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	122.55Å 122.55Å 158.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.50 57.16 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.2 (15.00-2.50) 93.3 (57.16-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.51Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.188 , 0.206 0.193 , 0.210	Depositor DCC
R_{free} test set	2174 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	54.1	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 73.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.107 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5835	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PC1, RET, SO4, TWT, PLM, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2865	0.57	0/3889
1	B	0.38	0/2847	0.57	0/3865
All	All	0.38	0/5712	0.57	0/7754

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2780	0	2765	90	0
1	B	2762	0	2751	86	0
2	A	20	0	28	4	0
2	B	20	0	28	2	0
3	A	20	0	27	2	0
3	B	20	0	27	2	0
4	A	34	0	62	1	0
4	B	34	0	62	1	0
5	A	22	0	46	0	0
5	B	22	0	46	0	0
6	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	39	0	49	1	0
8	A	28	0	0	4	0
8	B	29	0	0	1	0
All	All	5835	0	5891	180	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 180 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:150:ARG:H	1:B:150:ARG:HD2	1.28	0.96
1:B:354:ILE:HD13	1:B:354:ILE:H	1.30	0.94
1:A:21:PRO:HA	1:A:24:ARG:HD3	1.52	0.92
1:A:284:ALA:HB2	1:A:292:VAL:HG21	1.54	0.89
1:B:148:SER:HB2	1:B:150:ARG:HH21	1.43	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/448 (78%)	316 (91%)	30 (9%)	2 (1%)	25	43
1	B	345/448 (77%)	318 (92%)	22 (6%)	5 (1%)	11	20
All	All	693/896 (77%)	634 (92%)	52 (8%)	7 (1%)	15	28

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	226	SER
1	B	138	GLY
1	B	352	THR
1	B	223	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/369 (80%)	282 (95%)	14 (5%)	26	49
1	B	295/369 (80%)	275 (93%)	20 (7%)	16	30
All	All	591/738 (80%)	557 (94%)	34 (6%)	20	38

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	63	LYS
1	B	102	ILE
1	B	346	ASP
1	B	76	LEU
1	A	105	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	269	GLN
1	A	328	GLN
1	B	242	ASN
1	A	254	ASN
1	B	222	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	RET	B	1000	1	20,20,21	1.81	6 (30%)	27,27,28	1.69	6 (22%)
6	SO4	B	1006	-	4,4,4	0.25	0	6,6,6	0.12	0
3	RET	A	1000	1	20,20,21	1.84	5 (25%)	27,27,28	1.70	7 (25%)
4	PLM	A	1001	-	16,16,17	0.32	0	15,15,17	0.66	0
7	PC1	B	1004	-	38,38,53	1.64	2 (5%)	41,43,61	0.97	1 (2%)
2	BOG	B	1005	-	20,20,20	1.85	8 (40%)	25,25,25	2.86	11 (44%)
2	BOG	A	1005	-	20,20,20	1.85	8 (40%)	25,25,25	2.83	11 (44%)
5	TWT	A	1003	-	21,21,21	0.29	0	20,20,20	1.65	4 (20%)
5	TWT	B	1003	-	21,21,21	0.26	0	20,20,20	1.73	6 (30%)
4	PLM	B	1001	-	16,16,17	0.33	0	15,15,17	0.67	0
4	PLM	B	1002	1	16,16,17	0.32	0	15,15,17	0.66	0
4	PLM	A	1002	1	16,16,17	0.30	0	15,15,17	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RET	B	1000	1	-	1/13/30/31	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PC1	B	1004	-	-	14/42/42/57	-
3	RET	A	1000	1	-	1/13/30/31	0/1/1/1
4	PLM	A	1001	-	-	5/13/14/15	-
2	BOG	B	1005	-	-	6/11/31/31	0/1/1/1
2	BOG	A	1005	-	-	5/11/31/31	0/1/1/1
5	TWT	A	1003	-	-	9/19/19/19	-
5	TWT	B	1003	-	-	10/19/19/19	-
4	PLM	B	1001	-	-	7/13/14/15	-
4	PLM	B	1002	1	-	6/13/14/15	-
4	PLM	A	1002	1	-	6/13/14/15	-

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	1004	PC1	O21-C21	7.39	1.55	1.34
3	A	1000	RET	C14-C13	4.44	1.37	1.33
3	B	1000	RET	C14-C13	4.41	1.37	1.33
7	B	1004	PC1	O31-C31	4.39	1.46	1.33
2	A	1005	BOG	O5-C1	3.82	1.51	1.41

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1005	BOG	C1'-O1-C1	-8.63	99.54	113.84
2	A	1005	BOG	C1'-O1-C1	-8.33	100.03	113.84
2	A	1005	BOG	O1-C1-C2	-4.90	100.66	108.30
2	B	1005	BOG	O1-C1-C2	-4.80	100.81	108.30
2	B	1005	BOG	C1-C2-C3	-4.39	100.85	110.00

There are no chirality outliers.

5 of 70 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	B	1004	PC1	C1-O11-P-O12
7	B	1004	PC1	C1-O11-P-O14
7	B	1004	PC1	C32-C31-O31-C3
7	B	1004	PC1	O32-C31-O31-C3
2	A	1005	BOG	O1-C1'-C2'-C3'


There are no ring outliers.





7 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1000	RET	2	0
3	A	1000	RET	2	0
7	B	1004	PC1	1	0
2	B	1005	BOG	2	0
2	A	1005	BOG	4	0
4	B	1002	PLM	1	0
4	A	1002	PLM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/448 (78%)	0.15	16 (4%) 32 34	35, 55, 118, 140	0
1	B	347/448 (77%)	-0.02	1 (0%) 94 94	32, 49, 98, 121	0
All	All	697/896 (77%)	0.07	17 (2%) 59 62	32, 52, 108, 140	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	LEU	5.4
1	A	239	LYS	5.1
1	A	237	MET	4.5
1	A	234	MET	4.0
1	A	236	ALA	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PLM	B	1002	17/18	0.74	0.73	104,108,111,111	0
4	PLM	B	1001	17/18	0.82	0.38	53,70,74,74	0
4	PLM	A	1001	17/18	0.82	0.30	62,67,70,70	0
5	TWT	A	1003	22/22	0.86	0.69	57,93,102,102	0
2	BOG	B	1005	20/20	0.88	0.44	68,113,119,121	0
7	PC1	B	1004	39/54	0.89	0.18	42,67,120,120	0
5	TWT	B	1003	22/22	0.89	0.73	63,80,87,88	0
4	PLM	A	1002	17/18	0.89	0.32	96,102,110,110	0
2	BOG	A	1005	20/20	0.92	0.39	54,100,107,107	0
3	RET	A	1000	20/21	0.96	0.23	39,52,58,59	0
6	SO4	B	1006	5/5	0.96	0.12	96,96,96,97	0
3	RET	B	1000	20/21	0.97	0.23	33,38,40,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around PLM B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLM A 1001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TWT A 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around BOG B 1005:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PC1 B 1004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TWT B 1003:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLM A 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG A 1005:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around RET A 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RET B 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.