



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 06:10 pm BST

PDB ID : 5Z7J  
Title : Crystal structure of a lactonase double mutant in complex with ligand l  
Authors : Zheng, Y.Y.; Liu, W.D.; Chen, C.C.; Guo, R.T.  
Deposited on : 2018-01-29  
Resolution : 1.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>14%</div> </div> <div></div> </div>
1	B	264	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>15%</div> </div> <div></div> </div>
1	C	264	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>16%</div> </div> <div></div> </div>
1	D	264	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>19%</div> </div> <div></div> </div>
1	E	264	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>15%</div> </div> <div></div> </div>
1	F	264	<div> <div>11%</div> <div> <div></div> <div>83%</div> <div>15%</div> </div> <div></div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	264	<div><div></div><div>3%</div><div>84%</div><div>15%</div></div>
1	H	264	<div><div></div><div>10%</div><div>84%</div><div>14%</div><div></div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactonase for protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2048	1300	347	392	9			
1	B	263	Total	C	N	O	S	0	0	0
			2048	1300	347	392	9			
1	C	264	Total	C	N	O	S	0	0	0
			2053	1303	348	393	9			
1	D	264	Total	C	N	O	S	0	0	0
			2053	1303	348	393	9			
1	E	263	Total	C	N	O	S	0	0	0
			2043	1297	346	391	9			
1	F	263	Total	C	N	O	S	0	0	0
			2048	1300	347	392	9			
1	G	263	Total	C	N	O	S	0	0	0
			2043	1297	346	391	9			
1	H	263	Total	C	N	O	S	0	0	0
			2048	1300	347	392	9			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
A	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
B	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
B	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
C	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
C	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
D	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
D	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
E	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
E	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
F	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
F	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
G	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1
H	105	ALA	SER	engineered mutation	UNP A0A0D2ILK1
H	160	ALA	TYR	engineered mutation	UNP A0A0D2ILK1

- Molecule 2 is (3S,7R,11E)-7,14,16-trihydroxy-3-methyl-3,4,5,6,7,8,9,10-octahydro-1H-2-benzoxacyclotetradecin-1-one (three-letter code: 36J) (formula: C<sub>18</sub>H<sub>24</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			23	18	5		
2	B	1	Total	C	O	0	0
			23	18	5		
2	C	1	Total	C	O	0	0
			23	18	5		
2	D	1	Total	C	O	0	0
			23	18	5		
2	E	1	Total	C	O	0	0
			23	18	5		
2	F	1	Total	C	O	0	0
			23	18	5		
2	G	1	Total	C	O	0	0
			23	18	5		
2	H	1	Total	C	O	0	0
			23	18	5		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	286	Total	O	0	0
			286	286		
4	B	297	Total	O	0	0
			297	297		
4	C	226	Total	O	0	0
			226	226		
4	D	204	Total	O	0	0
			204	204		
4	E	217	Total	O	0	0
			217	217		
4	F	190	Total	O	0	0
			190	190		
4	G	268	Total	O	0	0
			268	268		
4	H	176	Total	O	0	0
			176	176		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lactonase for protein




- Molecule 1: Lactonase for protein




- Molecule 1: Lactonase for protein





- Molecule 1: Lactonase for protein









• Molecule 1: Lactonase for protein





• Molecule 1: Lactonase for protein



• Molecule 1: Lactonase for protein



• Molecule 1: Lactonase for protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.33Å 94.76Å 100.45Å 90.73° 92.36° 91.59°	Depositor
Resolution (Å)	24.71 – 1.98 24.71 – 1.98	Depositor EDS
% Data completeness (in resolution range)	97.3 (24.71-1.98) 92.0 (24.71-1.98)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 1.98Å)	Xtriage
Refinement program	PHENIX (1.12 _2829: ???)	Depositor
R, $R_{free}$	0.166 , 0.209 0.177 , 0.202	Depositor DCC
$R_{free}$ test set	2003 reflections (1.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l 0.016 for -h,k,-l 0.016 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	18446	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, 36J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	0/2106	0.56	0/2875
1	B	0.40	0/2106	0.57	0/2875
1	C	0.39	0/2111	0.56	0/2882
1	D	0.39	0/2111	0.55	0/2882
1	E	0.39	0/2101	0.56	0/2871
1	F	0.38	0/2106	0.54	0/2875
1	G	0.39	0/2101	0.56	0/2871
1	H	0.38	0/2106	0.55	0/2875
All	All	0.39	0/16848	0.56	0/23006

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2048	0	1989	25	0
1	B	2048	0	1989	29	0
1	C	2053	0	1994	41	0
1	D	2053	0	1994	36	0
1	E	2043	0	1981	36	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2048	0	1989	29	0
1	G	2043	0	1981	24	0
1	H	2048	0	1989	31	0
2	A	23	0	22	4	0
2	B	23	0	22	3	0
2	C	23	0	22	3	0
2	D	23	0	22	2	0
2	E	23	0	22	5	0
2	F	23	0	23	4	0
2	G	23	0	22	1	0
2	H	23	0	22	1	0
3	B	14	0	20	2	0
4	A	286	0	0	3	0
4	B	297	0	0	3	0
4	C	226	0	0	3	0
4	D	204	0	0	3	0
4	E	217	0	0	1	0
4	F	190	0	0	3	0
4	G	268	0	0	2	0
4	H	176	0	0	1	0
All	All	18446	0	16103	264	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 264 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:144:VAL:HB	1:C:148:THR:HG21	1.52	0.91
1:G:133:GLU:O	1:G:135:PRO:HD3	1.71	0.91
1:E:144:VAL:HB	1:E:148:THR:HG21	1.52	0.91
1:C:140:HIS:HB3	1:C:143:GLU:OE1	1.76	0.85
1:H:142:HIS:ND1	1:H:191:ARG:NH2	2.26	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/264 (99%)	254 (97%)	6 (2%)	1 (0%)	34	22
1	B	261/264 (99%)	251 (96%)	8 (3%)	2 (1%)	19	9
1	C	262/264 (99%)	255 (97%)	5 (2%)	2 (1%)	19	9
1	D	262/264 (99%)	253 (97%)	7 (3%)	2 (1%)	19	9
1	E	261/264 (99%)	250 (96%)	9 (3%)	2 (1%)	19	9
1	F	261/264 (99%)	250 (96%)	9 (3%)	2 (1%)	19	9
1	G	261/264 (99%)	250 (96%)	10 (4%)	1 (0%)	34	22
1	H	261/264 (99%)	252 (97%)	8 (3%)	1 (0%)	34	22
All	All	2090/2112 (99%)	2015 (96%)	62 (3%)	13 (1%)	25	14

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	163	ASN
1	E	163	ASN
1	B	189	TYR
1	C	189	TYR
1	D	189	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/222 (100%)	217 (98%)	5 (2%)	50	44

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	222/222 (100%)	218 (98%)	4 (2%)	59	51
1	C	222/222 (100%)	217 (98%)	5 (2%)	50	44
1	D	222/222 (100%)	217 (98%)	5 (2%)	50	44
1	E	221/222 (100%)	213 (96%)	8 (4%)	35	23
1	F	222/222 (100%)	212 (96%)	10 (4%)	27	15
1	G	221/222 (100%)	214 (97%)	7 (3%)	39	28
1	H	222/222 (100%)	215 (97%)	7 (3%)	39	28
All	All	1774/1776 (100%)	1723 (97%)	51 (3%)	42	31

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	225	ASN
1	F	5	ARG
1	H	95	ASP
1	E	246	TYR
1	F	12	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	36J	A	301	-	24,24,24	0.34	0	32,32,32	0.43	0
2	36J	G	301	-	24,24,24	0.34	0	32,32,32	0.46	0
2	36J	E	301	-	24,24,24	0.35	0	32,32,32	0.42	0
3	PEG	B	303	-	6,6,6	0.35	0	5,5,5	0.39	0
3	PEG	B	302	-	6,6,6	0.42	0	5,5,5	0.18	0
2	36J	D	301	-	24,24,24	0.34	0	32,32,32	0.41	0
2	36J	B	301	-	24,24,24	0.35	0	32,32,32	0.43	0
2	36J	H	301	-	24,24,24	0.33	0	32,32,32	0.53	0
2	36J	F	301	-	24,24,24	0.34	0	32,32,32	0.45	0
2	36J	C	301	-	24,24,24	0.34	0	32,32,32	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	36J	A	301	-	-	8/22/22/22	0/1/2/2
2	36J	G	301	-	-	13/22/22/22	0/1/2/2
2	36J	E	301	-	-	9/22/22/22	0/1/2/2
3	PEG	B	303	-	-	2/4/4/4	-
3	PEG	B	302	-	-	3/4/4/4	-
2	36J	D	301	-	-	13/22/22/22	0/1/2/2
2	36J	B	301	-	-	11/22/22/22	0/1/2/2
2	36J	H	301	-	-	7/22/22/22	0/1/2/2
2	36J	F	301	-	-	8/22/22/22	0/1/2/2
2	36J	C	301	-	-	7/22/22/22	0/1/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 81 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	301	36J	CAL-CAO-CAW-CAN
2	G	301	36J	CAL-CAO-CAW-OAE
2	G	301	36J	CAA-CAV-OAP-CAQ
2	G	301	36J	CAU-CAQ-OAP-CAV
2	G	301	36J	CAK-CAN-CAW-CAO

There are no ring outliers.

9 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	36J	4	0
2	G	301	36J	1	0
2	E	301	36J	5	0
3	B	303	PEG	2	0
2	D	301	36J	2	0
2	B	301	36J	3	0
2	H	301	36J	1	0
2	F	301	36J	4	0
2	C	301	36J	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## Ligand 36J G 301




Bond lengths



Bond angles




Torsions




Rings


## Ligand 36J E 301




## Ligand 36J D 301




## Ligand 36J B 301




Bond lengths



Bond angles




Torsions




Rings


## Ligand 36J H 301




Bond lengths



Bond angles





Torsions



Rings

## Ligand 36J F 301





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/264 (99%)	-0.09	6 (2%) 60 62	22, 29, 40, 58	0
1	B	263/264 (99%)	-0.10	6 (2%) 60 62	22, 28, 41, 56	0
1	C	264/264 (100%)	0.12	15 (5%) 23 25	23, 33, 55, 74	0
1	D	264/264 (100%)	0.20	16 (6%) 21 23	23, 34, 60, 76	0
1	E	263/264 (99%)	0.16	15 (5%) 23 25	25, 34, 50, 67	0
1	F	263/264 (99%)	0.40	28 (10%) 6 7	27, 38, 62, 77	0
1	G	263/264 (99%)	0.07	9 (3%) 45 48	24, 32, 45, 55	0
1	H	263/264 (99%)	0.45	27 (10%) 6 7	27, 38, 64, 83	0
All	All	2106/2112 (99%)	0.15	122 (5%) 23 25	22, 33, 53, 83	0

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	266	LYS	6.1
1	D	3	ALA	5.6
1	H	139	LEU	5.5
1	E	3	ALA	5.2
1	H	140	HIS	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.




## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PEG	B	302	7/7	0.73	0.35	32,35,37,44	0
2	36J	F	301	23/23	0.83	0.23	41,50,58,61	0
2	36J	B	301	23/23	0.84	0.19	27,37,45,48	0
2	36J	E	301	23/23	0.84	0.17	29,42,50,54	0
2	36J	H	301	23/23	0.85	0.20	41,49,54,57	0
2	36J	D	301	23/23	0.86	0.20	33,44,53,58	0
2	36J	C	301	23/23	0.86	0.19	29,43,50,55	0
2	36J	G	301	23/23	0.88	0.15	28,39,44,46	0
2	36J	A	301	23/23	0.90	0.15	29,40,45,49	0
3	PEG	B	303	7/7	0.91	0.20	32,35,37,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 36J B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 36J E 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 36J H 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 36J D 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 36J C 301:**



2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around 36J G 301:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 36J A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.