



Full wwPDB X-ray Structure Validation Report ⓘ

May 19, 2020 – 07:12 pm BST

PDB ID : 1Z82
Title : Crystal structure of glycerol-3-phosphate dehydrogenase (TM0378) from THERMOTOGA MARITIMA at 2.00 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2005-03-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>8% • 7%</div> </div> </div>
1	B	335	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>8% • 7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G3H	A	600	X	-	-	-
6	G3P	B	600	X	-	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycerol-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	Se	0	4	0
			2422	1548	415	445	3	11			
1	B	312	Total	C	N	O	S	Se	0	2	0
			2414	1543	413	444	3	11			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	LEADER SEQUENCE	GB 15644628
A	-10	GLY	-	LEADER SEQUENCE	GB 15644628
A	-9	SER	-	LEADER SEQUENCE	GB 15644628
A	-8	ASP	-	LEADER SEQUENCE	GB 15644628
A	-7	LYS	-	LEADER SEQUENCE	GB 15644628
A	-6	ILE	-	LEADER SEQUENCE	GB 15644628
A	-5	HIS	-	LEADER SEQUENCE	GB 15644628
A	-4	HIS	-	LEADER SEQUENCE	GB 15644628
A	-3	HIS	-	LEADER SEQUENCE	GB 15644628
A	-2	HIS	-	LEADER SEQUENCE	GB 15644628
A	-1	HIS	-	LEADER SEQUENCE	GB 15644628
A	0	HIS	-	LEADER SEQUENCE	GB 15644628
A	1	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	3	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	20	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	94	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	215	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	225	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	234	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	278	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	288	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	308	MSE	MET	MODIFIED RESIDUE	GB 15644628
A	312	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	-11	MET	-	LEADER SEQUENCE	GB 15644628
B	-10	GLY	-	LEADER SEQUENCE	GB 15644628

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	SER	-	LEADER SEQUENCE	GB 15644628
B	-8	ASP	-	LEADER SEQUENCE	GB 15644628
B	-7	LYS	-	LEADER SEQUENCE	GB 15644628
B	-6	ILE	-	LEADER SEQUENCE	GB 15644628
B	-5	HIS	-	LEADER SEQUENCE	GB 15644628
B	-4	HIS	-	LEADER SEQUENCE	GB 15644628
B	-3	HIS	-	LEADER SEQUENCE	GB 15644628
B	-2	HIS	-	LEADER SEQUENCE	GB 15644628
B	-1	HIS	-	LEADER SEQUENCE	GB 15644628
B	0	HIS	-	LEADER SEQUENCE	GB 15644628
B	1	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	3	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	20	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	94	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	215	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	225	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	234	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	278	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	288	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	308	MSE	MET	MODIFIED RESIDUE	GB 15644628
B	312	MSE	MET	MODIFIED RESIDUE	GB 15644628

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula: $C_3H_7O_6P$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is SN-GLYCEROL-3-PHOSPHATE (three-letter code: G3P) (formula: $C_3H_9O_6P$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	161	Total	O	0	0
			161	161		
7	B	162	Total	O	0	0
			162	162		

- Molecule 1: glycerol-3-phosphate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.09 Å 67.45 Å 75.60 Å 90.00° 113.67° 90.00°	Depositor
Resolution (Å)	29.81 – 2.00 29.81 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.3 (29.81-2.00) 98.4 (29.81-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.166 , 0.211 0.177 , 0.222	Depositor DCC
R_{free} test set	2024 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtriage
Anisotropy	0.804	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5339	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, G3P, MPD, NDP, G3H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/2470 (0.0%)	0.74	5/3321 (0.2%)
1	B	0.76	1/2453 (0.0%)	0.77	3/3298 (0.1%)
All	All	0.75	2/4923 (0.0%)	0.76	8/6619 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	166	CYS	CB-SG	-5.59	1.72	1.81
1	A	159	GLU	CB-CG	5.50	1.62	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	243	ARG	NE-CZ-NH2	-9.40	115.60	120.30
1	B	243	ARG	NE-CZ-NH1	7.55	124.08	120.30
1	B	109	ARG	NE-CZ-NH2	-6.85	116.88	120.30
1	A	87	ARG	NE-CZ-NH1	6.09	123.35	120.30
1	A	240	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	240	ARG	NE-CZ-NH2	-5.93	117.34	120.30
1	A	87	ARG	NE-CZ-NH2	-5.86	117.37	120.30
1	A	212	ARG	NE-CZ-NH2	-5.29	117.65	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2422	0	2452	25	0
1	B	2414	0	2438	21	0
2	A	48	0	26	1	0
2	B	48	0	26	1	0
3	A	10	0	5	1	0
4	A	8	0	14	0	0
4	B	32	0	56	1	0
5	A	24	0	42	5	0
6	B	10	0	7	1	0
7	A	161	0	0	2	0
7	B	162	0	0	4	0
All	All	5339	0	5066	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:MSE:SE	1:A:312:MSE:CE	2.15	1.44
1:A:308:MSE:HE1	1:A:312:MSE:SE	1.90	1.21
1:B:308:MSE:CE	1:B:312:MSE:SE	2.54	1.05
2:B:500:NDP:H8A	7:B:652:HOH:O	1.60	1.01
1:B:308:MSE:HE1	1:B:312:MSE:SE	2.16	0.96
1:A:308:MSE:CE	1:A:312:MSE:SE	2.67	0.92
1:B:308:MSE:HE3	1:B:312:MSE:SE	2.21	0.89
1:B:211:ALA:HB1	1:B:215:MSE:CE	2.12	0.79
1:B:211:ALA:HB1	1:B:215:MSE:HE3	1.65	0.79
2:A:500:NDP:H8A	7:A:664:HOH:O	1.89	0.72
1:A:222:LYS:HA	1:A:225[B]:MSE:HE3	1.76	0.68
1:B:44[A]:SER:OG	1:B:46:THR:HG22	1.97	0.64
1:B:215:MSE:HE2	1:B:221:GLN:HA	1.79	0.64
1:A:258:LEU:C	1:A:258:LEU:HD23	2.24	0.58
1:A:44:SER:O	1:A:46:THR:HG23	2.03	0.58
1:B:222:LYS:HA	1:B:225[A]:MSE:HE3	1.86	0.58
1:A:266:GLN:HB2	5:A:603:MPD:H51	1.85	0.57
1:A:191:GLY:HA3	1:A:252:ALA:HB3	1.89	0.55
1:A:308:MSE:HE2	1:A:312:MSE:HG3	1.88	0.55
1:B:211:ALA:HB1	1:B:215:MSE:HE1	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ARG:HD3	7:B:641:HOH:O	2.06	0.55
1:A:104:ILE:HA	1:A:276:ALA:HB2	1.91	0.52
5:A:602:MPD:H53	7:A:735:HOH:O	2.08	0.52
3:A:600:G3H:O4P	1:B:243:ARG:HD3	2.10	0.52
1:B:279:LYS:O	1:B:283:GLU:OE1	2.28	0.51
1:A:308:MSE:HE2	1:A:308:MSE:O	2.13	0.49
1:A:260:LEU:HD22	5:A:603:MPD:HM3	1.94	0.48
1:A:255:PHE:CD1	5:A:604:MPD:H11	2.49	0.47
1:A:179:LYS:NZ	6:B:600:G3P:H2	2.30	0.47
1:A:308:MSE:CE	1:A:312:MSE:HG3	2.45	0.47
1:B:308:MSE:HA	4:B:601:MRD:O2	2.15	0.46
1:A:8:LEU:HG	1:A:76:ILE:HD11	1.98	0.46
1:A:222:LYS:CA	1:A:225[B]:MSE:HE3	2.45	0.44
1:A:308:MSE:CE	1:A:312:MSE:CG	2.96	0.43
1:B:24:ASN:ND2	7:B:763:HOH:O	2.39	0.43
1:A:221:GLN:HG2	1:A:225[B]:MSE:HE2	2.00	0.42
1:A:282:LYS:HD3	1:A:282:LYS:HA	1.86	0.42
1:B:215:MSE:CE	1:B:221:GLN:HG3	2.49	0.42
1:B:278:MSE:O	1:B:282:LYS:HG2	2.20	0.42
1:B:143:THR:HA	1:B:164:TYR:O	2.20	0.41
1:A:255:PHE:CE1	5:A:604:MPD:H11	2.55	0.41
1:A:143[A]:THR:HG23	1:A:227:LEU:HB2	2.03	0.41
1:B:230:ILE:HD12	1:B:230:ILE:HA	1.92	0.41
1:B:100:LYS:HE2	1:B:126:SER:OG	2.21	0.41
1:B:137:LYS:HE3	7:B:713:HOH:O	2.20	0.41
1:B:222:LYS:HA	1:B:225[A]:MSE:CE	2.50	0.41
1:A:2:GLU:HG2	1:A:3:MSE:HE3	2.02	0.41
1:B:73:VAL:HA	1:B:96:LEU:O	2.21	0.40
1:A:50:VAL:HG12	1:A:53:SER:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/335 (94%)	308 (98%)	6 (2%)	0	100	100
1	B	312/335 (93%)	303 (97%)	9 (3%)	0	100	100
All	All	626/670 (93%)	611 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/272 (94%)	250 (98%)	6 (2%)	50	53
1	B	253/272 (93%)	246 (97%)	7 (3%)	43	44
All	All	509/544 (94%)	496 (97%)	13 (3%)	46	48

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	GLU
1	A	87	ARG
1	A	159	GLU
1	A	179	LYS
1	A	235	VAL
1	A	308	MSE
1	B	64	GLU
1	B	105	LYS
1	B	179	LYS
1	B	235	VAL
1	B	243	ARG
1	B	283	GLU
1	B	308	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	ASN
1	A	265	ASN
1	B	24	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MRD	B	602	-	7,7,7	0.37	0	9,10,10	0.88	0
5	MPD	A	603	-	7,7,7	0.40	0	9,10,10	0.54	0
5	MPD	A	604	-	7,7,7	0.24	0	9,10,10	0.81	0
2	NDP	A	500	-	45,52,52	1.69	6 (13%)	53,80,80	1.59	9 (16%)
4	MRD	B	601	-	7,7,7	0.31	0	9,10,10	0.37	0
4	MRD	A	601	-	7,7,7	0.41	0	9,10,10	0.56	0
4	MRD	B	603	-	7,7,7	0.32	0	9,10,10	0.40	0
4	MRD	B	604	-	7,7,7	0.35	0	9,10,10	0.67	0
6	G3P	B	600	-	9,9,9	0.69	0	11,12,12	1.23	1 (9%)
5	MPD	A	602	-	7,7,7	0.44	0	9,10,10	1.08	1 (11%)
2	NDP	B	500	-	45,52,52	1.77	6 (13%)	53,80,80	1.79	8 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	G3H	A	600	-	8,9,9	2.15	1 (12%)	10,12,12	1.14	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MRD	B	602	-	-	0/5/5/5	-
5	MPD	A	603	-	-	0/5/5/5	-
5	MPD	A	604	-	-	0/5/5/5	-
2	NDP	A	500	-	-	10/30/77/77	0/5/5/5
4	MRD	B	601	-	-	0/5/5/5	-
4	MRD	A	601	-	-	0/5/5/5	-
4	MRD	B	603	-	-	0/5/5/5	-
4	MRD	B	604	-	-	0/5/5/5	-
6	G3P	B	600	-	1/1/2/2	3/8/8/8	-
5	MPD	A	602	-	-	0/5/5/5	-
2	NDP	B	500	-	-	8/30/77/77	0/5/5/5
3	G3H	A	600	-	1/1/2/3	1/7/8/8	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NDP	O7N-C7N	7.32	1.41	1.24
2	B	500	NDP	O7N-C7N	6.71	1.40	1.24
3	A	600	G3H	O1-C1	5.59	1.42	1.19
2	B	500	NDP	C2A-N1A	5.04	1.43	1.33
2	B	500	NDP	C2A-N3A	4.94	1.40	1.32
2	A	500	NDP	C2A-N3A	3.90	1.38	1.32
2	A	500	NDP	C2A-N1A	3.49	1.40	1.33
2	A	500	NDP	C6N-C5N	2.99	1.38	1.33
2	B	500	NDP	C6A-C5A	-2.50	1.34	1.43
2	B	500	NDP	C6N-C5N	2.37	1.37	1.33
2	B	500	NDP	C4A-N3A	-2.37	1.32	1.35
2	A	500	NDP	C4A-N3A	-2.30	1.32	1.35
2	A	500	NDP	C6A-C5A	-2.02	1.35	1.43

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	NDP	C5A-C6A-N6A	-5.98	111.27	120.35
2	B	500	NDP	C1B-N9A-C4A	5.33	136.00	126.64
2	A	500	NDP	N3A-C2A-N1A	-5.03	120.81	128.68
2	B	500	NDP	C4A-C5A-N7A	4.09	113.66	109.40
2	B	500	NDP	O4D-C1D-N1N	4.02	115.92	108.06
2	A	500	NDP	O4B-C1B-C2B	-3.99	99.66	106.59
2	A	500	NDP	O4D-C1D-N1N	3.69	115.27	108.06
2	B	500	NDP	C3B-C2B-C1B	-3.37	96.55	102.89
2	B	500	NDP	O4B-C1B-C2B	-3.33	100.81	106.59
2	B	500	NDP	N6A-C6A-N1A	3.31	125.44	118.57
2	A	500	NDP	O4D-C1D-C2D	-2.85	100.42	106.64
2	A	500	NDP	C4A-C5A-N7A	2.84	112.36	109.40
5	A	602	MPD	CM-C2-C1	-2.75	104.84	110.57
2	A	500	NDP	C3B-C2B-C1B	-2.71	97.80	102.89
2	A	500	NDP	N6A-C6A-N1A	2.67	124.11	118.57
3	A	600	G3H	O4P-P-O3P	2.65	117.78	107.64
6	B	600	G3P	O2-C2-C1	2.51	120.18	109.12
2	A	500	NDP	PN-O3-PA	-2.41	124.56	132.83
2	B	500	NDP	N3A-C2A-N1A	-2.29	125.10	128.68
2	A	500	NDP	C3D-C2D-C1D	2.07	105.36	101.43

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	B	600	G3P	C2
3	A	600	G3H	C2

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	NDP	C5D-O5D-PN-O1N
2	B	500	NDP	C5D-O5D-PN-O1N
3	A	600	G3H	O1-C1-C2-C3
6	B	600	G3P	O2-C2-C3-O1P
6	B	600	G3P	C1-C2-C3-O1P
2	A	500	NDP	C5D-O5D-PN-O3
2	B	500	NDP	C5D-O5D-PN-O3
2	B	500	NDP	C2D-C1D-N1N-C6N
2	A	500	NDP	C5D-O5D-PN-O2N
2	B	500	NDP	C5D-O5D-PN-O2N
2	B	500	NDP	O4D-C1D-N1N-C6N
6	B	600	G3P	O1-C1-C2-O2
2	A	500	NDP	C2D-C1D-N1N-C6N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	500	NDP	C4B-C5B-O5B-PA
2	A	500	NDP	O4D-C1D-N1N-C6N
2	B	500	NDP	C4B-C5B-O5B-PA
2	A	500	NDP	C4D-C5D-O5D-PN
2	B	500	NDP	O4B-C4B-C5B-O5B
2	B	500	NDP	C4D-C5D-O5D-PN
2	A	500	NDP	C2B-O2B-P2B-O3X
2	A	500	NDP	O4B-C4B-C5B-O5B
2	A	500	NDP	PA-O3-PN-O1N


There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	603	MPD	2	0
5	A	604	MPD	2	0
2	A	500	NDP	1	0
4	B	601	MRD	1	0
6	B	600	G3P	1	0
5	A	602	MPD	1	0
2	B	500	NDP	1	0
3	A	600	G3H	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/335 (90%)	0.28	11 (3%) 42 42	22, 31, 40, 54	0
1	B	302/335 (90%)	0.13	5 (1%) 70 68	21, 31, 40, 51	0
All	All	604/670 (90%)	0.21	16 (2%) 56 54	21, 31, 40, 54	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	90	VAL	3.9
1	A	64	GLU	3.1
1	A	263	SER	3.0
1	A	52	GLU	2.8
1	B	283	GLU	2.7
1	A	39	ASP	2.5
1	A	285	LYS	2.5
1	A	262	GLU	2.4
1	B	98	LEU	2.4
1	A	216	PHE	2.4
1	A	167	GLU	2.4
1	A	258	LEU	2.3
1	B	52	GLU	2.3
1	A	98	LEU	2.3
1	B	233	LEU	2.2
1	B	176	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MPD	A	604	8/8	0.76	0.28	65,67,69,69	0
5	MPD	A	602	8/8	0.76	0.29	54,55,57,57	0
4	MRD	B	604	8/8	0.78	0.17	69,72,73,74	0
4	MRD	B	602	8/8	0.82	0.24	45,46,48,49	0
4	MRD	B	603	8/8	0.84	0.23	51,52,55,56	0
5	MPD	A	603	8/8	0.90	0.22	57,58,60,61	0
4	MRD	A	601	8/8	0.94	0.11	30,33,34,35	0
2	NDP	A	500	48/48	0.94	0.12	18,25,38,40	0
4	MRD	B	601	8/8	0.95	0.19	25,28,29,30	0
2	NDP	B	500	48/48	0.96	0.08	16,22,34,35	0
6	G3P	B	600	10/10	0.98	0.08	18,23,25,25	0
3	G3H	A	600	10/10	0.98	0.08	18,21,22,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NDP A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP B 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.