



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 10:47 am BST

PDB ID : 3ZBR  
Title : Catalytic domain of mouse 2',3'-cyclic nucleotide 3'- phosphodiesterase, with mutation H230S, crystallized with NADP  
Authors : Myllykoski, M.; Raasakka, A.; Lehtimäki, M.; Han, H.; Kursula, P.  
Deposited on : 2012-11-13  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	<div> <div>5%</div> <div>79%</div> <div>17%</div> <div>• •</div> </div>
1	B	221	<div> <div>17%</div> <div>76%</div> <div>19%</div> <div>• 5%</div> </div>



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	215	Total 3359	C 1074	H 1691	N 279	O 310	S 5	0	0	0
1	B	211	Total 3297	C 1054	H 1662	N 274	O 302	S 5	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	158	GLY	-	expression tag	UNP P16330
A	230	SER	HIS	engineered mutation	UNP P16330
B	158	GLY	-	expression tag	UNP P16330
B	230	SER	HIS	engineered mutation	UNP P16330

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		
2	B	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		
3	B	37	Total	O	0	0
			37	37		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE



- Molecule 1: 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.10Å 47.10Å 111.17Å 90.00° 90.26° 90.00°	Depositor
Resolution (Å)	29.12 – 2.30 29.12 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.12-2.30) 97.4 (29.12-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 2.31Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.220 , 0.261 0.222 , 0.266	Depositor DCC
$R_{free}$ test set	966 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtriage
Anisotropy	0.662	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 24.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.397 for h,-k,-l	Xtriage
Reported twinning fraction	0.460 for h,-k,-l	Depositor
Outliers	0 of 19304 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6890	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1704	0.48	0/2292
1	B	0.24	0/1669	0.46	0/2243
All	All	0.25	0/3373	0.47	0/4535

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1668	1691	1688	29	0
1	B	1635	1662	1657	23	3
2	A	48	25	25	0	0
2	B	48	25	25	0	0
3	A	51	0	0	4	0
3	B	37	0	0	2	0
All	All	3487	3403	3395	52	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:ALA:O	1:B:332:GLN:NE2	2.21	0.72
1:A:348:ARG:NH1	3:A:2046:HOH:O	2.24	0.69
1:A:322:GLN:OE1	3:A:2041:HOH:O	2.11	0.69
1:B:194:ASN:ND2	3:B:2006:HOH:O	2.24	0.68
1:B:271:THR:OG1	1:B:326:ASP:OD2	2.12	0.66
1:A:297:SER:O	3:A:2037:HOH:O	2.16	0.61
1:A:300:GLU:O	1:A:332:GLN:NE2	2.33	0.60
1:B:198:PHE:HA	1:B:359:TRP:CZ2	2.39	0.58
1:B:262:LYS:NZ	3:B:2023:HOH:O	2.28	0.58
1:A:218:VAL:O	1:A:224:ARG:NH2	2.38	0.56
1:B:342:ALA:HA	1:B:351:LEU:O	2.06	0.56
1:B:201:GLU:O	1:B:205:PHE:N	2.41	0.53
1:B:225:PRO:HG2	1:B:311:THR:O	2.09	0.53
1:A:192:LEU:HD21	1:A:270:VAL:HG21	1.92	0.51
1:B:246:GLU:N	1:B:246:GLU:OE1	2.43	0.51
1:A:251:GLU:O	1:A:255:ARG:HG3	2.12	0.50
1:B:201:GLU:HB3	1:B:204:HIS:HB2	1.93	0.50
1:B:167:LEU:N	1:B:376:TYR:O	2.44	0.50
1:B:165:LEU:HB3	1:B:166:PRO:CD	2.42	0.50
1:A:295:LYS:NZ	1:A:300:GLU:HB2	2.26	0.49
1:B:208:GLY:O	1:B:209:ASP:HB2	2.12	0.48
1:B:171:TRP:CZ3	1:B:233:THR:CG2	2.97	0.48
1:B:259:LYS:HD3	1:B:259:LYS:N	2.29	0.48
1:A:348:ARG:NH2	3:A:2047:HOH:O	2.48	0.47
1:A:298:ALA:HB1	1:A:328:LEU:HB3	1.97	0.47
1:A:300:GLU:HG2	1:A:301:GLY:N	2.29	0.47
1:B:329:ASP:OD1	1:B:358:ARG:NH1	2.48	0.47
1:A:192:LEU:CD2	1:A:270:VAL:HG21	2.45	0.46
1:A:165:LEU:HB3	1:A:166:PRO:CD	2.46	0.46
1:A:167:LEU:N	1:A:376:TYR:O	2.50	0.45
1:A:294:ASP:O	1:A:296:PRO:HD3	2.17	0.45
1:B:273:LYS:HD3	1:B:322:GLN:HG2	1.99	0.44
1:A:295:LYS:HZ2	1:A:300:GLU:HB2	1.81	0.44
1:A:223:LYS:HE2	1:A:315:ALA:O	2.18	0.44
1:A:295:LYS:HD2	1:A:295:LYS:C	2.37	0.44
1:B:315:ALA:HB3	1:B:318:VAL:CG2	2.48	0.44
1:A:260:ALA:HB1	1:A:370:LYS:HE2	2.00	0.43
1:B:319:GLN:H	1:B:319:GLN:CD	2.22	0.42
1:A:234:LYS:HB3	1:A:247:TYR:CE2	2.54	0.42
1:A:187:VAL:O	1:A:191:GLU:HG3	2.20	0.42
1:A:222:GLY:O	1:A:224:ARG:HD2	2.20	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:ASP:O	1:A:333:GLN:HG2	2.21	0.41
1:B:208:GLY:O	1:B:209:ASP:CB	2.68	0.41
1:B:208:GLY:C	1:B:209:ASP:OD1	2.59	0.41
1:A:296:PRO:O	1:A:298:ALA:N	2.53	0.41
1:A:294:ASP:CG	1:A:294:ASP:O	2.58	0.41
1:A:208:GLY:O	1:A:209:ASP:CB	2.69	0.40
1:A:256:SER:HA	1:A:259:LYS:HG2	2.02	0.40
1:B:166:PRO:HA	1:B:377:TYR:CD2	2.55	0.40
1:A:272:PRO:HD3	1:A:357:GLY:O	2.22	0.40
1:B:203:ARG:O	1:B:206:ILE:O	2.39	0.40
1:A:353:SER:HB2	1:A:359:TRP:CZ3	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:SER:O	1:B:257:TYR:OH[2_745]	1.97	0.23
1:B:240:LYS:O	1:B:255:ARG:NH1[2_645]	2.15	0.05
1:B:161:GLU:OE2	1:B:214:LYS:N[2_755]	2.15	0.05

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/221 (96%)	198 (94%)	12 (6%)	1 (0%)	29	35
1	B	205/221 (93%)	193 (94%)	12 (6%)	0	100	100
All	All	416/442 (94%)	391 (94%)	24 (6%)	1 (0%)	47	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	297	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/180 (98%)	173 (98%)	3 (2%)	60	76
1	B	172/180 (96%)	168 (98%)	4 (2%)	50	67
All	All	348/360 (97%)	341 (98%)	7 (2%)	55	72

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	ARG
1	A	295	LYS
1	A	319	GLN
1	B	207	SER
1	B	209	ASP
1	B	224	ARG
1	B	319	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	319	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	B	1379	-	45,52,52	0.84	1 (2%)	56,80,80	1.18	5 (8%)
2	NAP	A	1379	-	45,52,52	0.80	1 (2%)	56,80,80	1.21	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	1379	-	-	9/31/67/67	0/5/5/5
2	NAP	A	1379	-	-	9/31/67/67	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1379	NAP	C5A-C4A	2.56	1.47	1.40
2	A	1379	NAP	C5A-C4A	2.47	1.47	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1379	NAP	PN-O3-PA	-3.81	119.77	132.83
2	A	1379	NAP	N3A-C2A-N1A	-3.78	122.77	128.68
2	B	1379	NAP	N3A-C2A-N1A	-3.56	123.11	128.68
2	B	1379	NAP	C3D-C2D-C1D	3.38	106.07	100.98

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1379	NAP	PN-O3-PA	-3.18	121.91	132.83
2	A	1379	NAP	C3D-C2D-C1D	3.11	105.66	100.98
2	A	1379	NAP	C4A-C5A-N7A	-2.86	106.42	109.40
2	B	1379	NAP	C4A-C5A-N7A	-2.58	106.71	109.40
2	A	1379	NAP	C2A-N1A-C6A	2.19	122.50	118.75
2	B	1379	NAP	C2A-N1A-C6A	2.06	122.28	118.75

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1379	NAP	C5D-O5D-PN-O1N
2	B	1379	NAP	O4B-C4B-C5B-O5B
2	B	1379	NAP	C3B-C4B-C5B-O5B
2	B	1379	NAP	O4D-C4D-C5D-O5D
2	A	1379	NAP	C4D-C5D-O5D-PN
2	A	1379	NAP	O4B-C4B-C5B-O5B
2	B	1379	NAP	PN-O3-PA-O5B
2	A	1379	NAP	PA-O3-PN-O5D
2	A	1379	NAP	O4D-C4D-C5D-O5D
2	B	1379	NAP	C5D-O5D-PN-O3
2	A	1379	NAP	C2B-O2B-P2B-O3X
2	B	1379	NAP	C5D-O5D-PN-O2N
2	B	1379	NAP	C3D-C4D-C5D-O5D
2	A	1379	NAP	PN-O3-PA-O1A
2	A	1379	NAP	C3B-C4B-C5B-O5B
2	A	1379	NAP	C2B-O2B-P2B-O2X
2	B	1379	NAP	C5B-O5B-PA-O1A
2	A	1379	NAP	C3D-C4D-C5D-O5D


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/221 (97%)	0.68	12 (5%) 24 30	18, 34, 61, 112	0
1	B	211/221 (95%)	1.11	37 (17%) 1 1	28, 45, 78, 142	0
All	All	426/442 (96%)	0.90	49 (11%) 4 7	18, 40, 71, 142	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	355	GLY	7.5
1	B	208	GLY	7.4
1	B	356	LYS	5.4
1	B	189	LEU	4.7
1	B	205	PHE	4.6
1	A	227	GLY	4.5
1	B	293	LEU	4.2
1	A	293	LEU	4.1
1	B	298	ALA	4.1
1	A	298	ALA	3.8
1	B	343	VAL	3.6
1	B	302	LEU	3.5
1	B	206	ILE	3.4
1	B	301	GLY	3.4
1	B	244	ALA	3.2
1	B	203	ARG	3.2
1	B	321	VAL	3.1
1	A	336	GLY	3.1
1	B	339	GLN	2.9
1	B	199	LYS	2.8
1	B	354	LEU	2.7
1	B	281	LEU	2.7
1	A	161	GLU	2.7
1	B	207	SER	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	371	ALA	2.6
1	B	209	ASP	2.6
1	B	338	SER	2.5
1	B	337	GLY	2.5
1	B	365	LYS	2.5
1	B	308	ALA	2.4
1	B	187	VAL	2.4
1	A	209	ASP	2.4
1	B	346	LEU	2.4
1	A	202	LEU	2.3
1	A	208	GLY	2.3
1	B	336	GLY	2.3
1	B	245	GLU	2.3
1	A	291	SER	2.3
1	A	297	SER	2.3
1	B	358	ARG	2.2
1	B	215	LEU	2.2
1	B	196	LYS	2.2
1	A	164	PHE	2.2
1	B	335	LYS	2.1
1	B	252	VAL	2.1
1	B	216	GLU	2.1
1	B	370	LYS	2.1
1	A	228	VAL	2.1
1	B	279	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.


## 6.4 Ligands ⓘ


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAP	B	1379	48/48	0.88	0.17	25,57,79,209	0
2	NAP	A	1379	48/48	0.89	0.21	23,57,92,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.