



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 04:29 am BST

PDB ID : 4ZG4
Title : Myosin Vc Pre-powerstroke
Authors : Ropars, V.; Pylypenko, O.; Sweeney, L.; Houdusse, A.
Deposited on : 2015-04-22
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	764	<div> <div>12%</div> <div> <div></div> <div>89%</div> <div>•</div> <div>8%</div> </div> </div>
1	E	764	<div> <div>10%</div> <div> <div></div> <div>90%</div> <div>•</div> <div>7%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 21924 atoms, of which 10274 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called myosin-Vc.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	705	Total	C	H	N	O	S	0	5	0
			10590	3480	5116	928	1042	24			
1	E	711	Total	C	H	N	O	S	0	1	0
			10633	3497	5128	938	1047	23			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	756	GLY	-	expression tag	UNP Q9NQX4
B	757	ASP	-	expression tag	UNP Q9NQX4
B	758	TYR	-	expression tag	UNP Q9NQX4
B	759	LYS	-	expression tag	UNP Q9NQX4
B	760	ASP	-	expression tag	UNP Q9NQX4
B	761	ASP	-	expression tag	UNP Q9NQX4
B	762	ASP	-	expression tag	UNP Q9NQX4
B	763	ASP	-	expression tag	UNP Q9NQX4
B	764	LYS	-	expression tag	UNP Q9NQX4
E	756	GLY	-	expression tag	UNP Q9NQX4
E	757	ASP	-	expression tag	UNP Q9NQX4
E	758	TYR	-	expression tag	UNP Q9NQX4
E	759	LYS	-	expression tag	UNP Q9NQX4
E	760	ASP	-	expression tag	UNP Q9NQX4
E	761	ASP	-	expression tag	UNP Q9NQX4
E	762	ASP	-	expression tag	UNP Q9NQX4
E	763	ASP	-	expression tag	UNP Q9NQX4
E	764	LYS	-	expression tag	UNP Q9NQX4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	1	Total	Mg	0	0
			1	1		

- Molecule 3 is VANADATE ION (three-letter code: VO4) (formula: O₄V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	V	0	0
			5	4	1		
3	E	1	Total	O	V	0	0
			5	4	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
4	E	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	363	Total 363	O 363	0	0
6	E	238	Total 238	O 238	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: myosin-Vc



• Molecule 1: myosin-Vc



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	118.28Å 67.32Å 133.18Å 90.00° 108.48° 90.00°	Depositor
Resolution (Å)	49.54 – 2.36 49.54 – 2.36	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.54-2.36) 99.5 (49.54-2.36)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.37Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.173 , 0.213 0.179 , 0.216	Depositor DCC
R_{free} test set	4094 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.0	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21924	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VO4, MG, EDO, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.23	0/5604	0.40	0/7598
1	E	0.23	0/5616	0.40	0/7616
All	All	0.23	0/11220	0.40	0/15214

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	5474	5116	5099	11	0
1	E	5505	5128	5128	7	0
2	B	1	0	0	0	0
2	E	1	0	0	0	0
3	B	5	0	0	0	0
3	E	5	0	0	0	0
4	B	27	12	12	0	0
4	E	27	12	12	0	0
5	B	4	6	6	0	0
6	B	363	0	0	2	0
6	E	238	0	0	2	0
All	All	11650	10274	10257	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:648:ARG:NH2	6:B:901:HOH:O	2.28	0.66
1:E:656:LYS:NZ	6:E:902:HOH:O	2.32	0.62
1:B:34:GLY:O	1:B:36:LYS:N	2.35	0.59
1:B:741:ARG:O	1:B:743:GLY:N	2.36	0.59
1:E:466:HIS:ND1	6:E:901:HOH:O	2.32	0.59
1:B:461:GLN:NE2	1:B:491:GLN:OE1	2.38	0.56
1:B:42:LEU:N	1:B:45:GLY:O	2.42	0.52
1:B:389:MET:HE2	1:B:393:GLN:HB3	1.94	0.49
1:E:553:ASP:OD1	1:E:554:LYS:N	2.46	0.49
1:E:188:ALA:HB3	1:E:189:HIS:HA	1.97	0.47
1:B:526:TYR:HA	1:B:530:VAL:HG23	1.99	0.45
1:E:491:GLN:N	1:E:492:PRO:CD	2.81	0.44
1:E:446:SER:OG	1:E:447:PHE:N	2.52	0.42
1:B:446:SER:OG	1:B:447:PHE:N	2.50	0.42
1:B:491:GLN:N	1:B:492:PRO:CD	2.83	0.42
1:B:398:ARG:NH2	6:B:911:HOH:O	2.50	0.41
1:E:526:TYR:HA	1:E:530:VAL:HG23	2.03	0.41
1:B:391:ARG:HB3	1:B:392:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	696/764 (91%)	679 (98%)	16 (2%)	1 (0%)	51 63
1	E	696/764 (91%)	679 (98%)	16 (2%)	1 (0%)	51 63
All	All	1392/1528 (91%)	1358 (98%)	32 (2%)	2 (0%)	51 63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	241	LEU
1	B	241	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	553/680 (81%)	548 (99%)	5 (1%)	78	87
1	E	553/680 (81%)	541 (98%)	12 (2%)	52	63
All	All	1106/1360 (81%)	1089 (98%)	17 (2%)	65	76

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	86	ASN
1	B	195	LEU
1	B	437	TYR
1	B	478	GLU
1	B	648	ARG
1	E	3	VAL
1	E	86	ASN
1	E	189	HIS
1	E	357	VAL
1	E	365	GLU
1	E	366	SER
1	E	437	TYR
1	E	478	GLU
1	E	486	ASP
1	E	501	MET
1	E	532	ARG
1	E	576	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	EDO	B	804	-	3,3,3	0.47	0	2,2,2	0.29	0
4	ADP	B	803	3,2	24,29,29	0.94	1 (4%)	29,45,45	1.11	3 (10%)
4	ADP	E	803	3,2	24,29,29	1.09	2 (8%)	29,45,45	1.11	4 (13%)
3	VO4	E	802	2,4	1,4,4	0.70	0	-		
3	VO4	B	802	2,4	1,4,4	0.86	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	804	-	-	0/1/1/1	-
4	ADP	B	803	3,2	-	1/12/32/32	0/3/3/3
4	ADP	E	803	3,2	-	1/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	803	ADP	O4'-C1'	3.43	1.45	1.41
4	B	803	ADP	O4'-C1'	3.36	1.45	1.41
4	E	803	ADP	PA-O1A	2.35	1.59	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	803	ADP	C2'-C3'-C4'	-3.04	96.73	102.64
4	E	803	ADP	C2'-C3'-C4'	-2.72	97.37	102.64
4	E	803	ADP	C5-C6-N6	2.61	124.32	120.35
4	B	803	ADP	C5-C6-N6	2.54	124.22	120.35
4	E	803	ADP	C1'-N9-C4	-2.19	122.79	126.64
4	B	803	ADP	C1'-N9-C4	-2.06	123.02	126.64
4	E	803	ADP	C3'-C2'-C1'	-2.04	97.91	100.98

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	803	ADP	PA-O3A-PB-O3B
4	B	803	ADP	PA-O3A-PB-O3B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	705/764 (92%)	0.92	89 (12%) 3 6	39, 62, 138, 188	0
1	E	711/764 (93%)	0.80	76 (10%) 6 9	41, 75, 139, 178	0
All	All	1416/1528 (92%)	0.86	165 (11%) 4 7	39, 69, 139, 188	0

All (165) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	748	LEU	8.9
1	E	67	VAL	8.7
1	E	686	TYR	8.5
1	E	747	TYR	8.3
1	E	695	PHE	7.6
1	E	740	PHE	7.6
1	E	720	VAL	7.5
1	E	702	LEU	7.4
1	E	738	ILE	7.4
1	B	728	SER	7.3
1	E	696	TYR	7.0
1	E	717	CYS	7.0
1	E	729	ASN	7.0
1	B	7	TYR	6.8
1	B	6	LEU	6.6
1	B	25	ALA	6.6
1	B	60	LEU	6.5
1	B	740	PHE	6.4
1	B	13	VAL	6.4
1	B	724	LEU	6.3
1	B	745	VAL	6.3
1	B	725	ILE	6.2
1	E	66	LEU	6.2
1	E	690	TRP	6.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	686	TYR	5.9
1	E	728	SER	5.6
1	B	699	TYR	5.5
1	B	27	ILE	5.5
1	B	746	ALA	5.4
1	B	692	TYR	5.4
1	E	739	PHE	5.4
1	B	741	ARG	5.4
1	B	41	LEU	5.3
1	E	708	LEU	5.3
1	E	700	GLY	5.3
1	B	696	TYR	5.3
1	B	40	LEU	5.3
1	B	687	PRO	5.2
1	B	717	CYS	5.0
1	B	719	VAL	5.0
1	B	77	TYR	5.0
1	B	738	ILE	5.0
1	E	745	VAL	4.9
1	B	695	PHE	4.9
1	E	721	LEU	4.8
1	B	749	GLU	4.8
1	B	748	LEU	4.7
1	B	28	ALA	4.7
1	E	692	TYR	4.7
1	B	697	SER	4.6
1	E	697	SER	4.6
1	E	482	TRP	4.5
1	B	721	LEU	4.5
1	B	42	LEU	4.4
1	E	699	TYR	4.4
1	B	3	VAL	4.4
1	B	727	ASP	4.4
1	B	739	PHE	4.4
1	B	689	ARG	4.4
1	B	747	TYR	4.3
1	E	709	SER	4.3
1	E	733	PHE	4.3
1	B	744	GLN	4.2
1	B	723	ARG	4.2
1	E	719	VAL	4.2
1	E	735	LYS	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	754	ASP	4.1
1	E	475	TYR	4.0
1	E	742	ALA	4.0
1	E	731	TYR	4.0
1	E	693	ILE	3.9
1	B	688	SER	3.9
1	E	60	LEU	3.9
1	B	54	PRO	3.8
1	B	732	GLN	3.8
1	B	2	ALA	3.8
1	B	720	VAL	3.7
1	B	731	TYR	3.7
1	B	698	ARG	3.7
1	B	751	LEU	3.6
1	E	522	LEU	3.6
1	B	57	LEU	3.5
1	E	691	THR	3.5
1	E	50	TYR	3.5
1	B	722	HIS	3.5
1	E	25	ALA	3.5
1	E	707	GLU	3.5
1	E	7	TYR	3.4
1	B	750	LYS	3.4
1	E	685	SER	3.4
1	E	479	ASP	3.4
1	B	22	TRP	3.3
1	E	72	LEU	3.3
1	E	751	LEU	3.3
1	B	694	GLU	3.2
1	E	736	THR	3.2
1	B	726	GLN	3.2
1	B	716	VAL	3.2
1	E	726	GLN	3.2
1	B	714	LYS	3.2
1	B	45	GLY	3.2
1	E	732	GLN	3.1
1	B	14	TRP	3.1
1	B	48	LEU	3.1
1	E	507	LEU	3.0
1	E	65	ILE	3.0
1	E	698	ARG	3.0
1	B	736	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	718	LYS	3.0
1	E	734	GLY	3.0
1	E	701	ILE	3.0
1	B	55	GLU	3.0
1	B	59	PRO	2.9
1	E	478	GLU	2.9
1	B	693	ILE	2.9
1	B	84	LEU	2.9
1	B	58	PRO	2.9
1	B	730	GLN	2.9
1	E	741	ARG	2.9
1	B	38	LEU	2.8
1	E	752	ARG	2.8
1	B	5	GLU	2.8
1	E	485	ILE	2.8
1	B	50	TYR	2.7
1	E	724	LEU	2.7
1	B	30	ASP	2.7
1	B	743	GLY	2.6
1	E	694	GLU	2.6
1	B	12	ARG	2.6
1	E	633	TYR	2.6
1	B	8	THR	2.5
1	E	722	HIS	2.5
1	E	13	VAL	2.5
1	E	11	ASN	2.5
1	B	53	ASN	2.5
1	B	26	GLU	2.5
1	E	584	HIS	2.5
1	B	32	ARG	2.4
1	E	705	LYS	2.4
1	B	4	ALA	2.4
1	B	24	SER	2.4
1	B	729	ASN	2.4
1	B	36	LYS	2.4
1	E	476	MET	2.4
1	E	483	THR	2.4
1	B	480	ILE	2.3
1	B	189	HIS	2.3
1	B	47	GLU	2.2
1	E	737	LYS	2.2
1	E	472	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	150	ARG	2.2
1	B	650	ILE	2.2
1	E	574	LEU	2.2
1	E	727	ASP	2.2
1	E	480	ILE	2.1
1	B	49	ASP	2.1
1	B	563	LEU	2.1
1	E	484	LEU	2.1
1	E	288	MET	2.1
1	B	31	TYR	2.1
1	E	200	ILE	2.1
1	E	706	GLN	2.1
1	E	436	ILE	2.0
1	B	159	VAL	2.0
1	B	733	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	B	804	4/4	0.92	0.22	74,88,98,110	0
2	MG	B	801	1/1	0.93	0.23	44,44,44,44	0
2	MG	E	801	1/1	0.96	0.21	53,53,53,53	0
4	ADP	E	803	27/27	0.98	0.19	46,56,66,78	0
4	ADP	B	803	27/27	0.98	0.22	35,48,59,63	0
3	VO4	B	802	5/5	0.99	0.22	41,42,47,54	0
3	VO4	E	802	5/5	1.00	0.19	45,50,51,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.