



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 11, 2021 – 10:19 PM GMT

PDB ID : 6ZJ6
Title : Structure of the GH99 endo-alpha-mannanase from Bacteroides xylanisolvens in complex with cyclohexylmethyl-Glc-1,3-isofagomine
Authors : Thompson, A.J.; Sobala, L.F.; Fernandes, P.Z.; Hakki, Z.; Howe, J.D.; Hill, M.; Zitzmann, N.; Davies, S.; Stamatakis, Z.; Butters, T.D.; Alonzi, D.S.; Williams, S.J.; Davies, G.J.
Deposited on : 2020-06-27
Resolution : 1.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	385	<div> <div>4%</div> <div>85%</div> <div>8%</div> <div>6%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyl hydrolase family 71.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	360	Total	C	N	O	S	0	50	0
			3175	2074	509	583	9			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-4	MET	-	initiating methionine	UNP D6D1V7
AAA	-3	GLY	-	expression tag	UNP D6D1V7
AAA	-2	SER	-	expression tag	UNP D6D1V7
AAA	-1	SER	-	expression tag	UNP D6D1V7
AAA	0	HIS	-	expression tag	UNP D6D1V7
AAA	1	HIS	-	expression tag	UNP D6D1V7
AAA	2	HIS	-	expression tag	UNP D6D1V7
AAA	3	HIS	-	expression tag	UNP D6D1V7
AAA	4	HIS	-	expression tag	UNP D6D1V7
AAA	5	HIS	-	expression tag	UNP D6D1V7
AAA	6	SER	-	expression tag	UNP D6D1V7
AAA	7	SER	-	expression tag	UNP D6D1V7
AAA	8	GLY	-	expression tag	UNP D6D1V7
AAA	9	LEU	-	expression tag	UNP D6D1V7
AAA	10	VAL	-	expression tag	UNP D6D1V7
AAA	11	PRO	-	expression tag	UNP D6D1V7
AAA	12	ARG	-	expression tag	UNP D6D1V7
AAA	13	GLY	-	expression tag	UNP D6D1V7
AAA	14	SER	-	expression tag	UNP D6D1V7
AAA	15	HIS	-	expression tag	UNP D6D1V7
AAA	16	MET	-	expression tag	UNP D6D1V7

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	1
			4	2	2		
2	AAA	1	Total	C	O	0	0
			4	2	2		
2	AAA	1	Total	C	O	0	1
			4	2	2		
2	AAA	1	Total	C	O	0	1
			4	2	2		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is methylcyclohexane (three-letter code: QTE) (formula: C_7H_{14}) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	C	0	1
			14	14		

- Molecule 5 is 5-HYDROXYMETHYL-3,4-DIHYDROXYPIPERIDINE (three-letter code: IFM) (formula: $C_6H_{13}NO_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	AAA	1	Total	C	N	O	0	0
			10	6	1	3		

- Molecule 6 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	C	O	0	0
			11	6	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	463	Total	O	0	37
			468	468		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycosyl hydrolase family 71



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	107.98Å 107.98Å 67.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.35 – 1.09 76.35 – 1.09	Depositor EDS
% Data completeness (in resolution range)	95.9 (76.35-1.09) 95.9 (76.35-1.09)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.09Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.113 , 0.136 0.114 , 0.136	Depositor DCC
R_{free} test set	7638 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3698	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IFM, QTE, GLC, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.77	3/3412 (0.1%)	0.97	13/4637 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	377	GLU	CD-OE2	10.35	1.37	1.25
1	AAA	324	SER	CB-OG	6.33	1.50	1.42
1	AAA	324	SER	CA-CB	5.57	1.61	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	176[A]	ARG	NE-CZ-NH1	9.87	125.24	120.30
1	AAA	176[B]	ARG	NE-CZ-NH1	9.87	125.24	120.30
1	AAA	185	ARG	NE-CZ-NH1	-9.09	115.75	120.30
1	AAA	185	ARG	NE-CZ-NH2	7.87	124.24	120.30
1	AAA	316	ARG	CG-CD-NE	6.95	126.40	111.80
1	AAA	316	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	AAA	377	GLU	CB-CG-CD	-5.12	100.37	114.20
1	AAA	176[A]	ARG	NE-CZ-NH2	-5.11	117.75	120.30
1	AAA	176[B]	ARG	NE-CZ-NH2	-5.11	117.75	120.30
1	AAA	375	PHE	CB-CG-CD1	5.04	124.33	120.80
1	AAA	104[A]	ASN	CB-CA-C	5.02	120.45	110.40
1	AAA	104[B]	ASN	CB-CA-C	5.02	120.45	110.40
1	AAA	47	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3175	0	3151	26	0
2	AAA	16	0	24	1	0
3	AAA	4	0	3	1	0
4	AAA	14	0	0	0	0
5	AAA	10	0	12	1	0
6	AAA	11	0	9	0	0
7	AAA	468	0	0	13	0
All	All	3698	0	3199	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:198[B]:TYR:HB2	7:AAA:570:HOH:O	1.80	0.80
1:AAA:271:GLN:HE21	1:AAA:275[B]:LYS:HE3	1.55	0.71
1:AAA:198[B]:TYR:CD1	7:AAA:570:HOH:O	2.44	0.69
1:AAA:271:GLN:NE2	1:AAA:275[B]:LYS:HE3	2.10	0.66
1:AAA:198[B]:TYR:CG	7:AAA:570:HOH:O	2.52	0.62
1:AAA:309[B]:GLN:HG2	7:AAA:734:HOH:O	1.99	0.61
1:AAA:186:LYS:HE2	3:AAA:405:ACT:O	2.01	0.61
1:AAA:271:GLN:HE21	1:AAA:275[B]:LYS:CE	2.15	0.59
1:AAA:155:LEU:HD13	1:AAA:166[B]:LEU:HD11	1.86	0.57
1:AAA:317[A]:LYS:HE3	7:AAA:591:HOH:O	2.07	0.54
1:AAA:155:LEU:HD12	1:AAA:194[B]:ILE:CD1	2.40	0.52
1:AAA:148[A]:LYS:HE2	7:AAA:608:HOH:O	2.09	0.51
1:AAA:235[A]:THR:HG23	7:AAA:517:HOH:O	2.13	0.48
1:AAA:183:PHE:CZ	2:AAA:403[B]:EDO:H22	2.49	0.48
1:AAA:135[B]:LYS:NZ	7:AAA:508:HOH:O	2.45	0.48
1:AAA:293[B]:ARG:HD3	1:AAA:351:PHE:CZ	2.49	0.47
1:AAA:293[B]:ARG:HD2	7:AAA:771:HOH:O	2.15	0.46
1:AAA:148[B]:LYS:NZ	7:AAA:512:HOH:O	2.48	0.46
1:AAA:189[A]:LYS:NZ	7:AAA:504:HOH:O	2.34	0.44
1:AAA:375:PHE:CE2	1:AAA:379:LYS:HE2	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:309[A]:GLN:HG3	7:AAA:691:HOH:O	2.19	0.42
1:AAA:228:GLY:O	1:AAA:249:PHE:HA	2.18	0.42
1:AAA:230:TRP:C	1:AAA:230:TRP:CD1	2.94	0.41
1:AAA:252:TYR:CZ	5:AAA:407:IFM:H1C1	2.56	0.41
1:AAA:293[B]:ARG:HD3	1:AAA:351:PHE:CE1	2.56	0.40
1:AAA:293[A]:ARG:NH1	7:AAA:519:HOH:O	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	406/385 (106%)	395 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	353/330 (107%)	349 (99%)	4 (1%)	73	40

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	48	TRP
1	AAA	73	SER
1	AAA	315	TYR
1	AAA	333	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	AAA	405	-	1,3,3	1.07	0	0,3,3	0.00	-
4	QTE	AAA	406[B]	6	7,7,7	0.34	0	8,8,8	0.14	0
4	QTE	AAA	406[A]	6	7,7,7	0.42	0	8,8,8	0.30	0
6	GLC	AAA	408	5,4	11,11,12	1.50	3 (27%)	15,15,17	1.52	3 (20%)
2	EDO	AAA	404[B]	-	3,3,3	0.35	0	2,2,2	0.59	0
5	IFM	AAA	407	6	9,10,10	1.50	1 (11%)	9,13,13	0.99	0
2	EDO	AAA	403[B]	-	3,3,3	0.18	0	2,2,2	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	AAA	402	-	3,3,3	0.58	0	2,2,2	0.45	0
2	EDO	AAA	401[B]	-	3,3,3	0.34	0	2,2,2	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	QTE	AAA	406[A]	6	-	-	0/1/1/1
4	QTE	AAA	406[B]	6	-	-	0/1/1/1
6	GLC	AAA	408	5,4	-	0/2/19/22	0/1/1/1
2	EDO	AAA	404[B]	-	-	0/1/1/1	-
5	IFM	AAA	407	6	-	0/2/16/16	0/1/1/1
2	EDO	AAA	403[B]	-	-	0/1/1/1	-
2	EDO	AAA	402	-	-	0/1/1/1	-
2	EDO	AAA	401[B]	-	-	0/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	AAA	407	IFM	C2-C3	4.01	1.56	1.52
6	AAA	408	GLC	C2-C3	2.78	1.56	1.52
6	AAA	408	GLC	O3-C3	-2.46	1.37	1.43
6	AAA	408	GLC	O5-C5	-2.32	1.38	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	AAA	408	GLC	C1-O5-C5	4.11	117.75	112.19
6	AAA	408	GLC	O2-C2-C3	2.37	114.89	110.14
6	AAA	408	GLC	C6-C5-C4	-2.02	108.26	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	405	ACT	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	AAA	407	IFM	1	0
2	AAA	403[B]	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand QTE AAA 406 (A)



Ligand GLC AAA 408





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	AAA	360/385 (93%)	-0.38	14 (3%)	39 36	10, 15, 35, 75	8 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	74	GLY	5.3
1	AAA	380	GLN	4.1
1	AAA	21	PRO	4.0
1	AAA	22	SER	3.6
1	AAA	72	GLY	3.4
1	AAA	27[A]	ASN	3.1
1	AAA	25[A]	GLU	2.8
1	AAA	23	ASN	2.8
1	AAA	28[A]	GLY	2.7
1	AAA	73	SER	2.6
1	AAA	26[A]	ASN	2.3
1	AAA	75	GLN	2.2
1	AAA	71	GLY	2.2
1	AAA	70	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	QTE	AAA	406[B]	7/7	0.81	0.23	32,43,53,55	7
4	QTE	AAA	406[A]	7/7	0.81	0.23	26,30,35,36	7
2	EDO	AAA	402	4/4	0.93	0.16	25,34,38,39	0
3	ACT	AAA	405	4/4	0.95	0.09	20,21,23,26	0
6	GLC	AAA	408	11/12	0.96	0.06	14,16,20,22	0
2	EDO	AAA	401[B]	4/4	0.96	0.08	11,12,12,13	4
2	EDO	AAA	404[B]	4/4	0.97	0.07	12,13,15,15	4
5	IFM	AAA	407	10/10	0.98	0.05	11,11,14,14	0
2	EDO	AAA	403[B]	4/4	0.99	0.06	9,10,11,11	4

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around QTE AAA 406 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GLC AAA 408:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IFM AAA 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.