



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 5, 2022 – 09:12 pm GMT

PDB ID : 6ZK4  
Title : Plant nucleoside hydrolase - ZmNRh2b with a bound adenine  
Authors : Morera, S.; Vigouroux, A.; Kopecny, D.  
Deposited on : 2020-06-29  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.24  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.24


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>87%</span> <span>6%</span> <span>8%</span> </div> </div>
1	B	343	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>87%</span> <span>6%</span> <span>8%</span> </div> </div>
1	C	343	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">4%</span> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>87%</span> <span>5%</span> <span>8%</span> </div> </div>
1	D	343	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -10px; left: 0;">9%</span> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>86%</span> <span>6%</span> <span>8%</span> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyrimidine-specific ribonucleoside hydrolase rihA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	1	0
			2401	1545	401	445	10			
1	B	317	Total	C	N	O	S	0	0	0
			2408	1547	405	446	10			
1	C	317	Total	C	N	O	S	0	0	0
			2408	1547	405	446	10			
1	D	316	Total	C	N	O	S	0	0	0
			2397	1541	401	445	10			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP B6THD4
A	-16	GLY	-	expression tag	UNP B6THD4
A	-15	SER	-	expression tag	UNP B6THD4
A	-14	SER	-	expression tag	UNP B6THD4
A	-13	HIS	-	expression tag	UNP B6THD4
A	-12	HIS	-	expression tag	UNP B6THD4
A	-11	HIS	-	expression tag	UNP B6THD4
A	-10	HIS	-	expression tag	UNP B6THD4
A	-9	HIS	-	expression tag	UNP B6THD4
A	-8	HIS	-	expression tag	UNP B6THD4
A	-7	SER	-	expression tag	UNP B6THD4
A	-6	GLN	-	expression tag	UNP B6THD4
A	-5	ASP	-	expression tag	UNP B6THD4
A	-4	PRO	-	expression tag	UNP B6THD4
A	-3	ASN	-	expression tag	UNP B6THD4
A	-2	SER	-	expression tag	UNP B6THD4
A	-1	SER	-	expression tag	UNP B6THD4
A	0	SER	-	expression tag	UNP B6THD4
B	-17	MET	-	initiating methionine	UNP B6THD4
B	-16	GLY	-	expression tag	UNP B6THD4
B	-15	SER	-	expression tag	UNP B6THD4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	SER	-	expression tag	UNP B6THD4
B	-13	HIS	-	expression tag	UNP B6THD4
B	-12	HIS	-	expression tag	UNP B6THD4
B	-11	HIS	-	expression tag	UNP B6THD4
B	-10	HIS	-	expression tag	UNP B6THD4
B	-9	HIS	-	expression tag	UNP B6THD4
B	-8	HIS	-	expression tag	UNP B6THD4
B	-7	SER	-	expression tag	UNP B6THD4
B	-6	GLN	-	expression tag	UNP B6THD4
B	-5	ASP	-	expression tag	UNP B6THD4
B	-4	PRO	-	expression tag	UNP B6THD4
B	-3	ASN	-	expression tag	UNP B6THD4
B	-2	SER	-	expression tag	UNP B6THD4
B	-1	SER	-	expression tag	UNP B6THD4
B	0	SER	-	expression tag	UNP B6THD4
C	-17	MET	-	initiating methionine	UNP B6THD4
C	-16	GLY	-	expression tag	UNP B6THD4
C	-15	SER	-	expression tag	UNP B6THD4
C	-14	SER	-	expression tag	UNP B6THD4
C	-13	HIS	-	expression tag	UNP B6THD4
C	-12	HIS	-	expression tag	UNP B6THD4
C	-11	HIS	-	expression tag	UNP B6THD4
C	-10	HIS	-	expression tag	UNP B6THD4
C	-9	HIS	-	expression tag	UNP B6THD4
C	-8	HIS	-	expression tag	UNP B6THD4
C	-7	SER	-	expression tag	UNP B6THD4
C	-6	GLN	-	expression tag	UNP B6THD4
C	-5	ASP	-	expression tag	UNP B6THD4
C	-4	PRO	-	expression tag	UNP B6THD4
C	-3	ASN	-	expression tag	UNP B6THD4
C	-2	SER	-	expression tag	UNP B6THD4
C	-1	SER	-	expression tag	UNP B6THD4
C	0	SER	-	expression tag	UNP B6THD4
D	-17	MET	-	initiating methionine	UNP B6THD4
D	-16	GLY	-	expression tag	UNP B6THD4
D	-15	SER	-	expression tag	UNP B6THD4
D	-14	SER	-	expression tag	UNP B6THD4
D	-13	HIS	-	expression tag	UNP B6THD4
D	-12	HIS	-	expression tag	UNP B6THD4
D	-11	HIS	-	expression tag	UNP B6THD4
D	-10	HIS	-	expression tag	UNP B6THD4
D	-9	HIS	-	expression tag	UNP B6THD4

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-8	HIS	-	expression tag	UNP B6THD4
D	-7	SER	-	expression tag	UNP B6THD4
D	-6	GLN	-	expression tag	UNP B6THD4
D	-5	ASP	-	expression tag	UNP B6THD4
D	-4	PRO	-	expression tag	UNP B6THD4
D	-3	ASN	-	expression tag	UNP B6THD4
D	-2	SER	-	expression tag	UNP B6THD4
D	-1	SER	-	expression tag	UNP B6THD4
D	0	SER	-	expression tag	UNP B6THD4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0

- Molecule 3 is ADENINE (three-letter code: ADE) (formula: C<sub>5</sub>H<sub>5</sub>N<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			10	5	5		
3	B	1	Total	C	N	0	0
			10	5	5		
3	C	1	Total	C	N	0	1
			20	10	10		
3	D	1	Total	C	N	0	0
			10	5	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	193	Total	O	0	0
			193	193		
6	B	141	Total	O	0	0
			141	141		
6	C	106	Total	O	0	0
			106	106		
6	D	80	Total	O	0	0
			80	80		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA




- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA




- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA



- Molecule 1: Pyrimidine-specific ribonucleoside hydrolase rihA







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.64Å 148.18Å 88.41Å 90.00° 102.71° 90.00°	Depositor
Resolution (Å)	42.86 – 1.95 42.86 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (42.86-1.95) 99.0 (42.86-1.95)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.42 (at 1.95Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.175 , 0.203 0.185 , 0.215	Depositor DCC
$R_{free}$ test set	4299 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.4	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.039 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10206	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ADE, EDO, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2460	0.62	0/3341
1	B	0.52	0/2464	0.62	0/3345
1	C	0.49	0/2464	0.61	0/3345
1	D	0.47	0/2453	0.61	0/3331
All	All	0.51	0/9841	0.62	0/13362

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2401	0	2419	11	0
1	B	2408	0	2423	9	0
1	C	2408	0	2423	8	0
1	D	2397	0	2410	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	10	0	4	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	4	0	0
3	C	20	0	8	0	0
3	D	10	0	4	0	0
4	B	14	0	20	0	0
5	D	4	0	6	1	0
6	A	193	0	0	0	0
6	B	141	0	0	0	0
6	C	106	0	0	0	0
6	D	80	0	0	0	0
All	All	10206	0	9721	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:290:LYS:HE3	5:D:403:EDO:H21	1.81	0.63
1:B:42:THR:HB	1:B:113:ALA:HB1	1.82	0.61
1:C:42:THR:HB	1:C:113:ALA:HB1	1.82	0.61
1:D:42:THR:HB	1:D:113:ALA:HB1	1.83	0.59
1:A:42:THR:HB	1:A:113:ALA:HB1	1.83	0.59
1:A:182:GLU:HG2	1:C:93:SER:HB3	1.85	0.59
1:C:275:GLN:HG3	1:D:275:GLN:HE22	1.70	0.56
1:D:207:LEU:HG	1:D:248:LEU:HD11	1.91	0.53
1:A:207:LEU:HG	1:A:248:LEU:HD11	1.92	0.51
1:A:158:VAL:HB	1:A:196[B]:VAL:HG12	1.93	0.50
1:B:207:LEU:HG	1:B:248:LEU:HD11	1.94	0.49
1:C:207:LEU:HG	1:C:248:LEU:HD11	1.95	0.49
1:A:36:VAL:HG21	1:A:258:LEU:HD11	1.95	0.49
1:A:236:TRP:HE1	3:A:402:ADE:N6	2.11	0.48
1:A:188:PHE:CE1	1:A:196[B]:VAL:HG11	2.49	0.47
1:C:36:VAL:HG21	1:C:258:LEU:HD11	1.97	0.46
1:C:17:THR:HG21	1:C:27:ILE:HD11	1.97	0.46
1:A:283:LEU:HD11	1:B:172:ALA:HA	1.96	0.46
1:A:17:THR:HG21	1:A:27:ILE:HD11	1.99	0.45
1:B:32:ARG:NH2	1:B:222:HIS:H	2.15	0.45
1:C:283:LEU:HD11	1:D:172:ALA:HA	1.99	0.45
1:B:36:VAL:HG21	1:B:258:LEU:HD11	1.98	0.44
1:D:17:THR:HG21	1:D:27:ILE:HD11	1.99	0.44
1:B:17:THR:HG21	1:B:27:ILE:HD11	1.98	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:144:LYS:HD2	1:D:186:ILE:HD13	1.98	0.43
1:B:19:PRO:HA	1:B:20:GLY:HA3	1.84	0.43
1:D:36:VAL:HG21	1:D:258:LEU:HD11	2.01	0.41
1:B:43:THR:HG21	1:B:54:THR:HA	2.01	0.41
1:D:43:THR:HG21	1:D:54:THR:HA	2.01	0.41
1:A:172:ALA:HA	1:B:283:LEU:HD11	2.02	0.41
1:A:31:PHE:HE1	1:A:69:VAL:HG21	1.85	0.41
1:C:19:PRO:HA	1:C:20:GLY:HA3	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/343 (92%)	310 (98%)	5 (2%)	0	100	100
1	B	315/343 (92%)	309 (98%)	6 (2%)	0	100	100
1	C	315/343 (92%)	310 (98%)	5 (2%)	0	100	100
1	D	314/343 (92%)	309 (98%)	5 (2%)	0	100	100
All	All	1259/1372 (92%)	1238 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/283 (92%)	257 (99%)	2 (1%)	81	80
1	B	259/283 (92%)	256 (99%)	3 (1%)	71	68
1	C	259/283 (92%)	255 (98%)	4 (2%)	65	60
1	D	258/283 (91%)	253 (98%)	5 (2%)	57	50
All	All	1035/1132 (91%)	1021 (99%)	14 (1%)	67	62

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	199	ILE
1	A	234	ARG
1	B	199	ILE
1	B	234	ARG
1	B	316	SER
1	C	145	ARG
1	C	155	LYS
1	C	199	ILE
1	C	234	ARG
1	D	110	GLU
1	D	199	ILE
1	D	234	ARG
1	D	283	LEU
1	D	284	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	275	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	PEG	B	403	-	6,6,6	0.16	0	5,5,5	0.08	0
4	PEG	B	404	-	6,6,6	0.13	0	5,5,5	0.09	0
3	ADE	A	402	-	9,11,11	0.93	0	7,15,15	1.09	1 (14%)
3	ADE	C	402[A]	-	9,11,11	0.86	0	7,15,15	1.07	1 (14%)
3	ADE	C	402[B]	-	9,11,11	0.82	0	7,15,15	1.09	1 (14%)
5	EDO	D	403	-	3,3,3	0.57	0	2,2,2	0.27	0
3	ADE	B	402	-	9,11,11	0.87	0	7,15,15	1.05	1 (14%)
3	ADE	D	402	-	9,11,11	0.84	0	7,15,15	1.03	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	403	-	-	3/4/4/4	-
4	PEG	B	404	-	-	3/4/4/4	-
3	ADE	A	402	-	-	-	0/2/2/2
3	ADE	C	402[A]	-	-	-	0/2/2/2
5	EDO	D	403	-	-	0/1/1/1	-
3	ADE	C	402[B]	-	-	-	0/2/2/2
3	ADE	B	402	-	-	-	0/2/2/2
3	ADE	D	402	-	-	-	0/2/2/2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	402[B]	ADE	C5-C6-N6	2.31	123.87	120.35
3	A	402	ADE	C5-C6-N6	2.31	123.86	120.35
3	C	402[A]	ADE	C5-C6-N6	2.30	123.85	120.35
3	D	402	ADE	C5-C6-N6	2.19	123.69	120.35
3	B	402	ADE	C5-C6-N6	2.17	123.65	120.35

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	403	PEG	O1-C1-C2-O2
4	B	404	PEG	O1-C1-C2-O2
4	B	403	PEG	O2-C3-C4-O4
4	B	404	PEG	C4-C3-O2-C2
4	B	404	PEG	C1-C2-O2-C3
4	B	403	PEG	C4-C3-O2-C2


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	ADE	1	0
5	D	403	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## Ligand ADE C 402 (A)




Bond lengths



Bond angles





Torsions




Rings

## Ligand ADE C 402 (B)







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/343 (92%)	0.11	3 (0%) 84 89	28, 37, 56, 75	0
1	B	317/343 (92%)	-0.01	5 (1%) 72 79	30, 42, 70, 93	0
1	C	317/343 (92%)	0.29	15 (4%) 31 41	32, 52, 87, 118	0
1	D	316/343 (92%)	0.48	30 (9%) 8 13	29, 55, 99, 117	0
All	All	1266/1372 (92%)	0.22	53 (4%) 36 45	28, 45, 86, 118	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	248	LEU	7.1
1	D	243	PHE	4.7
1	D	239	LYS	4.4
1	D	233	TYR	4.4
1	D	325	PRO	3.9
1	D	242	GLY	3.8
1	D	89	PHE	3.7
1	B	213	LEU	3.4
1	D	105	SER	3.3
1	D	320	LYS	3.2
1	C	318	ILE	3.2
1	C	66	CYS	3.2
1	D	262	TYR	3.1
1	D	324	ALA	3.0
1	D	321	LEU	3.0
1	D	206	CYS	3.0
1	D	224	ALA	3.0
1	D	102	PRO	2.9
1	C	315	ILE	2.8
1	C	36	VAL	2.8
1	D	323	MET	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	85	ARG	2.7
1	D	247	PHE	2.7
1	D	312	PRO	2.6
1	C	151	SER	2.6
1	C	120	LYS	2.6
1	D	240	SER	2.6
1	D	99	LEU	2.5
1	A	196[A]	VAL	2.5
1	C	311	VAL	2.4
1	B	325	PRO	2.4
1	D	98	ASN	2.4
1	C	145	ARG	2.4
1	B	244	HIS	2.4
1	C	211	ASP	2.3
1	D	232	PHE	2.3
1	B	210	GLU	2.2
1	D	83	LYS	2.2
1	D	244	HIS	2.2
1	C	67	PRO	2.2
1	A	136	LEU	2.2
1	C	325	PRO	2.2
1	C	224	ALA	2.2
1	A	223	ALA	2.2
1	D	213	LEU	2.1
1	C	147	PRO	2.1
1	D	107	LYS	2.1
1	D	65	GLY	2.1
1	C	248	LEU	2.1
1	D	322	LEU	2.1
1	B	81	GLY	2.0
1	C	121	VAL	2.0
1	D	32	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PEG	B	403	7/7	0.77	0.17	57,60,62,64	0
3	ADE	A	402	10/10	0.80	0.14	45,48,51,52	0
3	ADE	C	402[B]	10/10	0.87	0.15	42,43,45,46	10
3	ADE	D	402	10/10	0.87	0.13	71,76,79,79	0
3	ADE	C	402[A]	10/10	0.87	0.15	38,40,43,44	10
3	ADE	B	402	10/10	0.88	0.11	51,54,56,56	0
4	PEG	B	404	7/7	0.88	0.12	65,66,69,70	0
5	EDO	D	403	4/4	0.94	0.08	66,67,68,69	0
2	CA	C	401	1/1	0.98	0.09	41,41,41,41	0
2	CA	D	401	1/1	0.98	0.07	40,40,40,40	0
2	CA	A	401	1/1	0.99	0.09	31,31,31,31	0
2	CA	B	401	1/1	0.99	0.09	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around ADE A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ADE C 402 (B):**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ADE D 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ADE C 402 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ADE B 402:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.