



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 07:59 pm BST

PDB ID : 3ZLI  
Title : Crystal structure of JmjC domain of human histone demethylase UTY  
Authors : Vollmar, M.; Gileadi, C.; Shrestha, L.; Goubin, S.; Johansson, C.; Krojer, T.; Raynor, J.W.; Bradley, A.; von Delft, F.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.; Oppermann, U.  
Deposited on : 2013-01-31  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	
1	B	478	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE DEMETHYLASE UTY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	11	1
			3573	2295	607	650	21			
1	B	439	Total	C	N	O	S	0	9	1
			3545	2278	601	646	20			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	877	MET	-	expression tag	UNP O14607
A	1348	ALA	-	expression tag	UNP O14607
A	1349	GLU	-	expression tag	UNP O14607
A	1350	ASN	-	expression tag	UNP O14607
A	1351	LEU	-	expression tag	UNP O14607
A	1352	TYR	-	expression tag	UNP O14607
A	1353	PHE	-	expression tag	UNP O14607
A	1354	GLN	-	expression tag	UNP O14607
B	877	MET	-	expression tag	UNP O14607
B	1348	ALA	-	expression tag	UNP O14607
B	1349	GLU	-	expression tag	UNP O14607
B	1350	ASN	-	expression tag	UNP O14607
B	1351	LEU	-	expression tag	UNP O14607
B	1352	TYR	-	expression tag	UNP O14607
B	1353	PHE	-	expression tag	UNP O14607
B	1354	GLN	-	expression tag	UNP O14607

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula:  $C_5H_6O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Fe	0	0
			1	1		
4	A	1	Total	Fe	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		
5	A	1	Total	Zn	0	0
			1	1		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	286	Total 286	O 286	0	0
6	B	263	Total 263	O 263	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: HISTONE DEMETHYLASE UTY

Chain A: 



#### • Molecule 1: HISTONE DEMETHYLASE UTY

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.58Å 110.25Å 118.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 1.80 49.96 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.01-1.80) 100.0 (49.96-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.176 , 0.204 0.184 , 0.212	Depositor DCC
$R_{free}$ test set	5494 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.5	Xtriage
Anisotropy	0.517	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 37.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7747	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AKG, ZN, FE2, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	1/3701 (0.0%)	0.76	2/5047 (0.0%)
1	B	0.69	0/3666	0.74	1/4999 (0.0%)
All	All	0.71	1/7367 (0.0%)	0.75	3/10046 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1344	SER	C-N	-5.13	1.22	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1156	ARG	NE-CZ-NH2	-7.40	116.60	120.30
1	A	1329	ASP	CB-CG-OD2	5.86	123.58	118.30
1	B	1156	ARG	NE-CZ-NH2	-5.71	117.44	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3573	0	3473	24	0
1	B	3545	0	3461	16	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	32	0	48	4	0
2	B	24	0	36	1	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	286	0	0	4	0
6	B	263	0	0	1	0
All	All	7747	0	7026	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:890:ILE:HD12	1:B:903:LEU:CD1	2.02	0.89
1:A:905[B]:GLN:OE1	6:A:2030:HOH:O	1.89	0.88
1:A:1224[B]:MET:SD	1:A:1228[B]:ILE:HG12	2.24	0.77
1:A:1247:LEU:HD21	1:A:1331:ILE:HD13	1.70	0.72
1:A:1068:HIS:ND1	1:A:1228[B]:ILE:HD11	2.10	0.66
1:B:890:ILE:CD1	1:B:903:LEU:CD1	2.75	0.65
1:A:1068:HIS:CE1	1:A:1228[B]:ILE:HD11	2.32	0.64
1:B:1075:GLY:HA2	1:B:1080:GLN:HG3	1.82	0.62
1:B:890:ILE:HD12	1:B:903:LEU:HD12	1.81	0.58
2:B:2350:EDO:H12	6:B:2114:HOH:O	2.04	0.58
1:A:1131:ASN:HB3	2:A:2345:EDO:H12	1.87	0.57
1:A:1075:GLY:HA2	1:A:1080:GLN:HG3	1.87	0.57
2:A:2351:EDO:H12	6:A:2133:HOH:O	2.06	0.54
1:B:1166:TRP:CH2	1:B:1188[A]:VAL:HG21	2.45	0.52
1:A:1224[B]:MET:CE	1:A:1228[B]:ILE:HG12	2.41	0.51
1:A:1224[B]:MET:HE3	1:A:1228[B]:ILE:HB	1.93	0.50
1:A:1342:SER:O	1:B:1245:LYS:HE3	2.11	0.49
1:A:1225:ALA:HB2	1:A:1243:LEU:HD11	1.94	0.49
1:B:1225:ALA:HB2	1:B:1243:LEU:HD11	1.94	0.49
1:B:890:ILE:CD1	1:B:903:LEU:HD12	2.41	0.49
1:A:1327:MET:O	1:A:1331:ILE:HG12	2.14	0.48
1:A:949:THR:HB	1:A:973[B]:ASN:OD1	2.14	0.47
1:A:1075:GLY:O	2:A:2351:EDO:H11	2.16	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1270:THR:HG22	1:B:1273:GLU:HB2	1.98	0.45
1:A:976:HIS:HE1	6:A:2053:HOH:O	2.00	0.44
1:A:1224[B]:MET:HE1	1:A:1228[B]:ILE:HG21	2.00	0.44
1:B:892:LEU:HD22	1:B:897:ASP:HB3	2.00	0.44
1:A:1342:SER:O	1:B:1245:LYS:CE	2.66	0.43
1:B:1270:THR:CG2	1:B:1273:GLU:HB2	2.49	0.43
1:A:967:ILE:HB	2:A:2345:EDO:H21	2.01	0.43
1:B:890:ILE:HD12	1:B:903:LEU:HD11	1.96	0.42
1:B:1166:TRP:CZ3	1:B:1188[A]:VAL:HG21	2.55	0.42
1:A:1224[B]:MET:HE3	1:A:1228[B]:ILE:CB	2.50	0.41
1:A:920:LEU:C	1:A:920:LEU:HD13	2.40	0.41
1:A:974:ARG:NH2	6:A:2053:HOH:O	2.53	0.41
1:B:1314:LYS:HD3	1:B:1314:LYS:HA	1.79	0.41
1:A:1227:ASN:C	1:A:1228[B]:ILE:CD1	2.89	0.41
1:B:1205:TRP:CZ2	1:B:1209:LYS:HE3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/478 (94%)	441 (98%)	9 (2%)	0	100	100
1	B	444/478 (93%)	436 (98%)	8 (2%)	0	100	100
All	All	894/956 (94%)	877 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/432 (90%)	389 (100%)	2 (0%)	88	87
1	B	391/432 (90%)	386 (99%)	5 (1%)	69	62
All	All	782/864 (90%)	775 (99%)	7 (1%)	81	75

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	896	ARG
1	A	1149	GLU
1	B	893	GLU
1	B	925	LYS
1	B	960	ASP
1	B	1329[A]	ASP
1	B	1329[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	2348	-	3,3,3	0.39	0	2,2,2	0.27	0
2	EDO	A	2351	-	3,3,3	0.79	0	2,2,2	0.77	0
2	EDO	A	2349	-	3,3,3	0.46	0	2,2,2	0.60	0
2	EDO	A	2347	-	3,3,3	0.57	0	2,2,2	0.66	0
2	EDO	B	2349	-	3,3,3	0.41	0	2,2,2	0.23	0
2	EDO	B	2346	-	3,3,3	0.38	0	2,2,2	0.39	0
2	EDO	B	2350	-	3,3,3	0.86	0	2,2,2	0.27	0
2	EDO	A	2350	-	3,3,3	0.32	0	2,2,2	0.57	0
3	AKG	A	4000	4	3,9,9	0.50	0	4,11,11	1.58	2 (50%)
2	EDO	A	2345	-	3,3,3	0.43	0	2,2,2	0.12	0
2	EDO	B	2348	-	3,3,3	0.33	0	2,2,2	0.07	0
3	AKG	B	4000	4	3,9,9	1.40	0	4,11,11	2.27	2 (50%)
2	EDO	B	2347	-	3,3,3	0.73	0	2,2,2	0.14	0
2	EDO	A	2352	-	3,3,3	0.34	0	2,2,2	0.60	0
2	EDO	B	2345	-	3,3,3	0.45	0	2,2,2	0.12	0
2	EDO	A	2346	-	3,3,3	0.33	0	2,2,2	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	2348	-	-	0/1/1/1	-
2	EDO	A	2351	-	-	0/1/1/1	-
2	EDO	A	2349	-	-	1/1/1/1	-
2	EDO	A	2347	-	-	1/1/1/1	-
2	EDO	B	2349	-	-	0/1/1/1	-
2	EDO	B	2346	-	-	1/1/1/1	-
2	EDO	B	2350	-	-	1/1/1/1	-
2	EDO	A	2350	-	-	0/1/1/1	-
3	AKG	A	4000	4	-	2/3/9/9	-
2	EDO	A	2345	-	-	0/1/1/1	-
2	EDO	B	2348	-	-	1/1/1/1	-
3	AKG	B	4000	4	-	2/3/9/9	-
2	EDO	B	2347	-	-	0/1/1/1	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	2352	-	-	1/1/1/1	-
2	EDO	B	2345	-	-	0/1/1/1	-
2	EDO	A	2346	-	-	0/1/1/1	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	4000	AKG	O5-C2-C3	-3.78	113.92	120.38
3	A	4000	AKG	O5-C2-C3	-2.33	116.40	120.38
3	B	4000	AKG	C3-C4-C5	-2.25	108.89	112.67
3	A	4000	AKG	C3-C4-C5	-2.05	109.23	112.67

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2347	EDO	O1-C1-C2-O2
3	B	4000	AKG	C1-C2-C3-C4
3	A	4000	AKG	C1-C2-C3-C4
3	B	4000	AKG	O5-C2-C3-C4
3	A	4000	AKG	O5-C2-C3-C4
2	B	2346	EDO	O1-C1-C2-O2
2	A	2352	EDO	O1-C1-C2-O2
2	A	2349	EDO	O1-C1-C2-O2
2	B	2348	EDO	O1-C1-C2-O2
2	B	2350	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2351	EDO	2	0
2	B	2350	EDO	1	0
2	A	2345	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	443/478 (92%)	-0.40	2 (0%) 91 89	19, 30, 48, 79	0
1	B	439/478 (91%)	-0.24	8 (1%) 68 64	21, 32, 54, 81	0
All	All	882/956 (92%)	-0.32	10 (1%) 80 78	19, 31, 51, 81	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	900	PHE	4.5
1	B	899	PHE	3.4
1	A	962	THR	2.9
1	B	1328	GLU	2.6
1	A	964	THR	2.5
1	B	893	GLU	2.4
1	B	901	PRO	2.3
1	B	962	THR	2.2
1	B	1026	PHE	2.2
1	B	897	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.



## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	A	2351	4/4	0.78	0.19	34,45,48,50	0
2	EDO	A	2352	4/4	0.82	0.25	45,51,51,58	0
2	EDO	B	2350	4/4	0.84	0.16	45,46,47,59	0
2	EDO	A	2347	4/4	0.84	0.16	32,43,45,53	0
2	EDO	A	2345	4/4	0.88	0.38	43,47,48,53	0
2	EDO	B	2348	4/4	0.90	0.15	53,57,58,60	0
2	EDO	A	2348	4/4	0.90	0.14	43,46,50,51	0
2	EDO	B	2347	4/4	0.91	0.13	33,41,42,46	0
2	EDO	A	2349	4/4	0.92	0.11	47,49,49,52	0
3	AKG	A	4000	10/10	0.94	0.10	24,29,33,33	0
2	EDO	B	2345	4/4	0.94	0.10	30,32,35,35	0
2	EDO	A	2350	4/4	0.95	0.09	42,43,45,47	0
3	AKG	B	4000	10/10	0.95	0.09	25,32,35,36	0
2	EDO	B	2346	4/4	0.95	0.14	47,48,50,52	0
2	EDO	B	2349	4/4	0.95	0.20	45,46,50,52	0
2	EDO	A	2346	4/4	0.96	0.09	31,31,34,34	0
5	ZN	B	4002	1/1	0.98	0.08	29,29,29,29	0
5	ZN	A	4002	1/1	1.00	0.09	27,27,27,27	0
4	FE2	B	4001	1/1	1.00	0.06	25,25,25,25	0
4	FE2	A	4001	1/1	1.00	0.05	24,24,24,24	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.