



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 10:16 am BST

PDB ID : 5ZMJ  
Title : Crystal structure of the Fab region of a neutralizing fully human antibody against GM-CSF  
Authors : Angkawidjaja, C.; Torashima, T.  
Deposited on : 2018-04-04  
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| $R_{free}$            | 130704                      | 7484 (1.84-1.80)                                      |
| Clashscore            | 141614                      | 8401 (1.84-1.80)                                      |
| Ramachandran outliers | 138981                      | 8290 (1.84-1.80)                                      |
| Sidechain outliers    | 138945                      | 8290 (1.84-1.80)                                      |
| RSRZ outliers         | 127900                      | 7371 (1.84-1.80)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain  |
|-----|-------|--------|---|
| 1   | H     | 248    | <div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 84%, grey 14%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>84%</span> <span>7%</span> <span>8%</span> </div> </div>  |
| 2   | L     | 235    | <div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 5%, green 84%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>84%</span> <span>5%</span> <span>10%</span> </div> </div> |

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of a Fab fraction of human IgG.

| Mol | Chain | Residues | Atoms |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 1   | H     | 227      | Total | C    | N   | O   | S | 0       | 1       | 0     |
|     |       |          | 1728  | 1102 | 288 | 333 | 5 |         |         |       |

- Molecule 2 is a protein called Light chain of a Fab fraction of human IgG.

| Mol | Chain | Residues | Atoms |     |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 2   | L     | 211      | Total | C   | N   | O   | S | 0       | 3       | 0     |
|     |       |          | 1588  | 990 | 269 | 325 | 4 |         |         |       |

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



| Mol | Chain | Residues | Atoms |   |   | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 3   | H     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |
| 3   | L     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Residues | Atoms |   |   | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 3   | L     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |
| 3   | L     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |
| 3   | L     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |
| 3   | L     | 1        | Total | O | S | 0       | 0       |
|     |       |          | 5     | 4 | 1 |         |         |

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


| Mol | Chain | Residues | Atoms |    | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 4   | H     | 2        | Total | Na | 0       | 0       |
|     |       |          | 2     | 2  |         |         |
| 4   | L     | 2        | Total | Na | 0       | 0       |
|     |       |          | 2     | 2  |         |         |

- Molecule 5 is water.

| Mol | Chain | Residues | Atoms |     | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 5   | H     | 264      | Total | O   | 0       | 0       |
|     |       |          | 264   | 264 |         |         |
| 5   | L     | 216      | Total | O   | 0       | 0       |
|     |       |          | 216   | 216 |         |         |



- Molecule 1: Heavy chain of a Fab fraction of human IgG



## 4 Data and refinement statistics

| Property  | Value   | Source           |
|---|---|------------------|
| Space group   | P 21 21 21  | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 59.12Å 66.69Å 129.04Å<br>90.00° 90.00° 90.00°               | Depositor        |
| Resolution (Å)  | 46.37 – 1.81<br>19.80 – 1.81                                | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 97.7 (46.37-1.81)<br>97.8 (19.80-1.81)                      | Depositor<br>EDS |
| $R_{merge}$   | 0.07  | Depositor        |
| $R_{sym}$   | (Not available)   | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 10.41 (at 1.81Å)  | Xtriage          |
| Refinement program  | REFMAC 5.8.0158   | Depositor        |
| R, $R_{free}$   | 0.161 , 0.191<br>0.172 , 0.198                              | Depositor<br>DCC |
| $R_{free}$ test set   | 2270 reflections (4.94%)                                    | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 16.6  | Xtriage          |
| Anisotropy  | 0.023   | Xtriage          |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.37 , 46.2   | EDS              |
| L-test for twinning <sup>2</sup>  | $\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$ | Xtriage          |
| Estimated twinning fraction   | No twinning to report.                                      | Xtriage          |
| $F_o, F_c$ correlation  | 0.95  | EDS              |
| Total number of atoms   | 3830  | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 20.0  | wwPDB-VP         |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |             | Bond angles |             |
|-----|-------|--------------|-------------|-------------|-------------|
|     |       | RMSZ         | $\# Z  > 5$ | RMSZ        | $\# Z  > 5$ |
| 1   | H     | 0.40         | 0/1779      | 0.52        | 0/2425      |
| 2   | L     | 0.32         | 0/1636      | 0.46        | 0/2233      |
| All | All   | 0.37         | 0/3415      | 0.49        | 0/4658      |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | H     | 1728  | 0        | 1692     | 15      | 0            |
| 2   | L     | 1588  | 0        | 1541     | 11      | 0            |
| 3   | H     | 5     | 0        | 0        | 0       | 0            |
| 3   | L     | 25    | 0        | 0        | 0       | 0            |
| 4   | H     | 2     | 0        | 0        | 0       | 0            |
| 4   | L     | 2     | 0        | 0        | 0       | 0            |
| 5   | H     | 264   | 0        | 0        | 4       | 2            |
| 5   | L     | 216   | 0        | 0        | 6       | 1            |
| All | All   | 3830  | 0        | 3233     | 25      | 2            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1             | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|--------------------|------------------|--------------------------|-------------------|
| 2:L:95:SER:HB3     | 5:L:402:HOH:O    | 1.76                     | 0.85              |
| 1:H:162:GLU:OE1    | 5:H:401:HOH:O    | 2.10                     | 0.69              |
| 1:H:3:HIS:HD2      | 5:H:419:HOH:O    | 1.82                     | 0.61              |
| 1:H:105:ASP:HA     | 2:L:33:TYR:OH    | 2.00                     | 0.61              |
| 1:H:157:LYS:HD2    | 5:L:540:HOH:O    | 2.02                     | 0.59              |
| 2:L:28:ASN:HB2     | 2:L:91:THR:CG2   | 2.33                     | 0.59              |
| 1:H:3:HIS:HB3      | 5:H:573:HOH:O    | 2.03                     | 0.57              |
| 2:L:19:ASN:ND2     | 5:L:404:HOH:O    | 2.31                     | 0.56              |
| 1:H:192:LEU:HD12   | 1:H:192:LEU:C    | 2.26                     | 0.56              |
| 2:L:132[B]:ASN:OD1 | 5:L:401:HOH:O    | 2.18                     | 0.53              |
| 1:H:135:VAL:HG12   | 1:H:223:LYS:HG3  | 1.95                     | 0.48              |
| 2:L:28:ASN:HB2     | 2:L:91:THR:HG21  | 1.94                     | 0.48              |
| 2:L:130:GLN:HA     | 2:L:130:GLN:HE21 | 1.79                     | 0.48              |
| 1:H:53:THR:HA      | 1:H:72:LEU:HD11  | 1.97                     | 0.46              |
| 1:H:215:LYS:HD3    | 1:H:215:LYS:O    | 2.17                     | 0.45              |
| 1:H:109:TYR:CE2    | 1:H:111:GLU:HG3  | 2.54                     | 0.43              |
| 1:H:135:VAL:CG1    | 1:H:223:LYS:HG3  | 2.49                     | 0.43              |
| 1:H:82:GLU:HG3     | 5:H:443:HOH:O    | 2.19                     | 0.42              |
| 1:H:144:SER:HB3    | 5:L:474:HOH:O    | 2.19                     | 0.42              |
| 1:H:128:ALA:HA     | 1:H:160:PHE:CE1  | 2.55                     | 0.42              |
| 2:L:154:ALA:HB2    | 2:L:195:TYR:CE2  | 2.54                     | 0.41              |
| 2:L:27:SER:OG      | 5:L:402:HOH:O    | 2.22                     | 0.41              |
| 2:L:63:PHE:CD1     | 2:L:76:ILE:HG12  | 2.55                     | 0.41              |
| 2:L:153:LYS:HB3    | 2:L:156:SER:HA   | 2.02                     | 0.41              |
| 1:H:215:LYS:N      | 1:H:216:PRO:CD   | 2.84                     | 0.40              |

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1        | Atom-2               | Interatomic distance (Å) | Clash overlap (Å) |
|---------------|----------------------|--------------------------|-------------------|
| 5:H:579:HOH:O | 5:L:583:HOH:O[3_755] | 1.96                     | 0.24              |
| 5:H:402:HOH:O | 5:H:553:HOH:O[3_755] | 2.11                     | 0.09              |



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Favoured  | Allowed | Outliers | Percentiles |     |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1   | H     | 226/248 (91%) | 218 (96%) | 8 (4%)  | 0        | 100         | 100 |
| 2   | L     | 212/235 (90%) | 206 (97%) | 6 (3%)  | 0        | 100         | 100 |
| All | All   | 438/483 (91%) | 424 (97%) | 14 (3%) | 0        | 100         | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Rotameric | Outliers | Percentiles |    |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1   | H     | 193/208 (93%) | 191 (99%) | 2 (1%)   | 76          | 70 |
| 2   | L     | 181/197 (92%) | 178 (98%) | 3 (2%)   | 60          | 50 |
| All | All   | 374/405 (92%) | 369 (99%) | 5 (1%)   | 67          | 61 |

All (5) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | H     | 99  | ARG  |
| 1   | H     | 215 | LYS  |
| 2   | L     | 95  | SER  |
| 2   | L     | 130 | GLN  |
| 2   | L     | 156 | SER  |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | H     | 185 | GLN  |
| 2   | L     | 130 | GLN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths |      |             | Bond angles |      |             |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
|     |      |       |     |      | Counts       | RMSZ | $\# Z  > 2$ | Counts      | RMSZ | $\# Z  > 2$ |
| 3   | SO4  | L     | 301 | -    | 4,4,4        | 0.33 | 0           | 6,6,6       | 0.10 | 0           |
| 3   | SO4  | H     | 301 | -    | 4,4,4        | 0.33 | 0           | 6,6,6       | 0.06 | 0           |
| 3   | SO4  | L     | 302 | -    | 4,4,4        | 0.28 | 0           | 6,6,6       | 0.15 | 0           |
| 3   | SO4  | L     | 304 | -    | 4,4,4        | 0.28 | 0           | 6,6,6       | 0.30 | 0           |
| 3   | SO4  | L     | 303 | -    | 4,4,4        | 0.35 | 0           | 6,6,6       | 0.18 | 0           |
| 3   | SO4  | L     | 305 | -    | 4,4,4        | 0.33 | 0           | 6,6,6       | 0.11 | 0           |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed      | <RSRZ> | #RSRZ>2      | OWAB(Å <sup>2</sup> ) | Q<0.9 |
|-----|-------|---------------|--------|--------------|-----------------------|-------|
| 1   | H     | 227/248 (91%) | -0.33  | 3 (1%) 77 74 | 10, 16, 34, 54        | 0     |
| 2   | L     | 211/235 (89%) | -0.29  | 3 (1%) 75 72 | 8, 17, 33, 50         | 0     |
| All | All   | 438/483 (90%) | -0.31  | 6 (1%) 75 72 | 8, 17, 34, 54         | 0     |

All (6) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1   | H     | 204 | GLY  | 4.8  |
| 1   | H     | 205 | THR  | 3.4  |
| 2   | L     | 156 | SER  | 3.2  |
| 2   | L     | 69  | ASP  | 2.4  |
| 2   | L     | 68  | SER  | 2.3  |
| 1   | H     | 107 | THR  | 2.3  |

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR  | B-factors( $\text{\AA}^2$ ) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|-----------------------------|-------|
| 3   | SO4  | L     | 303 | 5/5   | 0.94 | 0.22 | 43,49,61,64                 | 0     |
| 3   | SO4  | L     | 305 | 5/5   | 0.95 | 0.27 | 43,51,61,70                 | 0     |
| 3   | SO4  | L     | 301 | 5/5   | 0.97 | 0.19 | 41,42,42,43                 | 0     |
| 3   | SO4  | L     | 302 | 5/5   | 0.98 | 0.09 | 24,26,30,33                 | 0     |
| 4   | NA   | L     | 307 | 1/1   | 0.99 | 0.17 | 15,15,15,15                 | 0     |
| 3   | SO4  | L     | 304 | 5/5   | 0.99 | 0.08 | 17,20,22,22                 | 0     |
| 4   | NA   | H     | 303 | 1/1   | 0.99 | 0.22 | 10,10,10,10                 | 0     |
| 3   | SO4  | H     | 301 | 5/5   | 0.99 | 0.22 | 40,42,46,56                 | 0     |
| 4   | NA   | L     | 306 | 1/1   | 1.00 | 0.13 | 10,10,10,10                 | 0     |
| 4   | NA   | H     | 302 | 1/1   | 1.00 | 0.15 | 10,10,10,10                 | 0     |

## 6.5 Other polymers [i](#)

There are no such residues in this entry.