



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 05:24 PM EDT

PDB ID : 2ZNE  
Title : Crystal structure of Zn<sup>2+</sup>-bound form of des3-23ALG-2 complexed with Alix  
ABS peptide  
Authors : Suzuki, H.; Kawasaki, M.; Inuzuka, T.; Kakiuchi, T.; Shibata, H.; Wakatsuki,  
S.; Maki, M.  
Deposited on : 2008-04-22  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	 4% 76% 21% ..
1	B	169	 4% 79% 19% ..
2	C	16	 69% 12% 6% 6% 6%
2	D	16	 19% 56% 19% 25%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Programmed cell death protein 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	168	Total	C	N	O	S	0	0	0
			1399	889	242	264	4			
1	B	167	Total	C	N	O	S	0	0	0
			1391	885	241	261	4			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ALA	deletion	UNP O75340
A	?	-	TYR	deletion	UNP O75340
A	?	-	SER	deletion	UNP O75340
A	?	-	TYR	deletion	UNP O75340
A	?	-	ARG	deletion	UNP O75340
A	?	-	PRO	deletion	UNP O75340
A	?	-	GLY	deletion	UNP O75340
A	?	-	PRO	deletion	UNP O75340
A	?	-	GLY	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	GLY	deletion	UNP O75340
A	?	-	PRO	deletion	UNP O75340
A	?	-	GLY	deletion	UNP O75340
A	?	-	PRO	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	GLY	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	LEU	deletion	UNP O75340
A	?	-	PRO	deletion	UNP O75340
A	?	-	ALA	deletion	UNP O75340
A	?	-	TYR	deletion	UNP O75340
A	?	-	SER	deletion	UNP O75340
A	?	-	TYR	deletion	UNP O75340

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ARG	deletion	UNP O75340
B	?	-	PRO	deletion	UNP O75340
B	?	-	GLY	deletion	UNP O75340
B	?	-	PRO	deletion	UNP O75340
B	?	-	GLY	deletion	UNP O75340
B	?	-	ALA	deletion	UNP O75340
B	?	-	GLY	deletion	UNP O75340
B	?	-	PRO	deletion	UNP O75340
B	?	-	GLY	deletion	UNP O75340
B	?	-	PRO	deletion	UNP O75340
B	?	-	ALA	deletion	UNP O75340
B	?	-	ALA	deletion	UNP O75340
B	?	-	GLY	deletion	UNP O75340
B	?	-	ALA	deletion	UNP O75340
B	?	-	ALA	deletion	UNP O75340
B	?	-	LEU	deletion	UNP O75340
B	?	-	PRO	deletion	UNP O75340

- Molecule 2 is a protein called 16-meric peptide from Programmed cell death 6-interacting protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	15	Total	C	N	O	0	0	0
			117	79	16	22			
2	D	12	Total	C	N	O	0	0	0
			95	65	13	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	15	SER	CYS	engineered mutation	UNP Q8WUM4
D	15	SER	CYS	engineered mutation	UNP Q8WUM4

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		
3	B	4	Total	Zn	0	0
			4	4		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Na 1	0	0
4	B	1	Total 1	Na 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total 26	O 26	0	0
5	B	36	Total 36	O 36	0	0
5	C	1	Total 1	O 1	0	0
5	D	5	Total 5	O 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Programmed cell death protein 6



- Molecule 1: Programmed cell death protein 6



- Molecule 2: 16-meric peptide from Programmed cell death 6-interacting protein



- Molecule 2: 16-meric peptide from Programmed cell death 6-interacting protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.45Å 88.51Å 211.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.94 – 2.20 33.94 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.0 (33.94-2.20) 98.0 (33.94-2.20)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.73 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.223 , 0.274 0.222 , 0.270	Depositor DCC
$R_{free}$ test set	1163 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 38.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3080	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.64	0/1431	0.69	0/1933
1	B	0.65	0/1423	0.69	0/1922
2	C	0.83	0/125	1.01	2/174 (1.1%)
2	D	1.21	0/102	0.84	0/143
All	All	0.68	0/3081	0.71	2/4172 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	13	GLY	N-CA-C	-6.00	98.11	113.10
2	C	9	PRO	N-CA-C	-5.52	97.75	112.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	8	TYR	Peptide
2	D	9	PRO	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1399	0	1332	26	0
1	B	1391	0	1329	21	0
2	C	117	0	99	4	0
2	D	95	0	85	1	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	26	0	0	0	0
5	B	36	0	0	1	0
5	C	1	0	0	0	0
5	D	5	0	0	0	0
All	All	3080	0	2845	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:LYS:HE3	1:B:50:GLN:NE2	1.52	1.23
1:B:37:LYS:HE3	1:B:50:GLN:HE21	1.04	1.12
1:A:164:ILE:HG13	1:A:190:ILE:HD11	1.52	0.92
1:B:162:THR:O	1:B:166:ARG:HG3	1.74	0.88
1:B:37:LYS:CE	1:B:50:GLN:HE21	1.93	0.74
1:A:164:ILE:HG13	1:A:190:ILE:CD1	2.19	0.72
1:A:178:VAL:HG11	1:A:186:MET:HE1	1.75	0.68
2:D:8:TYR:CE2	2:D:10:GLY:HA3	2.28	0.68
2:C:8:TYR:CD2	2:C:8:TYR:C	2.69	0.65
1:A:111:ASP:HB3	1:A:145:GLN:HE22	1.61	0.65
1:A:113:ASN:O	1:A:117:GLN:HG2	1.98	0.64
1:B:108:GLY:O	1:B:109:MET:HE2	1.98	0.63
1:B:37:LYS:HE3	1:B:50:GLN:HE22	1.59	0.62
1:B:64:THR:OG1	1:B:153:GLN:HG2	2.02	0.59
1:A:68:ILE:HD13	1:A:92:ILE:HD11	1.88	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:VAL:CG2	1:B:191:VAL:HG22	2.36	0.55
1:A:46:THR:O	1:A:50:GLN:HG3	2.07	0.54
1:B:169:ASP:O	1:B:172:GLN:NE2	2.41	0.54
1:A:178:VAL:HG11	1:A:186:MET:CE	2.38	0.54
1:A:111:ASP:HB3	1:A:145:GLN:NE2	2.24	0.53
2:C:8:TYR:HB2	2:C:9:PRO:HD3	1.88	0.53
1:A:38:ASP:OD2	1:A:40:SER:HB3	2.09	0.53
1:B:31:VAL:O	1:B:35:VAL:HG23	2.11	0.51
1:A:74:ARG:HE	1:A:84:GLU:HG2	1.76	0.51
1:A:130:PHE:HA	1:A:133:ILE:HD12	1.93	0.50
1:A:166:ARG:NH2	2:C:6:PRO:O	2.45	0.50
1:A:112:LYS:NZ	1:A:128:ASP:OD1	2.46	0.48
1:B:28:LEU:HD13	1:B:86:THR:HG22	1.95	0.48
1:B:29:TRP:HA	1:B:82:PHE:CE1	2.49	0.48
1:A:191:VAL:HG21	1:B:191:VAL:HG22	1.96	0.47
1:A:109:MET:HB3	1:A:145:GLN:HB3	1.96	0.46
1:A:153:GLN:HA	1:A:153:GLN:OE1	2.15	0.46
1:A:135:ILE:HA	1:A:146:ILE:HD11	1.96	0.46
1:B:75:GLU:CG	1:B:77:LYS:HZ2	2.29	0.46
1:A:141:GLN:HE22	1:A:143:ARG:HB2	1.82	0.45
1:A:141:GLN:NE2	1:A:143:ARG:HB2	2.32	0.45
1:A:182:GLN:O	1:A:186:MET:HE2	2.17	0.44
1:A:139:ASP:HB2	1:A:146:ILE:HD13	1.99	0.44
1:B:98:VAL:HG11	1:B:122:PHE:HZ	1.81	0.44
1:A:165:PHE:CZ	1:A:176:ILE:HG23	2.53	0.43
1:B:64:THR:OG1	1:B:153:GLN:CG	2.66	0.43
1:A:122:PHE:O	2:C:6:PRO:HG3	2.17	0.43
1:B:60:PHE:O	1:B:61:ASN:C	2.56	0.43
1:B:165:PHE:CZ	1:B:176:ILE:HG23	2.54	0.43
1:B:66:ARG:HB3	5:B:1008:HOH:O	2.19	0.42
1:A:177:GLN:HB2	1:B:177:GLN:HG3	2.01	0.41
1:B:94:ASP:O	1:B:98:VAL:HG23	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	166/169 (98%)	159 (96%)	7 (4%)	0	100	100
1	B	165/169 (98%)	158 (96%)	7 (4%)	0	100	100
2	C	13/16 (81%)	7 (54%)	5 (38%)	1 (8%)	1	0
2	D	10/16 (62%)	8 (80%)	2 (20%)	0	100	100
All	All	354/370 (96%)	332 (94%)	21 (6%)	1 (0%)	41	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	8	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/153 (100%)	144 (94%)	9 (6%)	19	23
1	B	152/153 (99%)	146 (96%)	6 (4%)	32	41
2	C	12/13 (92%)	12 (100%)	0	100	100
2	D	10/13 (77%)	10 (100%)	0	100	100
All	All	327/332 (98%)	312 (95%)	15 (5%)	27	34

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ASP
1	A	46	THR
1	A	76	ASN
1	A	113	ASN
1	A	137	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	141	GLN
1	A	143	ARG
1	A	172	GLN
1	A	187	VAL
1	B	25	GLN
1	B	39	ARG
1	B	45	ASP
1	B	66	ARG
1	B	104	ARG
1	B	137	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	49	GLN
1	A	76	ASN
1	A	141	GLN
1	A	145	GLN
1	A	159	GLN
1	A	182	GLN
1	B	25	GLN
1	B	49	GLN
1	B	50	GLN
1	B	141	GLN
1	B	145	GLN
1	B	172	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	168/169 (99%)	0.22	7 (4%) 36 34	25, 38, 53, 63	0
1	B	167/169 (98%)	-0.01	6 (3%) 42 41	21, 34, 54, 76	0
2	C	15/16 (93%)	0.20	0 100 100	34, 39, 50, 55	0
2	D	12/16 (75%)	1.18	3 (25%) 0 0	23, 31, 55, 57	0
All	All	362/370 (97%)	0.14	16 (4%) 34 32	21, 36, 53, 76	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	143	ARG	4.0
1	B	25	GLN	3.8
1	A	141	GLN	3.6
1	A	136	ARG	3.6
1	A	39	ARG	3.6
1	B	28	LEU	3.5
1	A	104	ARG	3.2
2	D	12	PRO	3.0
1	B	27	PHE	2.7
1	B	74	ARG	2.7
1	A	38	ASP	2.5
1	B	191	VAL	2.4
2	D	11	TYR	2.2
1	A	113	ASN	2.2
1	B	26	SER	2.1
2	D	10	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NA	A	995	1/1	0.91	0.07	33,33,33,33	0
3	ZN	A	992	1/1	0.97	0.12	50,50,50,50	0
3	ZN	A	993	1/1	0.97	0.10	49,49,49,49	0
3	ZN	A	994	1/1	0.97	0.04	55,55,55,55	0
3	ZN	A	991	1/1	0.97	0.07	53,53,53,53	0
4	NA	B	1	1/1	0.98	0.07	26,26,26,26	0
3	ZN	B	994	1/1	0.99	0.10	46,46,46,46	0
3	ZN	B	992	1/1	0.99	0.07	35,35,35,35	0
3	ZN	B	993	1/1	0.99	0.13	37,37,37,37	0
3	ZN	B	991	1/1	1.00	0.10	41,41,41,41	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.