



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 03:29 AM BST

PDB ID : 5ZO2  
Title : Crystal structure of mouse nectin-like molecule 4 (mNectl-4) full ectodomain in complex with mouse nectin-like molecule 1 (mNectl-1) Ig1 domain, 3.3Å  
Authors : Liu, X.; An, T.; Li, D.; Fan, Z.; Xiang, P.; Li, C.; Ju, W.; Li, J.; Hu, G.; Qin, B.; Yin, B.; Wojdyla, J.A.; Wang, M.; Yuan, J.; Qiang, B.; Shu, P.; Cui, S.; Peng, X.  
Deposited on : 2018-04-12  
Resolution : 3.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>%</div> <div> <div></div> <div>44%</div> <div>21%</div> <div>34%</div> </div> </div>
1	C	299	<div> <div>%</div> <div> <div></div> <div>64%</div> <div>33%</div> <div>..</div> </div> </div>
2	B	147	<div> <div>%</div> <div> <div></div> <div>50%</div> <div>18%</div> <div>32%</div> </div> </div>
3	D	3	<div> <div></div> <div> <div>33%</div> <div>67%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell adhesion molecule 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1501	928	279	290	4			
1	C	293	Total	C	N	O	S	0	0	0
			2283	1413	421	443	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	GLN	ASN	engineered mutation	UNP Q8R464
A	262	GLN	ASN	engineered mutation	UNP Q8R464
A	286	GLN	ASN	engineered mutation	UNP Q8R464
A	318	HIS	-	expression tag	UNP Q8R464
A	319	HIS	-	expression tag	UNP Q8R464
A	320	HIS	-	expression tag	UNP Q8R464
A	321	HIS	-	expression tag	UNP Q8R464
A	322	HIS	-	expression tag	UNP Q8R464
A	323	HIS	-	expression tag	UNP Q8R464
C	31	GLN	ASN	engineered mutation	UNP Q8R464
C	262	GLN	ASN	engineered mutation	UNP Q8R464
C	286	GLN	ASN	engineered mutation	UNP Q8R464
C	318	HIS	-	expression tag	UNP Q8R464
C	319	HIS	-	expression tag	UNP Q8R464
C	320	HIS	-	expression tag	UNP Q8R464
C	321	HIS	-	expression tag	UNP Q8R464
C	322	HIS	-	expression tag	UNP Q8R464
C	323	HIS	-	expression tag	UNP Q8R464


- Molecule 2 is a protein called Cell adhesion molecule 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			769	479	131	156	3			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	GLY	-	expression tag	UNP Q99N28
B	-12	SER	-	expression tag	UNP Q99N28
B	-11	GLY	-	expression tag	UNP Q99N28
B	-10	MET	-	expression tag	UNP Q99N28
B	-9	LYS	-	expression tag	UNP Q99N28
B	-8	GLU	-	expression tag	UNP Q99N28
B	-7	THR	-	expression tag	UNP Q99N28
B	-6	ALA	-	expression tag	UNP Q99N28
B	-5	ALA	-	expression tag	UNP Q99N28
B	-4	ALA	-	expression tag	UNP Q99N28
B	-3	LYS	-	expression tag	UNP Q99N28
B	-2	PHE	-	expression tag	UNP Q99N28
B	-1	GLU	-	expression tag	UNP Q99N28
B	0	ARG	-	expression tag	UNP Q99N28
B	1	GLN	-	expression tag	UNP Q99N28
B	2	HIS	-	expression tag	UNP Q99N28
B	3	MET	-	expression tag	UNP Q99N28
B	4	ASP	-	expression tag	UNP Q99N28
B	5	SER	-	expression tag	UNP Q99N28
B	6	PRO	-	expression tag	UNP Q99N28
B	7	ASP	-	expression tag	UNP Q99N28
B	8	LEU	-	expression tag	UNP Q99N28
B	9	GLY	-	expression tag	UNP Q99N28
B	10	THR	-	expression tag	UNP Q99N28
B	11	ASP	-	expression tag	UNP Q99N28
B	12	ASP	-	expression tag	UNP Q99N28
B	13	ASP	-	expression tag	UNP Q99N28
B	14	ASP	-	expression tag	UNP Q99N28
B	15	LYS	-	expression tag	UNP Q99N28
B	16	ALA	-	expression tag	UNP Q99N28
B	17	MET	-	expression tag	UNP Q99N28
B	18	ALA	-	expression tag	UNP Q99N28
B	19	ASP	-	expression tag	UNP Q99N28
B	20	ILE	-	expression tag	UNP Q99N28
B	21	GLY	-	expression tag	UNP Q99N28
B	22	SER	-	expression tag	UNP Q99N28

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	38	22	2	14	0	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Cell adhesion molecule 4




#### • Molecule 1: Cell adhesion molecule 4



#### • Molecule 2: Cell adhesion molecule 3





- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	207.48 Å   207.48 Å   52.94 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	49.84 – 3.29 49.84 – 3.29	Depositor EDS
% Data completeness (in resolution range)	97.2 (49.84-3.29) 97.2 (49.84-3.29)	Depositor EDS
$R_{merge}$	0.22	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 3.33 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, $R_{free}$	0.243 , 0.302 0.251 , 0.302	Depositor DCC
$R_{free}$ test set	1003 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.0	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 27.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.045 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	4591	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/1528	0.91	2/2081 (0.1%)
1	C	0.58	0/2323	0.67	2/3158 (0.1%)
2	B	0.53	0/782	0.62	0/1065
All	All	0.56	0/4633	0.75	4/6304 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	274	GLU	OE1-CD-OE2	-27.88	89.85	123.30
1	A	274	GLU	CG-CD-OE2	13.37	145.03	118.30
1	C	110	ASP	CB-CG-OD2	5.15	122.93	118.30
1	C	303	LEU	CA-CB-CG	5.03	126.88	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	274	GLU	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1501	0	1482	56	3
1	C	2283	0	2238	82	0
2	B	769	0	756	23	0
3	D	38	0	34	2	0
All	All	4591	0	4510	159	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83:SER:O	1:C:85:ARG:N	1.83	1.12
1:C:75:GLU:OE2	1:C:78:GLN:NE2	1.87	1.07
1:A:138:GLY:O	1:A:187:ARG:NH1	1.92	1.02
1:A:140:GLU:OE1	1:A:185:ARG:NH1	1.93	0.99
1:C:99:GLU:OE1	1:C:176:LYS:NZ	1.95	0.97

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:GLU:CD	1:A:274:GLU:OE2[5_6710]	1.24	0.96
1:A:274:GLU:OE1	1:A:274:GLU:OE2[5_6710]	1.76	0.44
1:A:244:THR:OG1	1:A:244:THR:OG1[5_6710]	2.11	0.09

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/299 (65%)	174 (90%)	19 (10%)	1 (0%)	29	61
1	C	291/299 (97%)	256 (88%)	31 (11%)	4 (1%)	11	38
2	B	98/147 (67%)	91 (93%)	7 (7%)	0	100	100
All	All	583/745 (78%)	521 (89%)	57 (10%)	5 (1%)	17	48

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	84	PRO
1	A	125	GLU
1	C	193	ASP
1	C	207	PRO
1	C	279	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/251 (64%)	161 (100%)	0	100	100
1	C	245/251 (98%)	244 (100%)	1 (0%)	91	95
2	B	89/127 (70%)	89 (100%)	0	100	100
All	All	495/629 (79%)	494 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	82	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	GLN
1	A	174	ASN
1	C	57	GLN
2	B	89	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	D	1	1,3	14,14,15	0.47	0	17,19,21	1.04	1 (5%)
3	NAG	D	2	3	14,14,15	1.14	1 (7%)	17,19,21	2.70	5 (29%)
3	FUC	D	3	3	10,10,11	1.43	1 (10%)	14,14,16	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	3/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	NAG	C1-C2	4.07	1.58	1.52
3	D	3	FUC	O5-C1	-4.02	1.37	1.43

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	NAG	O4-C4-C5	-7.47	90.76	109.30
3	D	2	NAG	C1-O5-C5	4.82	118.72	112.19
3	D	2	NAG	C2-N2-C7	3.79	128.31	122.90
3	D	2	NAG	O4-C4-C3	-3.52	102.21	110.35
3	D	1	NAG	O4-C4-C3	3.26	117.89	110.35

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	1	NAG	O5-C5-C6-O6
3	D	2	NAG	C1-C2-N2-C7
3	D	2	NAG	C4-C5-C6-O6
3	D	1	NAG	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1	NAG	2	0
3	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	196/299 (65%)	0.08	4 (2%) 65 64	31, 68, 104, 142	0
1	C	293/299 (97%)	-0.08	3 (1%) 82 82	36, 58, 85, 106	0
2	B	100/147 (68%)	0.02	1 (1%) 82 82	50, 65, 84, 104	0
All	All	589/745 (79%)	-0.01	8 (1%) 75 75	31, 62, 95, 142	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	167	GLY	3.7
1	A	208	SER	3.4
1	A	209	GLY	3.0
1	A	201	ALA	2.7
1	C	79	LEU	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	D	2	14/15	0.86	0.25	87,100,107,108	0
3	FUC	D	3	10/11	0.89	0.24	93,97,107,111	0
3	NAG	D	1	14/15	0.92	0.15	80,89,97,99	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.