



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:34 am BST

PDB ID : 5ZPU
Title : LFS829 in complex with CRM1-Ran-RanBP1
Authors : Sun, Q.; Lei, Y.
Deposited on : 2018-04-16
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
2	B	140	<div> <div>2%</div> <div>83%</div> <div>14%</div> </div>
3	C	1024	<div> <div>2%</div> <div>95%</div> <div></div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	1	0
			1640	1056	282	296	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	ALA	LEU	engineered mutation	UNP P62826

- Molecule 2 is a protein called Ran-specific GTPase-activating protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	120	Total	C	N	O	S	0	0	0
			983	623	173	183	4			


- Molecule 3 is a protein called Exportin-1,Exportin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	1015	Total	C	N	O	S	0	4	0
			8210	5279	1349	1541	41			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P30822
C	-1	GLY	-	expression tag	UNP P30822
C	0	SER	-	expression tag	UNP P30822
C	537	GLY	ASP	engineered mutation	UNP P30822
C	539	CYS	THR	engineered mutation	UNP P30822
C	540	GLU	VAL	engineered mutation	UNP P30822
C	541	GLN	LYS	engineered mutation	UNP P30822
C	1022	CYS	TYR	engineered mutation	UNP P30822

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $\text{C}_{10}\text{H}_{16}\text{N}_5\text{O}_{14}\text{P}_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0

- Molecule 6 is (Z)-[3E]-4-[(R)-[3,5-bis(trifluoromethyl)phenyl]sulfinyl]but-3-en-1-yl]imino}methanethiol (three-letter code: D29) (formula: C₁₃H₁₁F₆NOS₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	C	1	Total	C	F	N	O	S	0	0
			23	13	6	1	1	2		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	53	Total	O	0	0
			53	53		
8	B	17	Total	O	0	0
			17	17		
8	C	240	Total	O	0	0
			240	240		



3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTP-binding nuclear protein Ran



- Molecule 2: Ran-specific GTPase-activating protein 1



- Molecule 3: Exportin-1,Exportin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.44Å 105.44Å 303.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.60 49.81 – 2.59	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.01-2.60) 99.8 (49.81-2.59)	Depositor EDS
R_{merge}	0.31	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.225 , 0.259 0.223 , 0.261	Depositor DCC
R_{free} test set	2743 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 16.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11200	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D29, GTP, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1679	0.64	0/2272
2	B	0.45	0/1000	0.67	0/1335
3	C	0.45	0/8379	0.63	0/11355
All	All	0.45	0/11058	0.63	0/14962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1640	0	1635	5	0
2	B	983	0	986	2	0
3	C	8210	0	8312	23	0
4	A	32	0	12	0	0
5	A	1	0	0	0	0
6	C	23	0	0	1	0
7	C	1	0	0	0	0
8	A	53	0	0	0	0
8	B	17	0	0	0	0
8	C	240	0	0	2	0
All	All	11200	0	10945	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:345:GLN:NE2	3:C:426:GLN:HE22	1.83	0.77
3:C:268:ASN:HD22	3:C:271:ILE:HD12	1.69	0.57
3:C:345:GLN:NE2	3:C:426:GLN:NE2	2.52	0.56
1:A:210:PRO:O	2:B:177:ARG:NH2	2.42	0.53
3:C:345:GLN:HE21	3:C:426:GLN:HE22	1.53	0.51
3:C:661:ARG:HD2	8:C:1218:HOH:O	2.10	0.51
3:C:436:VAL:HG11	3:C:464:LEU:HD23	1.93	0.50
3:C:686:ASN:HB2	3:C:687:PRO:HA	1.95	0.49
3:C:736:SER:OG	3:C:798:ASP:OD1	2.31	0.49
1:A:77:ASP:HA	1:A:80:TYR:CE2	2.48	0.49
3:C:190:PHE:CZ	3:C:224:ILE:HG21	2.48	0.48
2:B:155:SER:OG	2:B:177:ARG:HG3	2.15	0.47
3:C:600:ILE:O	3:C:604:GLN:HG3	2.16	0.46
3:C:345:GLN:HE22	3:C:426:GLN:NE2	2.13	0.46
3:C:526:ARG:NH2	8:C:1203:HOH:O	2.41	0.46
1:A:81:ILE:HD11	3:C:65:PHE:CD1	2.51	0.46
3:C:345:GLN:HE22	3:C:426:GLN:CD	2.19	0.46
1:A:85:CYS:HB2	1:A:164:LEU:HD22	1.98	0.46
1:A:117:ILE:HB	1:A:144:LEU:HD22	1.99	0.45
3:C:85:ASP:HA	3:C:88:ILE:HG22	1.99	0.44
3:C:859:PHE:CZ	3:C:863:LEU:HD11	2.55	0.42
3:C:686:ASN:HB3	3:C:688:THR:N	2.35	0.42
3:C:90:ILE:HA	3:C:93:PHE:CE2	2.55	0.41
3:C:293[A]:VAL:CG1	3:C:345:GLN:HB3	2.50	0.41
3:C:17:ASP:OD2	3:C:59:THR:HG22	2.20	0.41
3:C:548:LYS:HD2	6:C:1101:D29:N	2.36	0.41
3:C:707:VAL:O	3:C:711:THR:HG23	2.21	0.41
3:C:55:LEU:HG	3:C:97:MET:HE2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/216 (93%)	195 (98%)	5 (2%)	0	100	100
2	B	118/140 (84%)	111 (94%)	7 (6%)	0	100	100
3	C	1017/1024 (99%)	988 (97%)	28 (3%)	1 (0%)	51	75
All	All	1335/1380 (97%)	1294 (97%)	40 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	686	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/184 (96%)	176 (100%)	0	100	100
2	B	102/121 (84%)	100 (98%)	2 (2%)	55	78
3	C	931/933 (100%)	920 (99%)	11 (1%)	71	87
All	All	1209/1238 (98%)	1196 (99%)	13 (1%)	73	88

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	82	ASP
2	B	168	GLU
3	C	35	GLN
3	C	84	ASN
3	C	97	MET
3	C	262	LEU
3	C	273	ARG
3	C	291	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	623	GLN
3	C	940	PHE
3	C	954	LYS
3	C	988	GLN
3	C	991	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN
2	B	137	ASN
3	C	44	ASN
3	C	84	ASN
3	C	268	ASN
3	C	345	GLN
3	C	477	HIS
3	C	742	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GTP	A	301	5	26,34,34	1.19	3 (11%)	33,54,54	1.81	8 (24%)
6	D29	C	1101	3	19,23,23	3.91	1 (5%)	30,33,33	2.95	13 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	301	5	-	1/18/38/38	0/3/3/3
6	D29	C	1101	3	-	7/22/23/23	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1101	D29	C4-S1	-16.74	1.64	1.79
4	A	301	GTP	C6-C5	3.92	1.48	1.41
4	A	301	GTP	C5-C4	2.75	1.48	1.40
4	A	301	GTP	O4'-C1'	2.25	1.44	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1101	D29	O-S1-C4	-9.24	95.56	106.14
6	C	1101	D29	C3-C4-S1	5.94	127.39	119.10
6	C	1101	D29	C4-S1-C9	5.91	105.64	97.50
6	C	1101	D29	C5-C4-S1	-4.36	113.01	119.10
4	A	301	GTP	C5-C6-N1	-4.21	117.67	123.43
4	A	301	GTP	C6-N1-C2	4.18	122.58	115.93
6	C	1101	D29	F1-C8-C2	-3.90	104.36	112.93
6	C	1101	D29	F5-C7-C6	-3.82	104.54	112.93
4	A	301	GTP	C6-C5-C4	-3.76	117.20	120.80
4	A	301	GTP	C2-N3-C4	3.58	119.45	115.36
6	C	1101	D29	C12-N-C13	2.97	125.12	118.08
4	A	301	GTP	PA-O3A-PB	-2.85	123.05	132.83
4	A	301	GTP	N3-C2-N1	-2.82	123.46	127.22
6	C	1101	D29	C6-C5-C4	-2.60	119.14	121.62
4	A	301	GTP	PB-O3B-PG	-2.37	124.68	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1101	D29	F5-C7-F4	2.31	114.18	105.72
6	C	1101	D29	C11-C10-C9	-2.18	121.53	125.51
4	A	301	GTP	C3'-C2'-C1'	2.08	104.11	100.98
6	C	1101	D29	C5-C6-C1	2.07	123.03	118.66
6	C	1101	D29	C2-C3-C4	-2.05	119.66	121.62
6	C	1101	D29	C3-C2-C8	2.03	122.32	119.58

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	C	1101	D29	C5-C4-S1-O
6	C	1101	D29	C5-C4-S1-C9
6	C	1101	D29	C3-C4-S1-O
6	C	1101	D29	C3-C4-S1-C9
6	C	1101	D29	C10-C9-S1-O
6	C	1101	D29	C11-C12-N-C13
6	C	1101	D29	C10-C11-C12-N
4	A	301	GTP	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	1101	D29	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/216 (93%)	-0.13	7 (3%) 45 38	36, 51, 88, 134	0
2	B	120/140 (85%)	0.04	3 (2%) 57 51	45, 63, 85, 119	0
3	C	1015/1024 (99%)	-0.02	19 (1%) 66 62	32, 57, 92, 135	0
All	All	1338/1380 (96%)	-0.03	29 (2%) 62 56	32, 57, 91, 135	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	ALA	5.8
3	C	205	SER	4.4
3	C	690	LEU	4.0
1	A	193	LEU	3.9
3	C	971	ILE	3.9
3	C	967	TYR	3.7
3	C	204	GLY	3.3
3	C	1	MET	3.2
3	C	229	ILE	3.2
3	C	972	SER	3.2
1	A	181	ALA	3.2
2	B	158	TYR	3.0
3	C	500	TRP	2.9
1	A	182	ALA	2.9
3	C	201	LEU	2.6
3	C	230	TYR	2.6
2	B	124	ILE	2.5
3	C	978	GLU	2.5
3	C	687	PRO	2.5
3	C	541	GLN	2.3
1	A	197	TYR	2.3
3	C	966	VAL	2.3
1	A	47	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	591	VAL	2.1
2	B	125	LEU	2.1
3	C	89	GLY	2.1
3	C	979	ALA	2.1
1	A	185	PRO	2.0
3	C	545	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	CL	C	1102	1/1	0.88	0.13	80,80,80,80	0
6	D29	C	1101	23/23	0.94	0.43	81,94,101,104	0
4	GTP	A	301	32/32	0.98	0.13	41,45,49,52	0
5	MG	A	302	1/1	0.99	0.11	42,42,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D29 C 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.