



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 03:25 pm BST

PDB ID : 2ZQK  
Title : Crystal structure of intimin-Tir68 complex  
Authors : Ma, Y.; Gao, F.; Li, D.-F.; Gao, G.F.  
Deposited on : 2008-08-12  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	<div> <div></div> <div>57% 37% . .</div> </div>
1	B	189	<div> <div>12%</div> <div>47% 43% 7% .</div> </div>
2	C	77	<div> <div>12%</div> <div>31% 48% . 17%</div> </div>
2	D	77	<div> <div>6%</div> <div>36% 39% 6% 18%</div> </div>
2	M	77	<div> <div></div> <div>49% 30% . 17%</div> </div>
2	N	77	<div> <div></div> <div>44% 31% 6% 18%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intimin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	0	0	0
			1410	890	231	284	5			
1	B	184	Total	C	N	O	S	0	0	0
			1417	894	232	286	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP P43261
B	0	MET	-	INITIATING METHIONINE	UNP P43261

- Molecule 2 is a protein called Putative translocated intimin receptor protein (Translocated intimin receptor Tir).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	M	64	Total	C	N	O	0	0	0
			481	289	84	108			
2	N	63	Total	C	N	O	0	0	0
			475	286	83	106			
2	C	64	Total	C	N	O	0	0	0
			481	289	84	108			
2	D	63	Total	C	N	O	0	0	0
			475	286	83	106			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	0	MET	-	INITIATING METHIONINE	UNP Q7DB77
M	69	LEU	-	EXPRESSION TAG	UNP Q7DB77
M	70	GLU	-	EXPRESSION TAG	UNP Q7DB77
M	71	HIS	-	EXPRESSION TAG	UNP Q7DB77
M	72	HIS	-	EXPRESSION TAG	UNP Q7DB77

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
M	73	HIS	-	EXPRESSION TAG	UNP Q7DB77
M	74	HIS	-	EXPRESSION TAG	UNP Q7DB77
M	75	HIS	-	EXPRESSION TAG	UNP Q7DB77
M	76	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	0	MET	-	INITIATING METHIONINE	UNP Q7DB77
N	69	LEU	-	EXPRESSION TAG	UNP Q7DB77
N	70	GLU	-	EXPRESSION TAG	UNP Q7DB77
N	71	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	72	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	73	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	74	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	75	HIS	-	EXPRESSION TAG	UNP Q7DB77
N	76	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	0	MET	-	INITIATING METHIONINE	UNP Q7DB77
C	69	LEU	-	EXPRESSION TAG	UNP Q7DB77
C	70	GLU	-	EXPRESSION TAG	UNP Q7DB77
C	71	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	72	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	73	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	74	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	75	HIS	-	EXPRESSION TAG	UNP Q7DB77
C	76	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	0	MET	-	INITIATING METHIONINE	UNP Q7DB77
D	69	LEU	-	EXPRESSION TAG	UNP Q7DB77
D	70	GLU	-	EXPRESSION TAG	UNP Q7DB77
D	71	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	72	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	73	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	74	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	75	HIS	-	EXPRESSION TAG	UNP Q7DB77
D	76	HIS	-	EXPRESSION TAG	UNP Q7DB77



- Molecule 1: Intimin





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.68Å 42.31Å 182.08Å 90.00° 101.13° 90.00°	Depositor
Resolution (Å)	34.10 – 2.80 34.09 – 2.80	Depositor EDS
% Data completeness (in resolution range)	91.3 (34.10-2.80) 92.9 (34.09-2.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.51 (at 2.81Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.241 , 0.300 0.245 , 0.302	Depositor DCC
$R_{free}$ test set	2014 reflections (9.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.5	Xtriage
Anisotropy	1.109	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 60.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4739	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1439	0.74	0/1956
1	B	0.36	0/1446	0.63	0/1966
2	C	0.30	0/483	0.57	0/653
2	D	0.33	0/477	0.57	0/645
2	M	0.38	0/483	0.62	0/653
2	N	0.41	0/477	0.62	0/645
All	All	0.39	0/4805	0.65	0/6518

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1410	0	1379	56	0
1	B	1417	0	1386	93	0
2	C	481	0	462	42	0
2	D	475	0	457	34	0
2	M	481	0	462	25	0
2	N	475	0	457	25	0
All	All	4739	0	4603	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.



The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:GLU:HG2	1:A:161:SER:HA	1.50	0.93
1:B:96:MET:HG2	1:B:97:ILE:H	1.35	0.90
2:D:28:VAL:HG11	2:D:36:ALA:HB1	1.56	0.87
1:B:172:LEU:HD13	2:D:36:ALA:HB2	1.57	0.84
2:D:22:ASN:HB3	2:D:25:ASN:HD22	1.43	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/189 (96%)	151 (83%)	26 (14%)	4 (2%)	6	22
1	B	182/189 (96%)	148 (81%)	26 (14%)	8 (4%)	2	8
2	C	62/77 (80%)	49 (79%)	12 (19%)	1 (2%)	9	31
2	D	61/77 (79%)	56 (92%)	3 (5%)	2 (3%)	4	13
2	M	62/77 (80%)	55 (89%)	7 (11%)	0	100	100
2	N	61/77 (79%)	52 (85%)	7 (12%)	2 (3%)	4	13
All	All	609/686 (89%)	511 (84%)	81 (13%)	17 (3%)	5	17

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	177	VAL
1	A	110	SER
1	B	133	ALA
1	B	150	LYS
2	D	12	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/163 (97%)	146 (92%)	12 (8%)	13	36
1	B	159/163 (98%)	149 (94%)	10 (6%)	18	46
2	C	51/61 (84%)	49 (96%)	2 (4%)	32	66
2	D	50/61 (82%)	45 (90%)	5 (10%)	7	22
2	M	51/61 (84%)	47 (92%)	4 (8%)	12	35
2	N	50/61 (82%)	47 (94%)	3 (6%)	19	48
All	All	519/570 (91%)	483 (93%)	36 (7%)	15	41

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	17	LYS
1	B	17	ASP
2	D	44	ASP
2	N	64	GLN
1	B	21	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	N	35	ASN
1	B	33	GLN
2	D	35	ASN
2	N	64	GLN
1	B	21	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	183/189 (96%)	-0.19	0 <span>100</span> <span>100</span>	31, 54, 80, 97	0
1	B	184/189 (97%)	0.04	1 (0%) <span>91</span> <span>88</span>	50, 80, 109, 126	0
2	C	64/77 (83%)	0.76	9 (14%) <span>2</span> <span>1</span>	73, 107, 136, 153	0
2	D	63/77 (81%)	0.60	5 (7%) <span>12</span> <span>7</span>	70, 102, 130, 140	0
2	M	64/77 (83%)	-0.31	0 <span>100</span> <span>100</span>	42, 61, 85, 91	0
2	N	63/77 (81%)	-0.29	0 <span>100</span> <span>100</span>	39, 58, 102, 122	0
All	All	621/686 (90%)	0.03	15 (2%) <span>59</span> <span>49</span>	31, 71, 121, 153	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	63	GLN	5.5
2	C	66	ILE	4.8
2	C	64	GLN	3.3
2	C	60	GLU	3.2
2	C	65	ALA	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.