



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:27 am BST

PDB ID : 5ZQF  
Title : Crystal structure of human topoisomerase II beta in complex with 5-iodouridine-containing-DNA in space group P3221  
Authors : Chen, S.F.; Wang, Y.R.; Chan, N.L.  
Deposited on : 2018-04-18  
Resolution : 3.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1026 (4.12-3.64)
Clashscore	141614	1045 (4.10-3.66)
Ramachandran outliers	138981	1008 (4.10-3.66)
Sidechain outliers	138945	1001 (4.10-3.66)
RSRZ outliers	127900	1213 (4.16-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	803	
2	B	8	
3	C	9	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IU	C	20	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 2-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	708	Total	C	N	O	S	0	6	0
			5380	3462	906	989	23			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	419	MET	-	expression tag	UNP Q02880
A	420	ALA	-	expression tag	UNP Q02880
A	421	SER	-	expression tag	UNP Q02880
A	422	TRP	-	expression tag	UNP Q02880
A	423	SER	-	expression tag	UNP Q02880
A	424	HIS	-	expression tag	UNP Q02880
A	425	PRO	-	expression tag	UNP Q02880
A	426	GLN	-	expression tag	UNP Q02880
A	427	PHE	-	expression tag	UNP Q02880
A	428	GLU	-	expression tag	UNP Q02880
A	429	LYS	-	expression tag	UNP Q02880
A	430	GLY	-	expression tag	UNP Q02880
A	431	ALA	-	expression tag	UNP Q02880
A	432	ASP	-	expression tag	UNP Q02880
A	433	ASP	-	expression tag	UNP Q02880
A	434	ASP	-	expression tag	UNP Q02880
A	435	ASP	-	expression tag	UNP Q02880
A	436	LYS	-	expression tag	UNP Q02880
A	437	VAL	-	expression tag	UNP Q02880
A	438	PRO	-	expression tag	UNP Q02880
A	439	ASP	-	expression tag	UNP Q02880
A	440	PRO	-	expression tag	UNP Q02880
A	441	THR	-	expression tag	UNP Q02880
A	442	SER	-	expression tag	UNP Q02880
A	443	VAL	-	expression tag	UNP Q02880
A	444	ASP	-	expression tag	UNP Q02880
A	1202	GLY	-	expression tag	UNP Q02880

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	1203	ALA	-	expression tag	UNP Q02880
A	1204	PRO	-	expression tag	UNP Q02880
A	1205	GLY	-	expression tag	UNP Q02880
A	1206	PHE	-	expression tag	UNP Q02880
A	1207	SER	-	expression tag	UNP Q02880
A	1208	SER	-	expression tag	UNP Q02880
A	1209	ILE	-	expression tag	UNP Q02880
A	1210	SER	-	expression tag	UNP Q02880
A	1211	ALA	-	expression tag	UNP Q02880
A	1212	HIS	-	expression tag	UNP Q02880
A	1213	HIS	-	expression tag	UNP Q02880
A	1214	HIS	-	expression tag	UNP Q02880
A	1215	HIS	-	expression tag	UNP Q02880
A	1216	HIS	-	expression tag	UNP Q02880
A	1217	HIS	-	expression tag	UNP Q02880
A	1218	HIS	-	expression tag	UNP Q02880
A	1219	HIS	-	expression tag	UNP Q02880
A	1220	HIS	-	expression tag	UNP Q02880
A	1221	HIS	-	expression tag	UNP Q02880

- Molecule 2 is a DNA chain called DNA (5'-D(P\*AP\*GP\*CP\*CP\*GP\*AP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	8	Total	C	N	O	P	0	0	0
			165	77	34	46	8			

- Molecule 3 is DNA/RNA hybrid called DNA/RNA (5'-D(P\*AP\*GP\*C)-R(P\*(IU))-D(P\*C P\*GP\*GP\*C)-R(P\*(IU))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	I	N	O	P	0	0
			186	85	2	33	57	9		


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mn	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA topoisomerase 2-beta



- Molecule 2: DNA (5'-D(P\*AP\*GP\*CP\*CP\*GP\*AP\*GP\*C)-3')



- Molecule 3: DNA/RNA (5'-D(P\*AP\*GP\*C)-R(P\*(IU))-D(P\*CP\*GP\*GP\*C)-R(P\*(IU))-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.99 Å 94.99 Å 231.49 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.90 – 3.87 19.90 – 3.87	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.90-3.87) 99.8 (19.90-3.87)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.23 (at 3.82 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.203 , 0.247 0.203 , 0.247	Depositor DCC
$R_{free}$ test set	1178 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	105.0	Xtriage
Anisotropy	0.435	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.24 , 64.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.044 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5733	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	129.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, IU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	1/5509 (0.0%)	0.42	0/7480
2	B	0.52	0/185	0.75	0/283
3	C	0.59	0/161	0.75	0/246
All	All	0.29	1/5855 (0.0%)	0.45	0/8009

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	801	GLN	C-N	-6.42	1.22	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5380	0	5072	35	0
2	B	165	0	89	2	0
3	C	186	0	96	1	0
4	A	2	0	0	0	0
All	All	5733	0	5257	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:831:ARG:HH21	1:A:836:ALA:HA	1.25	1.01
1:A:831:ARG:NH2	1:A:836:ALA:HA	2.00	0.73
1:A:944:VAL:HG12	1:A:945:ARG:HG2	1.73	0.71
1:A:608:ILE:HG23	1:A:609:PRO:HD3	1.77	0.66
1:A:678:LYS:NZ	1:A:875:GLY:O	2.30	0.64
1:A:657:ALA:N	1:A:661:ASP:OD2	2.28	0.64
1:A:939:ILE:HB	1:A:984:PHE:HB2	1.86	0.56
1:A:1071:ILE:HG13	1:A:1149:LEU:HD23	1.87	0.56
1:A:560:GLN:HB3	1:A:722:PHE:HA	1.89	0.54
1:A:1171:LEU:HD13	1:A:1174:LYS:HD2	1.91	0.52
3:C:19:DC:H4'	3:C:20:IU:OP1	2.09	0.51
1:A:852:VAL:HG23	1:A:853:GLU:H	1.74	0.51
1:A:558:GLN:HG3	1:A:591:THR:O	2.12	0.50
1:A:739:LYS:HD3	2:B:6:DA:H5''	1.95	0.49
1:A:1072:LEU:HA	1:A:1075:ILE:HG22	1.96	0.48
1:A:457:LEU:HD22	1:A:529:ILE:HG12	1.95	0.48
1:A:670:SER:HB3	1:A:673:LYS:HG2	1.95	0.47
1:A:732:PRO:HG2	1:A:869:ALA:HB1	1.96	0.47
1:A:959:MET:HB3	1:A:970:ILE:HG12	1.97	0.47
1:A:554:ILE:HD12	1:A:566:LYS:HA	1.98	0.46
1:A:751:LYS:HB3	1:A:770:MET:HE1	1.98	0.45
1:A:842:LEU:HD21	1:A:858:ILE:HG22	1.98	0.45
1:A:752:ARG:NH1	1:A:754:ASP:OD2	2.50	0.45
1:A:459:ASP:OD2	1:A:548:ARG:NH2	2.50	0.45
1:A:915:ILE:HG12	1:A:925:VAL:HG22	1.97	0.44
2:B:6:DA:H2''	2:B:7:DG:H5''	1.99	0.44
1:A:952:LYS:HD2	1:A:975:GLU:OE2	2.17	0.44
1:A:906:PRO:HG2	1:A:915:ILE:HG21	1.99	0.43
1:A:805:GLN:OE1	1:A:807:GLY:N	2.52	0.43
1:A:767:VAL:O	1:A:771:SER:OG	2.24	0.43
1:A:555:MET:HE3	1:A:555:MET:HB2	1.85	0.42
1:A:993:LEU:O	1:A:997:GLU:HG2	2.19	0.42
1:A:844:PHE:HA	1:A:854:PRO:HA	2.02	0.41
1:A:1050:LYS:NZ	1:A:1174:LYS:O	2.38	0.41
1:A:1154:VAL:O	1:A:1158:ILE:HG13	2.21	0.41
1:A:562:GLY:O	1:A:566:LYS:HG3	2.20	0.41
1:A:810:LEU:HD12	1:A:948:THR:HB	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	704/803 (88%)	685 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	522/704 (74%)	521 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	979	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	IU	C	15	3,2	14,22,23	3.95	4 (28%)	16,32,35	1.72	3 (18%)
3	IU	C	20	3,2	14,22,23	3.88	4 (28%)	16,32,35	1.80	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IU	C	15	3,2	-	2/5/25/26	0/2/2/2
3	IU	C	20	3,2	-	0/5/25/26	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	15	IU	C6-C5	10.58	1.50	1.38
3	C	20	IU	C6-C5	10.15	1.50	1.38
3	C	20	IU	C4-N3	7.66	1.46	1.33
3	C	15	IU	C4-N3	7.58	1.46	1.33
3	C	20	IU	C2-N3	6.50	1.51	1.38
3	C	15	IU	C2-N3	6.40	1.50	1.38
3	C	20	IU	O4-C4	-2.21	1.19	1.24
3	C	15	IU	O4-C4	-2.20	1.19	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	20	IU	C4-N3-C2	6.07	120.27	115.14
3	C	15	IU	C4-N3-C2	5.75	119.99	115.14
3	C	20	IU	C5-C4-N3	-2.74	119.60	123.27
3	C	15	IU	C6-C5-I5	2.40	120.47	118.52
3	C	15	IU	C5-C4-N3	-2.36	120.11	123.27

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	15	IU	O4'-C1'-N1-C6
3	C	15	IU	C2'-C1'-N1-C6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	20	IU	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	708/803 (88%)	-0.53	4 (0%) 89 84	74, 117, 215, 260	0
2	B	8/8 (100%)	0.28	1 (12%) 3 4	103, 119, 172, 226	0
3	C	7/9 (77%)	0.18	0 100 100	100, 130, 185, 201	0
All	All	723/820 (88%)	-0.52	5 (0%) 87 82	74, 117, 215, 260	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	DA	3.4
1	A	596	ALA	2.7
1	A	591	THR	2.3
1	A	628	TYR	2.3
1	A	613	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IU	C	20	21/22	0.75	0.42	155,166,174,366	0
3	IU	C	15	21/22	0.93	0.18	52,95,118,265	0

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MN	A	1302	1/1	0.91	0.16	126,126,126,126	0
4	MN	A	1301	1/1	0.97	0.06	101,101,101,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.