



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2020 – 03:08 PM BST

PDB ID : 6ZSQ
Title : Crystal structure of the Cisplatin beta-Lactoglobulin adduct formed after 18 h of soaking
Authors : Balasco, N.; Ferraro, G.; Merlino, A.
Deposited on : 2020-07-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	162	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>.</div> </div> </div>
1	BBB	162	<div> <div>6%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NH3	AAA	201	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactoglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	162	Total	C	N	O	S	0	1	0
			1285	821	206	249	9			
1	BBB	161	Total	C	N	O	S	0	3	0
			1295	826	209	251	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	118	VAL	ALA	conflict	UNP P02754
BBB	118	VAL	ALA	conflict	UNP P02754

- Molecule 2 is AMMONIA (three-letter code: NH3) (formula: H₃N) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	1	Total	N	0	0
			1	1		

- Molecule 3 is PLATINUM (II) ION (three-letter code: PT) (formula: Pt) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	BBB	2	Total Pt 3 3	0	1
3	AAA	1	Total Pt 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	BBB	1	Total O S 5 4 1	0	0
4	BBB	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AAA	97	Total O 98 98	0	1
5	BBB	68	Total O 68 68	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactoglobulin



- Molecule 1: Beta-lactoglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.72Å 49.05Å 49.31Å 69.96° 68.59° 77.00°	Depositor
Resolution (Å)	23.23 – 2.00 23.22 – 2.00	Depositor EDS
% Data completeness (in resolution range)	86.9 (23.23-2.00) 86.9 (23.22-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.183 , 0.224 0.187 , 0.232	Depositor DCC
R_{free} test set	894 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2761	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NH3, PT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.68	0/1309	0.88	0/1771
1	BBB	0.68	0/1319	0.86	0/1783
All	All	0.68	0/2628	0.87	0/3554

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1285	0	1313	26	0
1	BBB	1295	0	1318	15	0
2	AAA	1	0	0	1	0
3	AAA	1	0	0	0	0
3	BBB	3	0	0	0	0
4	BBB	10	0	0	0	0
5	AAA	98	0	0	1	0
5	BBB	68	0	0	0	0
All	All	2761	0	2631	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:39:LEU:HD23	1:BBB:118:VAL:HG23	1.69	0.74
1:AAA:100:LYS:HG2	1:AAA:101:LYS:HG3	1.70	0.73
1:AAA:112:GLU:HG3	1:AAA:115:GLN:HG2	1.71	0.73
1:BBB:39:LEU:HB3	1:BBB:118:VAL:HG21	1.70	0.72
1:BBB:39:LEU:HD23	1:BBB:118:VAL:CG2	2.20	0.72
1:AAA:112:GLU:O	1:AAA:116:SER:HB2	1.89	0.72
1:AAA:31:LEU:HD22	1:AAA:39:LEU:HD22	1.72	0.71
1:AAA:7:MET:CE	1:AAA:10:LEU:HD13	2.24	0.68
1:AAA:152:ASN:H	1:AAA:155:GLN:HE21	1.43	0.66
1:BBB:152:ASN:H	1:BBB:155:GLN:HE21	1.41	0.66
1:AAA:7:MET:HE3	1:AAA:10:LEU:HB2	1.81	0.63
1:BBB:39:LEU:CB	1:BBB:118:VAL:HG21	2.31	0.60
1:AAA:88:ASN:ND2	5:AAA:303:HOH:O	2.39	0.55
1:AAA:7:MET:HE1	1:AAA:10:LEU:HD13	1.89	0.54
1:AAA:72:ILE:N	1:AAA:72:ILE:CD1	2.71	0.53
1:BBB:31:LEU:HD22	1:BBB:39:LEU:HD22	1.91	0.52
1:AAA:161:HIS:O	1:AAA:162:ILE:HG12	2.10	0.51
1:AAA:112:GLU:CG	1:AAA:115:GLN:HG2	2.39	0.50
1:AAA:77:LYS:N	1:AAA:77:LYS:HD2	2.29	0.48
1:AAA:33:ASP:O	1:AAA:34:ALA:HB3	2.14	0.47
1:AAA:59:GLN:HE21	1:AAA:68:GLN:HE22	1.61	0.47
1:BBB:152:ASN:H	1:BBB:155:GLN:NE2	2.12	0.47
1:BBB:66:CYS:SG	1:BBB:160:CYS:N	2.89	0.46
1:BBB:7:MET:HB2	1:BBB:96:ASP:HA	1.96	0.46
1:BBB:33:ASP:O	1:BBB:34:ALA:HB3	2.16	0.46
1:AAA:161:HIS:O	1:AAA:162:ILE:CG1	2.64	0.45
1:AAA:155:GLN:O	1:AAA:161:HIS:HA	2.17	0.44
1:BBB:39:LEU:HD23	1:BBB:118:VAL:HG21	1.96	0.44
1:AAA:33:ASP:O	1:AAA:37:ALA:HB2	2.17	0.44
1:AAA:112:GLU:OE2	1:AAA:114:GLU:OE1	2.36	0.43
1:AAA:162:ILE:OXT	1:AAA:162:ILE:HG13	2.18	0.43
1:BBB:138[B]:LYS:HE3	1:BBB:138[B]:LYS:CA	2.49	0.42
1:BBB:138[B]:LYS:HA	1:BBB:138[B]:LYS:HE3	2.00	0.42
1:AAA:72:ILE:HD12	1:AAA:72:ILE:N	2.34	0.42
1:AAA:44:GLU:OE2	1:AAA:68:GLN:OE1	2.38	0.41
1:AAA:7:MET:HE1	1:AAA:10:LEU:CD1	2.49	0.41
1:AAA:96:ASP:OD1	2:AAA:201:NH3:N	2.52	0.41
1:BBB:155:GLN:O	1:BBB:161:HIS:HA	2.20	0.41
1:AAA:59:GLN:HA	1:AAA:67:ALA:O	2.21	0.41
1:BBB:155:GLN:HG2	1:BBB:162:ILE:HB	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:12:ILE:HA	1:AAA:12:ILE:HD12	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	161/162 (99%)	153 (95%)	8 (5%)	0	100	100
1	BBB	162/162 (100%)	153 (94%)	9 (6%)	0	100	100
All	All	323/324 (100%)	306 (95%)	17 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	145/144 (101%)	134 (92%)	11 (8%)	13	8
1	BBB	146/144 (101%)	137 (94%)	9 (6%)	18	13
All	All	291/288 (101%)	271 (93%)	20 (7%)	16	11

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	4	THR
1	AAA	39	LEU
1	AAA	40	ARG
1	AAA	69	LYS
1	AAA	72	ILE
1	AAA	77	LYS
1	AAA	93	LEU
1	AAA	100	LYS
1	AAA	116	SER
1	AAA	138	LYS
1	AAA	161	HIS
1	BBB	39	LEU
1	BBB	62	GLU
1	BBB	66	CYS
1	BBB	68	GLN
1	BBB	69	LYS
1	BBB	131	GLU
1	BBB	150[A]	SER
1	BBB	150[B]	SER
1	BBB	162	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is modelled with single atom and 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	BBB	201	-	4,4,4	0.36	0	6,6,6	0.14	0
4	SO4	BBB	202	-	4,4,4	0.33	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	162/162 (100%)	-0.13	3 (1%) 66 65	9, 18, 42, 69	0
1	BBB	161/162 (99%)	0.09	9 (5%) 24 23	10, 21, 64, 123	0
All	All	323/324 (99%)	-0.02	12 (3%) 41 41	9, 19, 55, 123	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	4	THR	12.9
1	BBB	162	ILE	5.9
1	AAA	63	ASN	5.8
1	BBB	5	GLN	5.7
1	BBB	3	VAL	4.4
1	AAA	162	ILE	3.5
1	BBB	64	GLY	2.7
1	BBB	2	ILE	2.6
1	BBB	161	HIS	2.6
1	BBB	159	GLN	2.3
1	BBB	138[A]	LYS	2.1
1	AAA	159	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NH3	AAA	201	1/1	0.74	0.41	28,28,28,28	1
4	SO4	BBB	202	5/5	0.93	0.21	51,56,59,60	0
4	SO4	BBB	201	5/5	0.93	0.15	45,49,52,62	0
3	PT	BBB	203[A]	1/1	0.95	0.12	46,46,46,46	1
3	PT	BBB	203[B]	1/1	0.95	0.12	56,56,56,56	1
3	PT	BBB	204	1/1	0.98	0.08	46,46,46,46	1
3	PT	AAA	202	1/1	0.99	0.07	39,39,39,39	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NH3 AAA 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PT BBB 203 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PT BBB 203 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PT BBB 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PT AAA 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.