



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 01:43 pm BST

PDB ID : 1ZU5  
Title : Crystal structure of FtsY from Mycoplasma mycoides- space group H32  
Authors : Gariani, T.; Samuelsson, T.; Sauer-Eriksson, A.E.  
Deposited on : 2005-05-30  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div> <div>3%</div> <div> <div></div> <div>76%</div> <div>19%</div> <div>• •</div> </div> </div>
1	B	320	<div> <div>5%</div> <div> <div></div> <div>71%</div> <div>23%</div> <div>• •</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5029 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ftsY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2438	1553	406	467	12			
1	B	310	Total	C	N	O	S	0	0	0
			2452	1561	412	467	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	92	HIS	-	EXPRESSION TAG	GB 1929401
A	93	HIS	-	EXPRESSION TAG	GB 1929401
A	94	HIS	-	EXPRESSION TAG	GB 1929401
A	95	HIS	-	EXPRESSION TAG	GB 1929401
A	96	HIS	-	EXPRESSION TAG	GB 1929401
A	97	PRO	-	EXPRESSION TAG	GB 1929401
A	98	MET	-	EXPRESSION TAG	GB 1929401
A	236	ARG	SER	SEE REMARK 999	GB 1929401
B	92	HIS	-	EXPRESSION TAG	GB 1929401
B	93	HIS	-	EXPRESSION TAG	GB 1929401
B	94	HIS	-	EXPRESSION TAG	GB 1929401
B	95	HIS	-	EXPRESSION TAG	GB 1929401
B	96	HIS	-	EXPRESSION TAG	GB 1929401
B	97	PRO	-	EXPRESSION TAG	GB 1929401
B	98	MET	-	EXPRESSION TAG	GB 1929401
B	236	ARG	SER	SEE REMARK 999	GB 1929401


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	68	Total	O	0	0
			68	68		
2	B	71	Total	O	0	0
			71	71		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ftsY



#### • Molecule 1: ftsY



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	148.44Å 148.44Å 223.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.40 19.84 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.00-2.40) 95.0 (19.84-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.210 , 0.246 0.209 , 0.245	Depositor DCC
$R_{free}$ test set	1836 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.0	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5029	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	0/2469	0.82	1/3317 (0.0%)
1	B	0.85	1/2486 (0.0%)	0.85	4/3341 (0.1%)
All	All	0.82	1/4955 (0.0%)	0.83	5/6658 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	317	SER	CB-OG	6.87	1.51	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	266	ASP	CB-CG-OD1	7.71	125.23	118.30
1	B	236	ARG	NE-CZ-NH1	6.79	123.69	120.30
1	B	236	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	B	266	ASP	CB-CG-OD1	6.05	123.75	118.30
1	B	139	ASP	CB-CG-OD1	5.50	123.25	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2514	56	0
1	B	2452	0	2509	55	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	68	0	0	1	0
2	B	71	0	0	1	0
All	All	5029	0	5023	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:GLU:HG3	1:B:382:GLU:HG3	1.34	1.06
1:B:198:ILE:H	1:B:320:HIS:HD2	1.10	0.99
1:A:198:ILE:H	1:A:320:HIS:HD2	1.15	0.92
1:A:382:GLU:CG	1:B:382:GLU:HG3	2.04	0.86
1:A:175:GLN:HG2	1:A:180:ASN:HD21	1.39	0.86
1:A:296:ASN:HB3	1:A:299:ASN:CG	1.98	0.83
1:A:247:ILE:HD12	1:A:257:LEU:HB2	1.63	0.79
1:A:264:ASN:HD21	1:A:266:ASP:HB3	1.46	0.79
1:B:198:ILE:H	1:B:320:HIS:CD2	2.00	0.76
1:A:247:ILE:CD1	1:A:257:LEU:HB2	2.18	0.73
1:A:198:ILE:H	1:A:320:HIS:CD2	2.06	0.71
1:B:160:THR:HG22	1:B:164:ASN:HD22	1.57	0.70
1:A:175:GLN:HG2	1:A:180:ASN:ND2	2.07	0.69
1:A:382:GLU:HG3	1:B:382:GLU:CG	2.19	0.68
1:A:104:LYS:HE2	1:B:250:ARG:HD2	1.76	0.66
1:B:201:LEU:HD13	1:B:213:LEU:HD12	1.78	0.66
1:A:209:LYS:HE3	1:A:289:ASP:OD2	1.97	0.65
1:A:321:GLU:HG3	1:A:323:LEU:HD21	1.78	0.65
1:B:158:ARG:HD3	1:B:158:ARG:H	1.59	0.64
1:B:383:LYS:N	1:B:386:ASP:OD2	2.29	0.63
1:A:100:LYS:HD3	1:A:380:VAL:HG12	1.79	0.63
1:B:233:ASP:OD2	1:B:236:ARG:HD2	1.98	0.63
1:B:147:LYS:NZ	1:B:369:GLU:OE1	2.32	0.62
1:A:213:LEU:CD2	1:A:287:LEU:HB3	2.29	0.62
1:A:198:ILE:HD13	1:A:275:ILE:HD11	1.82	0.61
1:B:166:LYS:HD2	1:B:403:PHE:HD1	1.65	0.60
1:A:213:LEU:HD11	1:A:289:ASP:HB2	1.82	0.60
1:B:301:MET:HG3	1:B:344:VAL:HG22	1.82	0.60
1:A:101:ALA:HB2	1:A:380:VAL:HG22	1.82	0.59
1:A:156:THR:HG21	1:A:165:ILE:HD13	1.85	0.59
1:B:380:VAL:HG21	1:B:388:LEU:HD21	1.84	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:PHE:CD2	1:A:321:GLU:HB2	2.39	0.58
1:B:198:ILE:N	1:B:320:HIS:HD2	1.91	0.58
1:A:213:LEU:HD22	1:A:287:LEU:HB3	1.85	0.58
1:A:264:ASN:ND2	1:A:266:ASP:HB3	2.15	0.57
1:B:122:GLN:HB3	1:B:158:ARG:O	2.04	0.57
1:A:177:TYR:HB2	1:A:365:LEU:CD2	2.35	0.57
1:B:98:MET:HG3	1:B:394:GLN:HB3	1.87	0.56
1:B:175:GLN:HG3	1:B:180:ASN:HD22	1.69	0.56
1:B:339:GLU:HA	1:B:371:LEU:HD22	1.87	0.56
1:B:175:GLN:HG3	1:B:180:ASN:ND2	2.21	0.55
1:B:199:PHE:HB2	1:B:287:LEU:HD23	1.87	0.55
1:B:182:TRP:C	1:B:183:THR:HG23	2.27	0.55
1:A:213:LEU:CD1	1:A:289:ASP:HB2	2.37	0.55
1:A:380:VAL:HG12	1:A:380:VAL:O	2.06	0.55
1:A:198:ILE:N	1:A:320:HIS:HD2	1.95	0.54
1:B:158:ARG:CD	1:B:158:ARG:H	2.18	0.54
1:B:296:ASN:CG	1:B:299:ASN:HB3	2.27	0.54
1:B:296:ASN:ND2	1:B:299:ASN:HB3	2.25	0.52
1:A:199:PHE:HB2	1:A:287:LEU:HD23	1.92	0.52
1:A:160:THR:HG22	1:A:164:ASN:HB2	1.91	0.52
1:B:339:GLU:HA	1:B:371:LEU:CD2	2.40	0.51
1:B:132:GLU:O	1:B:136:ILE:HG13	2.09	0.51
1:A:271:VAL:HG21	1:A:307:MET:HG2	1.92	0.51
1:A:117:SER:HB3	1:A:162:PHE:CE2	2.46	0.51
1:A:177:TYR:HD1	1:A:365:LEU:HD23	1.76	0.51
1:B:275:ILE:CG2	1:B:314:VAL:HG21	2.41	0.51
1:B:182:TRP:O	1:B:183:THR:HG23	2.11	0.50
1:B:178:THR:O	1:B:179:ASP:HB2	2.10	0.50
1:B:187:TYR:OH	1:B:392:ILE:HG22	2.12	0.50
1:B:275:ILE:HG21	1:B:314:VAL:HG21	1.91	0.50
1:B:97:PRO:O	1:B:380:VAL:HG11	2.11	0.50
1:A:264:ASN:HD22	1:A:266:ASP:H	1.59	0.49
1:B:202:VAL:HB	1:B:341:PHE:CE1	2.48	0.49
1:A:382:GLU:CB	1:B:382:GLU:HG3	2.43	0.49
1:A:197:ASN:HA	1:A:320:HIS:CD2	2.48	0.49
1:B:247:ILE:HD13	1:B:251:LEU:HD12	1.95	0.49
1:A:275:ILE:HG13	1:A:311:ILE:HG22	1.94	0.48
1:A:296:ASN:HB3	1:A:299:ASN:OD1	2.11	0.48
1:A:233:ASP:OD2	1:A:236:ARG:HD2	2.15	0.46
1:A:380:VAL:CG1	1:A:380:VAL:O	2.63	0.46
1:B:101:ALA:HB2	1:B:380:VAL:HG13	1.97	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ILE:HG22	1:A:276:LYS:N	2.31	0.46
1:B:133:ASP:O	1:B:137:GLN:HG2	2.15	0.46
1:A:96:HIS:CD2	1:A:98:MET:HB2	2.50	0.46
1:B:101:ALA:HB2	1:B:380:VAL:CG1	2.45	0.46
1:B:92:HIS:N	2:B:444:HOH:O	2.49	0.46
1:A:252:ASN:OD1	1:A:254:LYS:HE3	2.16	0.46
1:A:157:LYS:HA	1:A:157:LYS:HE2	1.97	0.45
1:A:383:LYS:N	1:A:386:ASP:OD2	2.43	0.45
1:B:162:PHE:O	1:B:165:ILE:HB	2.16	0.45
1:B:178:THR:HB	1:B:181:ASP:O	2.17	0.44
1:B:169:LEU:O	1:B:173:LEU:HG	2.17	0.44
1:A:100:LYS:HD3	1:A:380:VAL:CG1	2.46	0.44
1:B:330:THR:OG1	1:B:334:GLY:HA3	2.17	0.44
1:A:332:GLN:O	1:A:336:ILE:HG13	2.17	0.44
1:B:102:MET:HB2	1:B:398:HIS:HB3	2.00	0.43
1:B:99:GLU:HG2	1:B:398:HIS:HE1	1.83	0.43
1:B:160:THR:HG22	1:B:164:ASN:HB2	2.00	0.43
1:A:163:GLU:HB2	2:A:453:HOH:O	2.19	0.43
1:A:336:ILE:O	1:A:339:GLU:HB2	2.18	0.43
1:A:380:VAL:HG21	1:A:388:LEU:HD21	2.01	0.43
1:A:380:VAL:HG21	1:A:388:LEU:CD2	2.49	0.42
1:B:148:VAL:O	1:B:152:VAL:HG23	2.17	0.42
1:A:381:GLY:N	1:A:386:ASP:HB3	2.35	0.42
1:B:228:LEU:HD11	1:B:258:VAL:HG23	2.02	0.42
1:A:377:MET:HB3	1:A:387:LEU:HD11	2.01	0.42
1:B:114:LYS:O	1:B:117:SER:HB3	2.20	0.41
1:A:339:GLU:OE2	1:A:370:LEU:HB3	2.21	0.41
1:A:275:ILE:CG1	1:A:311:ILE:HG22	2.50	0.41
1:B:166:LYS:HD2	1:B:403:PHE:CD1	2.52	0.41
1:A:96:HIS:HA	1:A:97:PRO:HD3	1.95	0.41
1:A:297:LYS:HB3	1:A:344:VAL:HG21	2.02	0.41
1:B:158:ARG:HD3	1:B:158:ARG:N	2.31	0.40
1:B:355:MET:HG2	1:B:378:ILE:HD11	2.03	0.40
1:B:96:HIS:HB2	1:B:99:GLU:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/320 (96%)	292 (95%)	14 (5%)	1 (0%)	41	55
1	B	306/320 (96%)	291 (95%)	13 (4%)	2 (1%)	22	32
All	All	613/640 (96%)	583 (95%)	27 (4%)	3 (0%)	29	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	313	GLN
1	A	295	GLN
1	B	384	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/277 (96%)	252 (94%)	15 (6%)	21	34
1	B	268/277 (97%)	253 (94%)	15 (6%)	21	34
All	All	535/554 (97%)	505 (94%)	30 (6%)	21	34

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LYS
1	A	119	LYS
1	A	157	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	192	LYS
1	A	222	GLU
1	A	264	ASN
1	A	269	SER
1	A	299	ASN
1	A	301	MET
1	A	321	GLU
1	A	341	PHE
1	A	344	VAL
1	A	363	ILE
1	A	382	GLU
1	A	385	ASP
1	B	99	GLU
1	B	153	ARG
1	B	158	ARG
1	B	192	LYS
1	B	222	GLU
1	B	252	ASN
1	B	253	ASN
1	B	280	GLU
1	B	294	LEU
1	B	299	ASN
1	B	339	GLU
1	B	341	PHE
1	B	382	GLU
1	B	388	LEU
1	B	393	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	150	ASN
1	A	180	ASN
1	A	264	ASN
1	A	320	HIS
1	A	333	ASN
1	B	108	ASN
1	B	150	ASN
1	B	164	ASN
1	B	180	ASN
1	B	320	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	394	GLN
1	B	398	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/320 (96%)	-0.12	10 (3%) 47 46	25, 46, 64, 75	0
1	B	310/320 (96%)	-0.04	16 (5%) 27 26	28, 46, 72, 81	0
All	All	619/640 (96%)	-0.08	26 (4%) 36 35	25, 46, 69, 81	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263	LEU	4.3
1	B	93	HIS	4.2
1	A	180	ASN	3.8
1	A	96	HIS	3.7
1	B	182	TRP	3.6
1	A	264	ASN	3.5
1	A	253	ASN	3.1
1	B	264	ASN	2.9
1	B	119	LYS	2.9
1	A	263	LEU	2.9
1	A	280	GLU	2.8
1	B	316	LYS	2.7
1	B	188	ARG	2.7
1	B	157	LYS	2.7
1	A	313	GLN	2.7
1	B	183	THR	2.6
1	B	95	HIS	2.6
1	A	194	ASN	2.5
1	B	118	LYS	2.4
1	B	286	LEU	2.4
1	B	313	GLN	2.3
1	B	194	ASN	2.2
1	B	295	GLN	2.2
1	A	295	GLN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	286	LEU	2.1
1	B	287	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.