



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 04:49 AM BST

PDB ID : 3ZUX
Title : Crystal structure of a bacterial homologue of the bile acid sodium symporter ASBT.
Authors : Hu, N.-J.; Iwata, S.; Cameron, A.D.; Drew, D.
Deposited on : 2011-07-21
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	<div> <div>8%</div> <div>88%</div> <div>5%</div> <div>7%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2490 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSPORTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2259	1509	355	381	14			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	MET	-	expression tag	UNP Q9K0A9
A	-3	VAL	-	expression tag	UNP Q9K0A9
A	-2	ALA	-	expression tag	UNP Q9K0A9
A	-1	ALA	-	expression tag	UNP Q9K0A9
A	0	SER	-	expression tag	UNP Q9K0A9
A	87	CYS	LEU	engineered mutation	UNP A9M359
A	138	THR	ILE	cloning artifact	UNP Q9K0A9
A	256	ALA	THR	cloning artifact	UNP Q9K0A9
A	316	LEU	-	expression tag	UNP Q9K0A9
A	317	ASP	-	expression tag	UNP Q9K0A9
A	318	ARG	-	expression tag	UNP Q9K0A9
A	319	ALA	-	expression tag	UNP Q9K0A9
A	320	GLY	-	expression tag	UNP Q9K0A9
A	321	SER	-	expression tag	UNP Q9K0A9
A	322	GLU	-	expression tag	UNP Q9K0A9
A	323	ASN	-	expression tag	UNP Q9K0A9
A	324	LEU	-	expression tag	UNP Q9K0A9
A	325	TYR	-	expression tag	UNP Q9K0A9
A	326	PHE	-	expression tag	UNP Q9K0A9
A	327	GLN	-	expression tag	UNP Q9K0A9


- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Na	0	0
			2	2		

- Molecule 3 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Hg 1 1	0	0

- Molecule 4 is TAUROCHOLIC ACID (three-letter code: TCH) (formula: $\text{C}_{26}\text{H}_{45}\text{NO}_7\text{S}$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 35	C 26	N 1	O 7	S 1	0	0
4	A	1	Total 22	C 19	O 3	0			0

- Molecule 5 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: $\text{C}_{14}\text{H}_{31}\text{NO}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 10 8 1 1	0	0
5	A	1	Total C N O 16 14 1 1	0	0
5	A	1	Total C N O 16 14 1 1	0	0
5	A	1	Total C 9 9	0	0
5	A	1	Total C N O 9 7 1 1	0	0

- Molecule 6 is PHOSPHATIDYLETHANOLAMINE (three-letter code: PTY) (formula: $C_{40}H_{80}NO_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			40	30	1	8	1		
6	A	1	Total	C	N	O	P	0	0
			34	24	1	8	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	37	Total	O	0	0
			37	37		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRANSPORTER



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	73.33Å 73.33Å 163.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.64 – 2.20 24.44 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (24.64-2.20) 99.6 (24.44-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.19Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.197 , 0.229 0.208 , 0.243	Depositor DCC
R_{free} test set	1180 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 71.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2490	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, LDA, TCH, PTY, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2313	0.60	0/3158

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2259	0	2400	14	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	57	0	74	4	0
5	A	60	0	106	2	0
6	A	74	0	97	4	0
7	A	37	0	0	0	0
All	All	2490	0	2677	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:PRO:HG2	5:A:1317:LDA:HM12	1.74	0.68
1:A:116:VAL:HG11	4:A:1313:TCH:H22	1.83	0.60
1:A:184:LEU:HB2	1:A:188:THR:HG23	1.85	0.57
1:A:173:PRO:HG3	5:A:1317:LDA:H81	1.87	0.57
1:A:241:PHE:HZ	6:A:1319:PTY:HC22	1.70	0.56
1:A:199:SER:HB3	4:A:1313:TCH:H1	1.94	0.49
1:A:303:THR:HG23	6:A:1318:PTY:HC6	1.95	0.48
1:A:53:LEU:HD21	1:A:191:LEU:HD23	1.97	0.46
1:A:13:LYS:HG2	4:A:1313:TCH:H26A	2.01	0.42
1:A:303:THR:CG2	6:A:1318:PTY:HC6	2.50	0.42
1:A:13:LYS:HE2	4:A:1313:TCH:H26A	2.03	0.41
1:A:297:SER:HB2	6:A:1319:PTY:H171	2.03	0.40
1:A:162:MET:HG2	1:A:272:LEU:HD22	2.04	0.40
1:A:172:LEU:HB3	1:A:173:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/332 (92%)	302 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/253 (92%)	233 (100%)	1 (0%)	91	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	294	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PTY	A	1319	-	33,33,49	1.20	4 (12%)	36,38,54	2.17	8 (22%)
5	LDA	A	1316	-	12,15,15	0.87	1 (8%)	14,17,17	0.73	0
4	TCH	A	1314	-	25,25,38	0.83	0	37,41,60	1.71	12 (32%)
4	TCH	A	1313	-	38,38,38	1.02	2 (5%)	59,60,60	1.77	14 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PTY	A	1318	-	39,39,49	1.08	2 (5%)	42,44,54	2.02	10 (23%)
5	LDA	A	1315	-	6,9,15	0.89	1 (16%)	8,11,17	0.67	0
5	LDA	A	1321	-	5,8,15	1.54	1 (20%)	7,10,17	0.76	0
5	LDA	A	1320	-	8,8,15	0.40	0	7,7,17	0.72	0
5	LDA	A	1317	-	12,15,15	0.85	1 (8%)	14,17,17	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PTY	A	1319	-	-	18/37/37/53	-
5	LDA	A	1316	-	-	6/13/13/13	-
4	TCH	A	1314	-	-	-	0/4/4/4
4	TCH	A	1313	-	-	2/16/81/81	0/4/4/4
6	PTY	A	1318	-	-	21/43/43/53	-
5	LDA	A	1320	-	-	2/6/6/13	-
5	LDA	A	1321	-	-	3/6/6/13	-
5	LDA	A	1315	-	-	2/7/7/13	-
5	LDA	A	1317	-	-	7/13/13/13	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1321	LDA	O1-N1	3.35	1.50	1.42
6	A	1318	PTY	P1-O13	3.32	1.62	1.50
4	A	1313	TCH	C26-S26	3.12	1.81	1.77
6	A	1319	PTY	P1-O13	3.08	1.61	1.50
6	A	1319	PTY	O4-C30	2.84	1.41	1.33
6	A	1318	PTY	O7-C6	-2.80	1.39	1.46
5	A	1317	LDA	O1-N1	2.65	1.48	1.42
5	A	1316	LDA	O1-N1	2.64	1.48	1.42
4	A	1313	TCH	O1S-S26	2.62	1.52	1.45
6	A	1319	PTY	O7-C8	2.29	1.40	1.34
6	A	1319	PTY	O30-C30	-2.02	1.16	1.22
5	A	1315	LDA	O1-N1	2.00	1.46	1.42

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1319	PTY	C6-O7-C8	6.71	134.32	117.79
6	A	1318	PTY	C6-O7-C8	5.34	130.94	117.79
4	A	1313	TCH	O3S-S26-O2S	5.10	123.75	111.27
6	A	1318	PTY	O4-C30-O30	-5.06	110.81	123.59
6	A	1319	PTY	O4-C30-O30	-4.89	111.25	123.59
6	A	1319	PTY	O7-C8-C11	4.64	121.49	111.50
6	A	1319	PTY	O4-C30-C31	4.46	125.91	111.91
4	A	1313	TCH	O3S-S26-O1S	-4.10	101.27	111.27
4	A	1314	TCH	C19-C10-C1	-3.98	101.85	108.26
4	A	1313	TCH	O2S-S26-C26	3.96	111.69	106.92
6	A	1319	PTY	O7-C8-O10	-3.72	114.70	123.70
6	A	1318	PTY	O7-C8-C11	3.64	119.34	111.50
4	A	1313	TCH	C5-C4-C3	-3.62	107.45	112.76
6	A	1318	PTY	O4-C30-C31	3.54	123.03	111.91
4	A	1313	TCH	O2S-S26-O1S	-3.52	101.75	113.95
6	A	1318	PTY	O7-C8-O10	-3.52	115.19	123.70
6	A	1318	PTY	C42-C41-C40	-3.51	96.62	114.42
4	A	1313	TCH	O3S-S26-C26	3.39	111.25	105.77
4	A	1314	TCH	C1-C2-C3	3.25	114.64	110.47
6	A	1318	PTY	O7-C6-C5	3.14	119.79	108.40
6	A	1319	PTY	O7-C6-C5	3.14	119.77	108.40
4	A	1314	TCH	C19-C10-C9	-3.10	106.91	111.18
4	A	1313	TCH	C23-C22-C20	-3.08	108.90	114.52
6	A	1318	PTY	O12-P1-O14	2.96	121.49	107.75
6	A	1319	PTY	O12-P1-O14	2.76	120.55	107.75
6	A	1318	PTY	C40-C39-C38	-2.70	100.71	114.42
4	A	1314	TCH	C9-C8-C7	-2.69	108.65	111.88
4	A	1314	TCH	C14-C8-C9	2.69	113.41	109.71
4	A	1313	TCH	C9-C10-C5	2.63	112.28	108.58
4	A	1313	TCH	C19-C10-C5	-2.54	106.05	110.36
4	A	1314	TCH	C10-C9-C8	-2.53	109.11	111.82
6	A	1318	PTY	O11-C3-C2	-2.52	99.67	109.10
4	A	1314	TCH	C1-C10-C5	2.52	111.49	107.77
4	A	1313	TCH	C13-C17-C20	-2.51	116.50	119.50
4	A	1313	TCH	O1S-S26-C26	-2.40	104.03	106.92
4	A	1314	TCH	C11-C9-C8	2.39	114.37	110.88
4	A	1313	TCH	O7-C7-C8	2.31	114.59	109.43
4	A	1314	TCH	C4-C5-C10	2.26	115.06	112.66
4	A	1314	TCH	C2-C1-C10	2.22	116.59	112.78
4	A	1313	TCH	C1-C10-C5	2.16	110.96	107.77
6	A	1319	PTY	C1-O4-C30	2.12	124.96	117.12
4	A	1314	TCH	C1-C10-C9	2.10	114.65	111.35
4	A	1314	TCH	C6-C5-C10	2.03	114.82	112.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1313	TCH	C6-C5-C4	-2.02	108.86	111.19

There are no chirality outliers.

All (61) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1319	PTY	N1-C2-C3-O11
6	A	1319	PTY	C31-C30-O4-C1
6	A	1319	PTY	O30-C30-O4-C1
5	A	1316	LDA	N1-C1-C2-C3
4	A	1313	TCH	C25-C26-S26-O1S
4	A	1313	TCH	C25-C26-S26-O2S
6	A	1318	PTY	N1-C2-C3-O11
6	A	1318	PTY	C5-O14-P1-O11
6	A	1318	PTY	C5-O14-P1-O12
6	A	1318	PTY	C5-O14-P1-O13
5	A	1317	LDA	C2-C1-N1-CM1
5	A	1317	LDA	C2-C1-N1-CM2
6	A	1318	PTY	C31-C30-O4-C1
6	A	1319	PTY	C11-C8-O7-C6
6	A	1318	PTY	O30-C30-O4-C1
6	A	1319	PTY	O10-C8-O7-C6
6	A	1318	PTY	C11-C8-O7-C6
6	A	1318	PTY	O10-C8-O7-C6
6	A	1319	PTY	C32-C33-C34-C35
6	A	1318	PTY	C35-C36-C37-C38
5	A	1316	LDA	C4-C5-C6-C7
6	A	1319	PTY	C30-C31-C32-C33
5	A	1316	LDA	C2-C3-C4-C5
5	A	1317	LDA	C11-C10-C9-C8
5	A	1320	LDA	C2-C3-C4-C5
6	A	1318	PTY	C14-C15-C16-C17
6	A	1318	PTY	C15-C16-C17-C18
6	A	1318	PTY	C36-C37-C38-C39
6	A	1319	PTY	O14-C5-C6-C1
5	A	1316	LDA	C5-C6-C7-C8
5	A	1315	LDA	C3-C4-C5-C6
6	A	1318	PTY	C41-C42-C43-C44
6	A	1319	PTY	O4-C1-C6-C5
6	A	1319	PTY	C16-C17-C18-C19
5	A	1320	LDA	C3-C4-C5-C6
6	A	1319	PTY	O14-C5-C6-O7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	1318	PTY	C16-C17-C18-C19
6	A	1318	PTY	C12-C13-C14-C15
6	A	1318	PTY	C34-C35-C36-C37
5	A	1315	LDA	C2-C3-C4-C5
5	A	1316	LDA	C3-C4-C5-C6
5	A	1317	LDA	C2-C3-C4-C5
5	A	1321	LDA	C2-C1-N1-CM1
6	A	1318	PTY	C3-O11-P1-O14
6	A	1319	PTY	C31-C32-C33-C34
5	A	1317	LDA	C2-C1-N1-O1
5	A	1316	LDA	C7-C8-C9-C10
6	A	1318	PTY	C6-C5-O14-P1
6	A	1319	PTY	C3-O11-P1-O14
6	A	1319	PTY	C15-C16-C17-C18
5	A	1317	LDA	C7-C8-C9-C10
6	A	1319	PTY	C11-C12-C13-C14
6	A	1319	PTY	C13-C14-C15-C16
5	A	1321	LDA	C2-C3-C4-C5
6	A	1319	PTY	O4-C1-C6-O7
6	A	1318	PTY	C8-C11-C12-C13
5	A	1321	LDA	C2-C1-N1-CM2
5	A	1317	LDA	C5-C6-C7-C8
6	A	1319	PTY	C3-O11-P1-O13
6	A	1318	PTY	C32-C33-C34-C35
6	A	1318	PTY	C5-C6-O7-C8


There are no ring outliers.


4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1319	PTY	2	0
4	A	1313	TCH	4	0
6	A	1318	PTY	2	0
5	A	1317	LDA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/332 (92%)	-0.01	26 (8%) 11 9	32, 50, 102, 144	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ILE	9.2
1	A	2	ASN	7.1
1	A	4	LEU	6.4
1	A	190	LYS	4.2
1	A	183	VAL	4.1
1	A	246	TRP	3.8
1	A	28	PHE	3.5
1	A	119	TYR	3.4
1	A	7	ILE	3.4
1	A	93	ASN	3.1
1	A	122	ARG	3.0
1	A	5	SER	3.0
1	A	188	THR	2.9
1	A	251	TYR	2.9
1	A	42	TRP	2.8
1	A	86	TRP	2.8
1	A	309	ALA	2.7
1	A	184	LEU	2.4
1	A	193	ASP	2.4
1	A	182	LYS	2.3
1	A	187	LYS	2.2
1	A	242	PHE	2.2
1	A	91	LEU	2.2
1	A	176	LEU	2.1
1	A	90	LYS	2.0
1	A	179	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	LDA	A	1316	16/16	0.68	0.35	77,80,90,92	0
6	PTY	A	1319	34/50	0.71	0.28	84,92,105,106	0
6	PTY	A	1318	40/50	0.76	0.21	67,80,106,106	0
4	TCH	A	1313	35/35	0.82	0.25	56,81,144,180	0
5	LDA	A	1321	9/16	0.84	0.34	79,82,85,86	0
5	LDA	A	1320	9/16	0.84	0.16	67,68,69,69	0
4	TCH	A	1314	22/35	0.86	0.17	65,85,110,116	0
5	LDA	A	1317	16/16	0.88	0.19	78,79,86,87	0
5	LDA	A	1315	10/16	0.94	0.23	64,65,66,67	0
3	HG	A	1312	1/1	0.97	0.05	129,129,129,129	0
2	NA	A	1310	1/1	0.98	0.06	42,42,42,42	0
2	NA	A	1311	1/1	0.99	0.10	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PTY A 1319:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PTY A 1318:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TCH A 1313:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TCH A 1314:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.