



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:21 am BST

PDB ID : 4ZU3
Title : Halohydrin hydrogen-halide-lyases, HheB
Authors : Watanabe, F.; Yu, F.; Ohtaki, A.; Yamanaka, Y.; Noguchi, K.; Yohda, M.;
Odaka, M.
Deposited on : 2015-05-15
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>%</div> <div> <div></div> <div>96%</div> <div>..</div> </div> </div>
1	B	227	<div> <div></div> <div>96%</div> <div>..</div> </div>
1	C	227	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>
1	D	227	<div> <div>5%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	E	227	<div> <div>%</div> <div> <div></div> <div>95%</div> <div>..</div> </div> </div>
1	F	227	<div> <div>4%</div> <div> <div></div> <div>96%</div> <div>..</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 21766 atoms, of which 10260 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Halohydrin epoxidase B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	224	Total	C	H	N	O	S	0	0	0
			3463	1116	1701	308	334	4			
1	B	225	Total	C	H	N	O	S	0	0	0
			3477	1120	1707	310	336	4			
1	C	224	Total	C	H	N	O	S	0	0	0
			3463	1116	1701	308	334	4			
1	D	224	Total	C	H	N	O	S	0	0	0
			3463	1116	1701	308	334	4			
1	E	225	Total	C	H	N	O	S	0	0	0
			3477	1120	1707	310	336	4			
1	F	224	Total	C	H	N	O	S	0	0	0
			3463	1116	1701	308	334	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q46347
A	2	LYS	-	expression tag	UNP Q46347
B	1	MET	-	initiating methionine	UNP Q46347
B	2	LYS	-	expression tag	UNP Q46347
C	1	MET	-	initiating methionine	UNP Q46347
C	2	LYS	-	expression tag	UNP Q46347
D	1	MET	-	initiating methionine	UNP Q46347
D	2	LYS	-	expression tag	UNP Q46347
E	1	MET	-	initiating methionine	UNP Q46347
E	2	LYS	-	expression tag	UNP Q46347
F	1	MET	-	initiating methionine	UNP Q46347
F	2	LYS	-	expression tag	UNP Q46347

- Molecule 2 is 3-hydroxypentanedinitrile (three-letter code: 4SD) (formula: C₅H₆N₂O).



- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	130	Total	O	0	0
			130	130		
5	B	167	Total	O	0	0
			167	167		
5	C	137	Total	O	0	0
			137	137		
5	D	106	Total	O	0	0
			106	106		
5	E	198	Total	O	0	0
			198	198		
5	F	118	Total	O	0	0
			118	118		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Halohydrin epoxidase B



- Molecule 1: Halohydrin epoxidase B




- Molecule 1: Halohydrin epoxidase B




- Molecule 1: Halohydrin epoxidase B



- Molecule 1: Halohydrin epoxidase B



- Molecule 1: Halohydrin epoxidase B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	122.91Å 215.64Å 104.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.01 – 2.20 47.01 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.01-2.20) 99.5 (47.01-2.19)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.56 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.10-2152-000)	Depositor
R, R_{free}	0.163 , 0.208 0.166 , 0.211	Depositor DCC
R_{free} test set	3586 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.005 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.014 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21766	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, 4SD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1800	0.49	0/2443
1	B	0.31	0/1808	0.50	0/2454
1	C	0.29	0/1800	0.49	0/2443
1	D	0.33	0/1800	0.50	0/2443
1	E	0.31	0/1808	0.50	0/2454
1	F	0.30	0/1800	0.50	0/2443
All	All	0.31	0/10816	0.50	0/14680

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1762	1701	1701	2	0
1	B	1770	1707	1707	3	0
1	C	1762	1701	1701	5	0
1	D	1762	1701	1701	7	0
1	E	1770	1707	1707	6	0
1	F	1762	1701	1701	3	0
2	A	8	6	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	16	12	12	1	0
2	C	8	6	6	0	0
2	D	8	6	6	0	0
2	E	8	6	6	1	0
2	F	8	6	6	0	0
3	A	5	0	0	1	0
4	E	1	0	0	0	0
5	A	130	0	0	1	0
5	B	167	0	0	0	0
5	C	137	0	0	2	1
5	D	106	0	0	4	1
5	E	198	0	0	2	0
5	F	118	0	0	2	0
All	All	11506	10260	10260	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:123:ARG:NH1	5:C:401:HOH:O	1.88	1.04
1:E:88:GLU:OE1	5:E:401:HOH:O	1.94	0.85
1:D:186:ARG:NH1	5:D:402:HOH:O	2.23	0.70
3:A:302:SO4:O3	5:A:401:HOH:O	2.16	0.62
1:D:80:GLU:OE1	5:D:401:HOH:O	2.15	0.59
1:D:167:LYS:HG3	5:D:437:HOH:O	2.03	0.57
1:E:167:LYS:NZ	5:E:404:HOH:O	2.32	0.51
1:A:30:GLU:OE2	1:A:197:ARG:NH2	2.43	0.51
1:C:58:ILE:O	1:C:108:ARG:NH2	2.44	0.51
1:F:107:GLU:OE1	5:F:401:HOH:O	2.19	0.49
1:B:3:ASN:HD22	1:B:3:ASN:N	2.10	0.49
1:B:123:ARG:HG2	1:D:123:ARG:NH1	2.28	0.49
1:C:77:GLU:OE2	5:C:402:HOH:O	2.19	0.48
1:F:22:GLU:HB3	5:F:407:HOH:O	2.12	0.48
1:E:131:TYR:HH	2:E:301:4SD:H6	1.62	0.47
1:B:142:VAL:HG21	1:B:158:PHE:HB2	1.96	0.47
1:C:184:MET:HE1	1:C:191:GLY:C	2.35	0.47
1:D:109:ASN:ND2	5:D:404:HOH:O	2.31	0.46
1:F:5:ARG:HG2	1:F:5:ARG:HH21	1.82	0.44
1:A:10:ARG:NH1	1:A:57:HIS:O	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:117:GLY:HA3	1:C:158:PHE:CZ	2.54	0.43
1:E:84:GLN:O	1:E:88:GLU:HG2	2.19	0.42
1:D:184:MET:SD	1:D:193:LEU:HD13	2.60	0.42
1:D:71:PHE:HA	1:D:127:GLY:O	2.21	0.41
1:E:11:VAL:HG22	1:E:60:VAL:HB	2.02	0.41
2:B:302:4SD:H1	1:E:173:GLU:HB3	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:434:HOH:O	5:D:408:HOH:O[6_454]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/227 (98%)	214 (96%)	8 (4%)	0	100	100
1	B	223/227 (98%)	214 (96%)	9 (4%)	0	100	100
1	C	222/227 (98%)	213 (96%)	9 (4%)	0	100	100
1	D	222/227 (98%)	214 (96%)	8 (4%)	0	100	100
1	E	223/227 (98%)	216 (97%)	7 (3%)	0	100	100
1	F	222/227 (98%)	212 (96%)	10 (4%)	0	100	100
All	All	1334/1362 (98%)	1283 (96%)	51 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/178 (98%)	173 (99%)	2 (1%)	73	85
1	B	176/178 (99%)	170 (97%)	6 (3%)	37	47
1	C	175/178 (98%)	170 (97%)	5 (3%)	42	54
1	D	175/178 (98%)	171 (98%)	4 (2%)	50	63
1	E	176/178 (99%)	172 (98%)	4 (2%)	50	63
1	F	175/178 (98%)	170 (97%)	5 (3%)	42	54
All	All	1052/1068 (98%)	1026 (98%)	26 (2%)	47	60

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	GLU
1	A	182	GLU
1	B	3	ASN
1	B	101	VAL
1	B	123	ARG
1	B	136	PHE
1	B	182	GLU
1	B	183	ASP
1	C	19	TYR
1	C	109	ASN
1	C	162	HIS
1	C	179	GLU
1	C	184	MET
1	D	19	TYR
1	D	53	GLU
1	D	151	ARG
1	D	173	GLU
1	E	3	ASN
1	E	19	TYR
1	E	151	ARG
1	E	167	LYS
1	F	5	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	17	ASP
1	F	19	TYR
1	F	182	GLU
1	F	184	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	4SD	B	302	-	7,7,7	0.46	0	6,7,7	0.97	0
2	4SD	D	301	-	7,7,7	0.43	0	6,7,7	1.53	2 (33%)
3	SO4	A	302	-	4,4,4	0.11	0	6,6,6	0.21	0
2	4SD	B	301	-	7,7,7	0.45	0	6,7,7	1.14	0
2	4SD	A	301	-	7,7,7	0.57	0	6,7,7	1.45	0
2	4SD	C	301	-	7,7,7	0.37	0	6,7,7	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	4SD	F	301	-	7,7,7	0.47	0	6,7,7	1.42	1 (16%)
2	4SD	E	301	-	7,7,7	0.42	0	6,7,7	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4SD	B	302	-	-	2/4/6/6	-
2	4SD	D	301	-	-	0/4/6/6	-
2	4SD	B	301	-	-	3/4/6/6	-
2	4SD	A	301	-	-	0/4/6/6	-
2	4SD	C	301	-	-	0/4/6/6	-
2	4SD	F	301	-	-	4/4/6/6	-
2	4SD	E	301	-	-	2/4/6/6	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	301	4SD	O1-C1-C4	-2.94	102.20	109.72
2	D	301	4SD	O1-C1-C4	-2.44	103.48	109.72
2	D	301	4SD	C1-C4-C5	-2.26	106.34	112.82

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	302	4SD	C4-C1-C2-C3
2	B	302	4SD	O1-C1-C2-C3
2	B	301	4SD	C4-C1-C2-C3
2	F	301	4SD	C4-C1-C2-C3
2	F	301	4SD	O1-C1-C2-C3
2	F	301	4SD	C2-C1-C4-C5
2	F	301	4SD	O1-C1-C4-C5
2	E	301	4SD	C2-C1-C4-C5
2	B	301	4SD	O1-C1-C4-C5
2	E	301	4SD	O1-C1-C4-C5
2	B	301	4SD	C2-C1-C4-C5

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	4SD	1	0
3	A	302	SO4	1	0
2	E	301	4SD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/227 (98%)	-0.22	3 (1%) 77 75	20, 30, 50, 72	0
1	B	225/227 (99%)	-0.50	1 (0%) 92 91	18, 25, 42, 69	0
1	C	224/227 (98%)	-0.13	8 (3%) 42 41	19, 29, 50, 70	0
1	D	224/227 (98%)	0.20	12 (5%) 25 24	21, 32, 59, 78	0
1	E	225/227 (99%)	-0.35	3 (1%) 77 75	14, 22, 38, 61	0
1	F	224/227 (98%)	-0.16	8 (3%) 42 41	18, 30, 49, 71	0
All	All	1346/1362 (98%)	-0.19	35 (2%) 56 53	14, 28, 50, 78	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	176	ALA	6.3
1	C	176	ALA	4.6
1	F	54	ARG	4.5
1	D	177	THR	4.3
1	C	179	GLU	4.1
1	D	32	GLY	3.5
1	F	177	THR	3.4
1	C	7	ALA	2.9
1	D	219	ILE	2.7
1	D	26	GLN	2.7
1	D	218	SER	2.6
1	E	219	ILE	2.6
1	D	179	GLU	2.6
1	C	182	GLU	2.5
1	F	178	ASP	2.4
1	C	219	ILE	2.4
1	D	178	ASP	2.4
1	C	56	GLY	2.4
1	A	179	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	179	GLU	2.3
1	E	218	SER	2.2
1	C	26	GLN	2.2
1	D	122	MET	2.1
1	D	54	ARG	2.1
1	F	218	SER	2.1
1	C	122	MET	2.1
1	E	122	MET	2.1
1	D	158	PHE	2.1
1	F	180	PHE	2.1
1	F	121	ALA	2.1
1	A	218	SER	2.1
1	F	122	MET	2.0
1	D	182	GLU	2.0
1	B	219	ILE	2.0
1	A	44	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	302	5/5	0.84	0.15	64,67,69,84	0
2	4SD	A	301	8/8	0.91	0.14	22,27,34,36	0
2	4SD	B	302	8/8	0.92	0.18	33,39,44,50	0
2	4SD	D	301	8/8	0.92	0.18	24,32,46,46	0
2	4SD	C	301	8/8	0.92	0.17	28,35,38,38	0
2	4SD	F	301	8/8	0.93	0.15	27,35,45,45	0
2	4SD	E	301	8/8	0.93	0.20	22,27,33,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	4SD	B	301	8/8	0.95	0.13	21,29,37,37	0
4	MG	E	302	1/1	0.98	0.06	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.