



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:10 am BST

PDB ID : 5ZYD  
Title : Crystal Structure of Glucose Isomerase Soaked with Glucose  
Authors : Nam, K.H.  
Deposited on : 2018-05-24  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 10px; left: 0; width: 100%; text-align: center;">88% <span style="float: right;">9% ..</span></div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	14	0
			3157	1979	578	591	9			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	522	Total 522	O 522	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.50 Å 97.82 Å 102.30 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.70 – 1.40 30.75 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.4 (70.70-1.40) 99.4 (30.75-1.40)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.37 (at 1.40 Å)	Xtriage
Refinement program	REFMAC 5.8.0218	Depositor
R, $R_{free}$	0.151 , 0.165 0.159 , 0.172	Depositor DCC
$R_{free}$ test set	4675 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.3	Xtriage
Anisotropy	0.417	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.005 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3684	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.14	9/3230 (0.3%)	1.28	39/4368 (0.9%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	204	GLU	CD-OE1	7.74	1.34	1.25
1	A	186[A]	GLU	CD-OE2	5.62	1.31	1.25
1	A	186[B]	GLU	CD-OE2	5.62	1.31	1.25
1	A	280	TYR	CD2-CE2	5.60	1.47	1.39
1	A	204	GLU	CG-CD	5.42	1.60	1.51
1	A	280	TYR	CE1-CZ	-5.35	1.31	1.38
1	A	331[A]	ARG	CZ-NH1	-5.16	1.26	1.33
1	A	331[B]	ARG	CZ-NH1	-5.16	1.26	1.33
1	A	337	GLU	CD-OE2	-5.11	1.20	1.25

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340[A]	ARG	NE-CZ-NH1	11.09	125.85	120.30
1	A	340[B]	ARG	NE-CZ-NH1	11.09	125.85	120.30
1	A	113	ARG	NE-CZ-NH1	10.72	125.66	120.30
1	A	205[A]	ARG	NE-CZ-NH1	-9.89	115.35	120.30
1	A	205[B]	ARG	NE-CZ-NH1	-9.89	115.35	120.30
1	A	266[A]	ARG	NE-CZ-NH2	-8.89	115.85	120.30
1	A	266[B]	ARG	NE-CZ-NH2	-8.89	115.85	120.30
1	A	340[A]	ARG	NE-CZ-NH2	-8.77	115.91	120.30
1	A	340[B]	ARG	NE-CZ-NH2	-8.77	115.91	120.30
1	A	188	ARG	NE-CZ-NH1	8.41	124.50	120.30
1	A	255[A]	ASP	CB-CG-OD2	-7.80	111.28	118.30
1	A	255[B]	ASP	CB-CG-OD2	-7.80	111.28	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	292	ARG	NE-CZ-NH2	7.79	124.20	120.30
1	A	257	ASP	CB-CG-OD1	7.41	124.97	118.30
1	A	113	ARG	NE-CZ-NH2	-7.31	116.64	120.30
1	A	157	ARG	NE-CZ-NH2	7.25	123.92	120.30
1	A	188	ARG	NE-CZ-NH2	-7.19	116.70	120.30
1	A	284	ARG	NE-CZ-NH2	-6.91	116.84	120.30
1	A	110	ASP	CB-CG-OD1	6.89	124.50	118.30
1	A	292	ARG	NE-CZ-NH1	-6.65	116.97	120.30
1	A	331[A]	ARG	NE-CZ-NH1	6.63	123.61	120.30
1	A	331[B]	ARG	NE-CZ-NH1	6.63	123.61	120.30
1	A	321	ARG	NE-CZ-NH2	6.33	123.46	120.30
1	A	245	ASP	CB-CG-OD2	-5.89	113.00	118.30
1	A	374[A]	ARG	NE-CZ-NH1	-5.72	117.44	120.30
1	A	374[B]	ARG	NE-CZ-NH1	-5.72	117.44	120.30
1	A	117	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	157	ARG	NE-CZ-NH1	-5.52	117.54	120.30
1	A	10	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	A	32	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	361	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	337	GLU	CG-CD-OE1	5.28	128.86	118.30
1	A	331[A]	ARG	NH1-CZ-NH2	-5.25	113.62	119.40
1	A	331[B]	ARG	NH1-CZ-NH2	-5.25	113.62	119.40
1	A	212	TYR	CB-CG-CD1	-5.23	117.86	121.00
1	A	381	ASP	CB-CG-OD1	5.23	123.00	118.30
1	A	55	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	331[A]	ARG	CD-NE-CZ	5.11	130.75	123.60
1	A	331[B]	ARG	CD-NE-CZ	5.11	130.75	123.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3157	0	3019	28	0
2	A	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	3	1	0
4	A	522	0	0	15	7
All	All	3684	0	3022	28	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:PRO:HB2	1:A:26[B]:PHE:CE1	2.04	0.93
1:A:331[A]:ARG:NH1	4:A:502:HOH:O	2.08	0.81
1:A:374[A]:ARG:NH2	4:A:504:HOH:O	2.13	0.81
1:A:262:ALA:O	4:A:501:HOH:O	2.01	0.79
1:A:132:GLU:HG2	4:A:535:HOH:O	1.82	0.78
1:A:3:TYR:N	4:A:505:HOH:O	2.18	0.76
1:A:337:GLU:HG2	1:A:340[A]:ARG:NH2	2.04	0.72
1:A:255[B]:ASP:OD2	4:A:503:HOH:O	2.10	0.70
1:A:132:GLU:CG	4:A:535:HOH:O	2.40	0.67
1:A:3:TYR:N	4:A:508:HOH:O	2.29	0.66
1:A:25:PRO:HB2	1:A:26[B]:PHE:CD1	2.33	0.62
1:A:374[B]:ARG:HG3	1:A:374[B]:ARG:HH21	1.68	0.58
1:A:261:GLY:C	4:A:501:HOH:O	2.46	0.53
1:A:374[A]:ARG:CZ	4:A:504:HOH:O	2.53	0.51
1:A:223:MET:SD	1:A:255[A]:ASP:HB3	2.51	0.51
1:A:354[A]:ARG:NH1	1:A:359:GLU:OE2	2.44	0.50
1:A:24:ASP:HB2	1:A:25:PRO:CD	2.42	0.49
1:A:9:ASP:O	3:A:402:ACT:H2	2.12	0.49
1:A:157:ARG:HH11	1:A:160[B]:GLU:CD	2.17	0.48
1:A:26[B]:PHE:CD1	1:A:26[B]:PHE:N	2.82	0.48
1:A:289:LYS:NZ	4:A:515:HOH:O	2.41	0.48
1:A:374[B]:ARG:NH2	4:A:518:HOH:O	2.43	0.46
1:A:74:ARG:NH1	4:A:530:HOH:O	2.50	0.45
1:A:158[A]:MET:CE	4:A:973:HOH:O	2.65	0.44
1:A:283:PRO:HD2	4:A:555:HOH:O	2.20	0.41
1:A:25:PRO:CB	1:A:26[B]:PHE:CE1	2.90	0.40
1:A:24:ASP:HB2	1:A:25:PRO:HD3	2.01	0.40
1:A:205[B]:ARG:HA	1:A:205[B]:ARG:HD2	1.78	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:718:HOH:O	4:A:718:HOH:O[3_656]	1.89	0.31
4:A:519:HOH:O	4:A:832:HOH:O[2_655]	1.92	0.28
4:A:941:HOH:O	4:A:979:HOH:O[6_554]	2.04	0.16
4:A:979:HOH:O	4:A:1016:HOH:O[6_555]	2.05	0.15
4:A:800:HOH:O	4:A:890:HOH:O[4_556]	2.13	0.07
4:A:855:HOH:O	4:A:932:HOH:O[7_555]	2.16	0.04
4:A:565:HOH:O	4:A:881:HOH:O[7_545]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	396/388 (102%)	381 (96%)	13 (3%)	2 (0%)	29 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186[A]	GLU
1	A	186[B]	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	315/304 (104%)	315 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	A	402	-	1,3,3	2.36	1 (100%)	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	ACT	CH3-C	2.36	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	384/388 (98%)	-0.11	2 (0%) 91 89	6, 10, 21, 32	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	175	ASP	3.4
1	A	66	SER	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	A	402	4/4	0.75	0.25	20,23,28,41	0
2	MG	A	401	1/1	1.00	0.16	14,14,14,14	0

## 6.5 Other polymers

There are no such residues in this entry.