



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2024 – 08:39 AM EDT

PDB ID : 3CCB
Title : Crystal Structure of Human DPP4 in complex with a benzimidazole derivative
Authors : Wallace, M.B.; Skene, R.J.
Deposited on : 2008-02-25
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	740	<div> <div>7%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>..</div> </div> </div>
1	B	740	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>..</div> </div> </div>
1	C	740	<div> <div>9%</div> <div> <div></div> <div>83%</div> <div>14%</div> <div>..</div> </div> </div>
1	D	740	<div> <div>11%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>..</div> </div> </div>
2	E	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%
2	G	2	 100%
2	H	2	 50% 50%
2	I	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	804	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	724	Total	C	N	O	S	0	1	0
			5935	3812	977	1120	26			
1	B	729	Total	C	N	O	S	0	0	0
			5965	3830	983	1126	26			
1	C	724	Total	C	N	O	S	0	1	0
			5936	3813	977	1120	26			
1	D	724	Total	C	N	O	S	0	0	0
			5929	3809	974	1120	26			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	ALA	-	expression tag	UNP P27487
A	28	ASP	-	expression tag	UNP P27487
A	29	PRO	-	expression tag	UNP P27487
A	30	GLY	-	expression tag	UNP P27487
A	31	GLY	-	expression tag	UNP P27487
A	32	SER	-	expression tag	UNP P27487
A	33	HIS	-	expression tag	UNP P27487
A	34	HIS	-	expression tag	UNP P27487
A	35	HIS	-	expression tag	UNP P27487
A	36	HIS	-	expression tag	UNP P27487
A	37	HIS	-	expression tag	UNP P27487
A	38	HIS	-	expression tag	UNP P27487
B	27	ALA	-	expression tag	UNP P27487
B	28	ASP	-	expression tag	UNP P27487
B	29	PRO	-	expression tag	UNP P27487
B	30	GLY	-	expression tag	UNP P27487
B	31	GLY	-	expression tag	UNP P27487
B	32	SER	-	expression tag	UNP P27487
B	33	HIS	-	expression tag	UNP P27487
B	34	HIS	-	expression tag	UNP P27487
B	35	HIS	-	expression tag	UNP P27487

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	36	HIS	-	expression tag	UNP P27487
B	37	HIS	-	expression tag	UNP P27487
B	38	HIS	-	expression tag	UNP P27487
C	27	ALA	-	expression tag	UNP P27487
C	28	ASP	-	expression tag	UNP P27487
C	29	PRO	-	expression tag	UNP P27487
C	30	GLY	-	expression tag	UNP P27487
C	31	GLY	-	expression tag	UNP P27487
C	32	SER	-	expression tag	UNP P27487
C	33	HIS	-	expression tag	UNP P27487
C	34	HIS	-	expression tag	UNP P27487
C	35	HIS	-	expression tag	UNP P27487
C	36	HIS	-	expression tag	UNP P27487
C	37	HIS	-	expression tag	UNP P27487
C	38	HIS	-	expression tag	UNP P27487
D	27	ALA	-	expression tag	UNP P27487
D	28	ASP	-	expression tag	UNP P27487
D	29	PRO	-	expression tag	UNP P27487
D	30	GLY	-	expression tag	UNP P27487
D	31	GLY	-	expression tag	UNP P27487
D	32	SER	-	expression tag	UNP P27487
D	33	HIS	-	expression tag	UNP P27487
D	34	HIS	-	expression tag	UNP P27487
D	35	HIS	-	expression tag	UNP P27487
D	36	HIS	-	expression tag	UNP P27487
D	37	HIS	-	expression tag	UNP P27487
D	38	HIS	-	expression tag	UNP P27487

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 1-biphenyl-2-ylmethanamine (three-letter code: B2Y) (formula: $C_{13}H_{13}N$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			14	13	1		
4	B	1	Total	C	N	0	0
			14	13	1		
4	C	1	Total	C	N	0	0
			14	13	1		
4	D	1	Total	C	N	0	0
			14	13	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	206	Total	O	0	0
			206	206		
5	B	191	Total	O	0	0
			191	191		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	189	Total	O	0	0
			189	189		
5	D	90	Total	O	0	0
			90	90		


- Molecule 1: Dipeptidyl peptidase 4



• Molecule 1: Dipeptidyl peptidase 4



• Molecule 1: Dipeptidyl peptidase 4



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.09Å 123.01Å 144.65Å 90.00° 114.84° 90.00°	Depositor
Resolution (Å)	32.80 – 2.49 32.80 – 2.49	Depositor EDS
% Data completeness (in resolution range)	98.3 (32.80-2.49) 98.5 (32.80-2.49)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.48Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.200 , 0.244 0.234 , 0.270	Depositor DCC
R_{free} test set	6729 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	24805	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, B2Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/6111	0.62	1/8311 (0.0%)
1	B	0.44	0/6138	0.62	1/8348 (0.0%)
1	C	0.44	0/6111	0.61	1/8311 (0.0%)
1	D	0.42	0/6100	0.58	0/8296
All	All	0.44	0/24460	0.61	3/33266 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	415	LEU	CA-CB-CG	5.83	128.71	115.30
1	A	415	LEU	CA-CB-CG	5.37	127.66	115.30
1	B	415	LEU	CA-CB-CG	5.13	127.11	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5935	0	5657	72	0
1	B	5965	0	5672	63	0
1	C	5936	0	5660	65	0
1	D	5929	0	5651	61	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	28	0	25	0	0
2	F	28	0	25	0	0
2	G	28	0	25	1	0
2	H	28	0	25	0	0
2	I	28	0	25	0	0
3	A	56	0	52	0	0
3	B	56	0	52	1	0
3	C	28	0	26	0	0
3	D	28	0	26	1	0
4	A	14	0	13	0	0
4	B	14	0	13	1	0
4	C	14	0	13	0	0
4	D	14	0	13	0	0
5	A	206	0	0	2	0
5	B	191	0	0	0	0
5	C	189	0	0	1	0
5	D	90	0	0	0	0
All	All	24805	0	22973	253	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 253 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:343:ARG:HD2	1:C:389:ILE:HG22	1.42	0.98
1:C:153:GLN:HE22	1:C:170:ASN:H	1.14	0.93
1:A:153:GLN:HE22	1:A:170:ASN:H	1.18	0.90
1:A:253:ARG:HH21	1:B:253:ARG:HH21	1.14	0.90
1:C:54:ARG:HG2	1:C:54:ARG:HH21	1.39	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	721/740 (97%)	684 (95%)	36 (5%)	1 (0%)	48	69
1	B	725/740 (98%)	698 (96%)	25 (3%)	2 (0%)	37	56
1	C	721/740 (97%)	688 (95%)	31 (4%)	2 (0%)	37	56
1	D	720/740 (97%)	684 (95%)	34 (5%)	2 (0%)	37	56
All	All	2887/2960 (98%)	2754 (95%)	126 (4%)	7 (0%)	44	64

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	474	GLY
1	A	474	GLY
1	B	37	HIS
1	B	474	GLY
1	D	474	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	650/662 (98%)	611 (94%)	39 (6%)	16	33
1	B	652/662 (98%)	611 (94%)	41 (6%)	15	30
1	C	650/662 (98%)	614 (94%)	36 (6%)	18	37
1	D	649/662 (98%)	617 (95%)	32 (5%)	21	42
All	All	2601/2648 (98%)	2453 (94%)	148 (6%)	17	35

5 of 148 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	710	ASN
1	D	575	VAL
1	D	90	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	280	THR
1	B	230	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	710	ASN
1	D	572	ASN
1	C	153	GLN
1	D	508	GLN
1	D	748	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	1,2	14,14,15	0.60	0	17,19,21	0.87	1 (5%)
2	NAG	E	2	2	14,14,15	0.46	0	17,19,21	1.04	0
2	NAG	F	1	1,2	14,14,15	0.55	0	17,19,21	1.45	3 (17%)
2	NAG	F	2	2	14,14,15	0.57	0	17,19,21	0.77	0
2	NAG	G	1	1,2	14,14,15	0.63	0	17,19,21	0.93	0
2	NAG	G	2	2	14,14,15	0.53	0	17,19,21	0.93	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	H	1	1,2	14,14,15	0.64	0	17,19,21	0.98	0
2	NAG	H	2	2	14,14,15	0.62	0	17,19,21	1.26	2 (11%)
2	NAG	I	1	1,2	14,14,15	0.45	0	17,19,21	1.02	1 (5%)
2	NAG	I	2	2	14,14,15	0.61	0	17,19,21	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	4/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	4/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	NAG	C4-C3-C2	3.87	116.68	111.02
2	F	1	NAG	C1-O5-C5	2.94	116.13	112.19
2	F	1	NAG	O5-C1-C2	-2.66	107.17	111.29
2	I	1	NAG	C1-O5-C5	2.54	115.59	112.19
2	F	1	NAG	C2-N2-C7	-2.35	119.76	122.90

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	1	NAG	C8-C7-N2-C2
2	F	1	NAG	O7-C7-N2-C2
2	F	2	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	F	2	NAG	O7-C7-N2-C2
2	G	2	NAG	C8-C7-N2-C2


There are no ring outliers.


2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	NAG	1	0
2	G	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	B2Y	B	800	-	14,15,15	0.70	0	15,19,19	0.69	0
3	NAG	C	801	1	14,14,15	0.50	0	17,19,21	1.40	1 (5%)
3	NAG	C	802	1	14,14,15	0.64	0	17,19,21	1.32	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	804	1	14,14,15	0.79	1 (7%)	17,19,21	1.61	3 (17%)
4	B2Y	C	800	-	14,15,15	0.77	0	15,19,19	0.86	0
4	B2Y	A	800	-	14,15,15	0.81	0	15,19,19	0.87	0
3	NAG	B	803	1	14,14,15	0.63	0	17,19,21	1.21	1 (5%)
3	NAG	B	801	1	14,14,15	0.78	1 (7%)	17,19,21	1.39	2 (11%)
3	NAG	D	801	1	14,14,15	0.60	0	17,19,21	1.06	1 (5%)
3	NAG	A	802	1	14,14,15	0.58	0	17,19,21	1.41	1 (5%)
3	NAG	B	806	1	14,14,15	0.56	0	17,19,21	1.64	1 (5%)
3	NAG	B	802	1	14,14,15	0.53	0	17,19,21	1.17	1 (5%)
4	B2Y	D	800	-	14,15,15	0.72	0	15,19,19	0.77	0
3	NAG	A	808	1	14,14,15	0.57	0	17,19,21	1.41	1 (5%)
3	NAG	A	801	1	14,14,15	0.59	0	17,19,21	1.44	1 (5%)
3	NAG	A	803	1	14,14,15	0.59	0	17,19,21	0.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	B2Y	B	800	-	-	0/2/6/6	0/2/2/2
3	NAG	C	801	1	-	0/6/23/26	0/1/1/1
3	NAG	C	802	1	-	2/6/23/26	0/1/1/1
3	NAG	D	804	1	1/1/5/7	5/6/23/26	0/1/1/1
4	B2Y	C	800	-	-	1/2/6/6	0/2/2/2
4	B2Y	A	800	-	-	1/2/6/6	0/2/2/2
3	NAG	B	803	1	-	0/6/23/26	0/1/1/1
3	NAG	B	801	1	-	3/6/23/26	0/1/1/1
3	NAG	D	801	1	-	0/6/23/26	0/1/1/1
3	NAG	A	802	1	-	0/6/23/26	0/1/1/1
3	NAG	B	806	1	-	4/6/23/26	0/1/1/1
3	NAG	B	802	1	-	1/6/23/26	0/1/1/1
4	B2Y	D	800	-	-	0/2/6/6	0/2/2/2
3	NAG	A	808	1	-	0/6/23/26	0/1/1/1
3	NAG	A	801	1	-	2/6/23/26	0/1/1/1
3	NAG	A	803	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	804	NAG	C1-C2	2.53	1.55	1.52
3	B	801	NAG	C1-C2	2.18	1.55	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	806	NAG	C1-O5-C5	5.75	119.89	112.19
3	A	802	NAG	C1-O5-C5	5.49	119.54	112.19
3	A	801	NAG	C1-O5-C5	4.90	118.75	112.19
3	A	808	NAG	C1-O5-C5	4.70	118.49	112.19
3	C	801	NAG	C1-O5-C5	4.48	118.19	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	804	NAG	C1

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	803	NAG	C8-C7-N2-C2
3	A	803	NAG	O7-C7-N2-C2
3	B	801	NAG	C8-C7-N2-C2
3	B	801	NAG	O7-C7-N2-C2
3	B	806	NAG	C8-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	800	B2Y	1	0
3	D	801	NAG	1	0
3	B	802	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	724/740 (97%)	0.65	50 (6%)	24 22	31, 48, 68, 104	1 (0%)
1	B	729/740 (98%)	0.52	30 (4%)	42 39	36, 47, 67, 83	0
1	C	724/740 (97%)	0.72	64 (8%)	17 16	28, 48, 69, 101	1 (0%)
1	D	724/740 (97%)	0.85	83 (11%)	11 10	36, 51, 69, 107	0
All	All	2901/2960 (98%)	0.68	227 (7%)	20 19	28, 48, 68, 107	2 (0%)

The worst 5 of 227 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	135	TYR	6.1
1	C	474	GLY	5.4
1	A	135	TYR	5.2
1	A	81	ALA	5.1
1	C	95	PHE	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	F	1	14/15	0.66	0.15	73,76,79,80	0
2	NAG	F	2	14/15	0.71	0.16	83,84,85,85	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	H	2	14/15	0.77	0.14	73,76,78,78	0
2	NAG	I	2	14/15	0.79	0.15	67,68,72,72	0
2	NAG	G	2	14/15	0.80	0.15	80,82,83,83	0
2	NAG	E	2	14/15	0.84	0.11	68,70,71,71	0
2	NAG	I	1	14/15	0.86	0.13	55,58,61,63	0
2	NAG	G	1	14/15	0.86	0.13	67,69,73,77	0
2	NAG	H	1	14/15	0.89	0.11	63,65,68,72	0
2	NAG	E	1	14/15	0.92	0.10	58,60,62,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	A	801	14/15	0.58	0.18	72,73,74,74	0
3	NAG	B	801	14/15	0.64	0.22	78,79,80,80	0
3	NAG	D	804	14/15	0.67	0.18	78,80,82,82	0
3	NAG	C	801	14/15	0.70	0.15	55,55,56,57	0
3	NAG	B	803	14/15	0.71	0.15	63,65,68,68	0
3	NAG	A	802	14/15	0.73	0.16	65,66,68,68	0
3	NAG	A	803	14/15	0.73	0.14	71,73,76,77	0
3	NAG	D	801	14/15	0.75	0.15	58,60,62,63	0
3	NAG	B	806	14/15	0.75	0.16	67,69,73,74	0
3	NAG	C	802	14/15	0.76	0.14	68,70,74,74	0
3	NAG	A	808	14/15	0.79	0.12	61,63,64,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	802	14/15	0.86	0.10	63,64,66,66	0
4	B2Y	C	800	14/14	0.89	0.16	50,54,55,55	0
4	B2Y	A	800	14/14	0.90	0.17	53,56,58,58	0
4	B2Y	B	800	14/14	0.91	0.19	60,62,63,64	0
4	B2Y	D	800	14/14	0.92	0.12	42,44,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.