



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2024 – 01:25 AM EDT

PDB ID : 2PTS  
Title : Crystal structure of wild type Escherichia coli adenylosuccinate lyase  
Authors : Tsai, M.; Howell, P.L.  
Deposited on : 2007-05-08  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	 2% 76% 19%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylosuccinate lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	Se	0	0	0
			3514	2244	610	650	2	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P0AB89
A	134	MSE	MET	modified residue	UNP P0AB89
A	154	LEU	ILE	conflict	UNP P0AB89
A	184	MSE	MET	modified residue	UNP P0AB89
A	191	MSE	MET	modified residue	UNP P0AB89
A	298	MSE	MET	modified residue	UNP P0AB89
A	393	MSE	MET	modified residue	UNP P0AB89
A	419	MSE	MET	modified residue	UNP P0AB89
A	439	MSE	MET	modified residue	UNP P0AB89
A	451	MSE	MET	modified residue	UNP P0AB89
A	457	HIS	-	expression tag	UNP P0AB89
A	458	HIS	-	expression tag	UNP P0AB89
A	459	HIS	-	expression tag	UNP P0AB89
A	460	HIS	-	expression tag	UNP P0AB89
A	461	HIS	-	expression tag	UNP P0AB89
A	462	HIS	-	expression tag	UNP P0AB89


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	457	Total	O	0	0
			457	457		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenylosuccinate lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.37Å 98.41Å 136.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.00) 99.7 (50.00-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	4.80	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	45.42 (at 2.00Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.165 , 0.202 0.170 , 0.206	Depositor DCC
$R_{free}$ test set	1542 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.3	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3971	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/3583	0.56	1/4855 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	GLU	N-CA-C	-5.26	96.79	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	0	3443	82	0
2	A	457	0	0	8	1
All	All	3971	0	3443	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ASN:HD21	1:A:206:LYS:H	1.16	0.94

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:GLN:HE22	1:A:263:ARG:HE	1.34	0.75
1:A:443:ASN:C	1:A:443:ASN:HD22	1.91	0.74
1:A:322:GLN:HG2	2:A:879:HOH:O	1.89	0.73
1:A:198:LEU:HD13	1:A:260:CYS:SG	2.31	0.71
1:A:303:ASN:HD22	1:A:305:ILE:H	1.38	0.71
1:A:142:VAL:O	1:A:145:PRO:HG2	1.92	0.70
1:A:215:ASN:H	1:A:215:ASN:HD22	1.39	0.70
1:A:321:LEU:HD13	1:A:349:GLY:HA3	1.75	0.69
1:A:167:LEU:HD21	1:A:174:PRO:HB3	1.73	0.69
1:A:213:ASN:HB2	1:A:215:ASN:HD21	1.59	0.67
1:A:346:LEU:O	1:A:350:ILE:HD13	1.95	0.66
1:A:286:GLN:HE22	1:A:302:VAL:HG23	1.61	0.66
1:A:260:CYS:HB2	2:A:788:HOH:O	1.94	0.65
1:A:107:ILE:HD12	1:A:107:ILE:N	2.11	0.64
1:A:440:THR:H	1:A:443:ASN:ND2	1.95	0.64
1:A:309:ASN:HB2	2:A:632:HOH:O	2.00	0.61
1:A:77:GLU:HG3	2:A:582:HOH:O	2.01	0.59
1:A:401:PRO:O	1:A:405:LEU:HD23	2.02	0.59
1:A:108:PRO:HG2	1:A:109:GLU:OE2	2.02	0.59
1:A:104:VAL:HG13	1:A:110:LEU:HB2	1.85	0.58
1:A:303:ASN:ND2	1:A:305:ILE:H	2.01	0.58
1:A:456:LYS:HE3	2:A:781:HOH:O	2.03	0.58
1:A:334:GLN:O	1:A:335:ARG:HB3	2.04	0.57
1:A:42:ARG:HH11	1:A:45:GLN:HE22	1.53	0.57
1:A:214:TYR:OH	1:A:227:HIS:HD2	1.89	0.56
1:A:429:PRO:HG2	1:A:432:GLU:HG2	1.88	0.56
1:A:213:ASN:HB2	1:A:215:ASN:ND2	2.21	0.56
1:A:432:GLU:HA	1:A:432:GLU:OE2	2.06	0.56
1:A:443:ASN:HD22	1:A:444:TYR:N	2.04	0.55
1:A:104:VAL:HG13	1:A:110:LEU:CB	2.36	0.55
1:A:50:HIS:CD2	1:A:52:ALA:HB3	2.43	0.54
1:A:128:ASN:HD21	1:A:206:LYS:N	1.96	0.53
1:A:393:MSE:SE	1:A:405:LEU:HD21	2.59	0.53
1:A:440:THR:H	1:A:443:ASN:HD21	1.57	0.53
1:A:303:ASN:HD22	1:A:303:ASN:C	2.12	0.52
1:A:390:GLN:HG2	1:A:401:PRO:HB2	1.92	0.52
1:A:38:GLN:O	1:A:42:ARG:HG2	2.10	0.51
1:A:215:ASN:H	1:A:215:ASN:ND2	2.05	0.51
1:A:57:PRO:HG2	2:A:598:HOH:O	2.10	0.51
1:A:350:ILE:N	1:A:350:ILE:HD12	2.26	0.51
1:A:128:ASN:ND2	1:A:206:LYS:H	1.98	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:441:PRO:HG2	2:A:495:HOH:O	2.11	0.50
1:A:443:ASN:C	1:A:443:ASN:ND2	2.63	0.50
1:A:151:ILE:O	1:A:155:LYS:HG3	2.12	0.50
1:A:335:ARG:C	1:A:335:ARG:HD3	2.32	0.50
1:A:29:GLU:O	1:A:33:LEU:HD13	2.11	0.49
1:A:197:GLN:HB2	2:A:788:HOH:O	2.11	0.49
1:A:144:LEU:N	1:A:145:PRO:HD2	2.27	0.49
1:A:208:ASN:OD1	1:A:212:GLY:HA2	2.12	0.49
1:A:286:GLN:NE2	1:A:302:VAL:HG23	2.26	0.48
1:A:321:LEU:HD11	1:A:350:ILE:HD12	1.95	0.48
1:A:334:GLN:O	1:A:335:ARG:CB	2.61	0.48
1:A:450:THR:O	1:A:454:GLU:HG3	2.14	0.48
1:A:207:ILE:O	1:A:207:ILE:HG23	2.12	0.48
1:A:35:PHE:O	1:A:39:VAL:HG23	2.14	0.47
1:A:370:ASN:HD22	1:A:370:ASN:C	2.17	0.47
1:A:456:LYS:NZ	1:A:457:HIS:NE2	2.63	0.46
1:A:98:TYR:O	1:A:102:GLU:HG3	2.15	0.46
1:A:208:ASN:CG	1:A:212:GLY:HA2	2.36	0.46
1:A:173:GLN:HB3	1:A:174:PRO:HD2	1.99	0.44
1:A:268:LEU:HD11	1:A:360:THR:HG21	1.98	0.44
1:A:327:LYS:HD3	1:A:327:LYS:C	2.38	0.44
1:A:91:HIS:HE1	1:A:93:VAL:HB	1.83	0.44
1:A:228:GLN:O	1:A:232:GLU:HG3	2.18	0.43
1:A:151:ILE:HG22	1:A:155:LYS:HE2	2.01	0.43
1:A:429:PRO:HG2	1:A:432:GLU:CG	2.48	0.43
1:A:408:LEU:HD22	1:A:422:PHE:CG	2.54	0.42
1:A:286:GLN:NE2	1:A:302:VAL:HA	2.34	0.42
1:A:48:ALA:HA	1:A:56:VAL:O	2.19	0.42
1:A:158:ALA:HA	1:A:181:GLY:HA2	2.01	0.42
1:A:217:HIS:HB3	1:A:226:TRP:CE2	2.55	0.42
1:A:400:LYS:O	1:A:404:LYS:HG3	2.19	0.42
1:A:242:ASN:C	1:A:242:ASN:HD22	2.23	0.41
1:A:408:LEU:HD21	1:A:419:MSE:HE1	2.02	0.41
1:A:392:VAL:HG21	1:A:436:LEU:HD21	2.03	0.41
1:A:318:ASN:O	1:A:322:GLN:HB2	2.20	0.41
1:A:194:GLN:NE2	1:A:263:ARG:HE	2.10	0.40
1:A:231:GLU:O	1:A:235:THR:HG23	2.20	0.40
1:A:176:THR:HA	1:A:177:PRO:HD3	1.89	0.40
1:A:423:ILE:HD13	1:A:436:LEU:HB3	2.04	0.40
1:A:171:HIS:HD2	1:A:173:GLN:OE1	2.05	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-



metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:676:HOH:O	2:A:676:HOH:O[3_555]	1.14	1.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	441/462 (96%)	427 (97%)	12 (3%)	2 (0%)	25 21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	ARG
1	A	207	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/384 (95%)	353 (97%)	11 (3%)	36 37

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LEU
1	A	128	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	206	LYS
1	A	215	ASN
1	A	242	ASN
1	A	303	ASN
1	A	321	LEU
1	A	335	ARG
1	A	370	ASN
1	A	439	MSE
1	A	443	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	45	GLN
1	A	111	HIS
1	A	128	ASN
1	A	131	HIS
1	A	171	HIS
1	A	194	GLN
1	A	197	GLN
1	A	215	ASN
1	A	227	HIS
1	A	228	GLN
1	A	242	ASN
1	A	286	GLN
1	A	303	ASN
1	A	358	GLN
1	A	370	ASN
1	A	421	GLN
1	A	443	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	438/462 (94%)	-0.41	10 (2%) 61 59	4, 11, 29, 45	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	409	THR	3.9
1	A	289	ILE	3.5
1	A	322	GLN	3.2
1	A	402	TYR	2.9
1	A	66	TYR	2.8
1	A	413	ARG	2.6
1	A	107	ILE	2.3
1	A	104	VAL	2.3
1	A	288	THR	2.2
1	A	406	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.