



wwPDB EM Validation Summary Report ⓘ

Oct 27, 2024 – 09:45 AM EDT

PDB ID : 8TAS
EMDB ID : EMD-41141
Title : PRC2 monomer bound to nucleosome
Authors : Sauer, P.V.; Pavlenko, E.; Nogales, E.; Poepsel, S.
Deposited on : 2023-06-27
Resolution : 4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	619	
2	E	753	
3	G	441	
4	I	136	
4	W	136	
5	H	215	
6	J	106	
6	X	106	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	R	133	 54% 27% 19%
7	U	133	 53% 28% 19%
8	S	123	 58% 20% 22%
8	V	123	 51% 26% 23%
9	T	215	 38% 50% 10%
10	O	425	 43% 60% 32% 7%
11	Y	303	 35% 31% 11% 58%

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 29440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polycomb protein SUZ12.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	D	439	3633	2309	652	645	27	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	67	MET	-	initiating methionine	UNP Q15022

- Molecule 2 is a protein called Histone-lysine N-methyltransferase EZH2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	E	570	4580	2871	821	846	42	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	SER	-	expression tag	UNP Q15910
E	0	ASN	-	expression tag	UNP Q15910
E	1	ALA	-	expression tag	UNP Q15910
E	14	ALA	CYS	conflict	UNP Q15910
E	15	CYS	TRP	conflict	UNP Q15910

- Molecule 3 is a protein called Polycomb protein EED.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	G	362	2930	1855	513	541	21	0	0

- Molecule 4 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	114	Total	C	N	O	S	0	0
			917	576	179	159	3		
4	W	99	Total	C	N	O	S	0	0
			817	515	158	141	3		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	102	ALA	GLY	conflict	UNP P84233
W	102	ALA	GLY	conflict	UNP P84233

- Molecule 5 is a DNA chain called DNA (226-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	H	193	Total	C	N	O	P	0	0
			3972	1873	761	1145	193		

- Molecule 6 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	83	Total	C	N	O	S	0	0
			661	418	129	113	1		
6	X	87	Total	C	N	O	S	0	0
			666	419	129	117	1		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	103	SER	-	expression tag	UNP P62799
J	104	SER	-	expression tag	UNP P62799
J	105	GLY	-	expression tag	UNP P62799
X	103	SER	-	expression tag	UNP P62799
X	104	SER	-	expression tag	UNP P62799
X	105	GLY	-	expression tag	UNP P62799

- Molecule 7 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	R	108	Total	C	N	O	S	0	0
			829	522	162	144	1		
7	U	108	Total	C	N	O	S	0	0
			825	519	161	144	1		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-3	SER	-	expression tag	UNP Q6AZJ8
R	-2	ASN	-	expression tag	UNP Q6AZJ8
R	-1	ALA	-	expression tag	UNP Q6AZJ8
R	119	CYS	LYS	conflict	UNP Q6AZJ8
U	-3	SER	-	expression tag	UNP Q6AZJ8
U	-2	ASN	-	expression tag	UNP Q6AZJ8
U	-1	ALA	-	expression tag	UNP Q6AZJ8
U	119	CYS	LYS	conflict	UNP Q6AZJ8

- Molecule 8 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	S	96	Total	C	N	O	S	0	0
			742	466	133	141	2		
8	V	95	Total	C	N	O	S	0	0
			737	463	132	140	2		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	0	MET	-	initiating methionine	UNP P02281
S	29	THR	SER	conflict	UNP P02281
V	0	MET	-	initiating methionine	UNP P02281
V	29	THR	SER	conflict	UNP P02281

- Molecule 9 is a DNA chain called DNA (226-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
9	T	193	Total	C	N	O	P	0	0
			3919	1853	724	1149	193		

- Molecule 10 is a protein called Histone-binding protein RBBP4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	O	395	Total	C	N	O	S	0	0
			3139	1981	535	613	10		


- Molecule 11 is a protein called Zinc finger protein AEBP2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	Y	126	1047	664	206	174	3	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	7	SER	-	expression tag	UNP Q6ZN18
Y	8	ASN	-	expression tag	UNP Q6ZN18
Y	9	ALA	-	expression tag	UNP Q6ZN18
Y	10	TYR	-	expression tag	UNP Q6ZN18
Y	11	THR	-	expression tag	UNP Q6ZN18
Y	12	ARG	-	expression tag	UNP Q6ZN18
Y	13	ARG	-	expression tag	UNP Q6ZN18
Y	14	TYR	-	expression tag	UNP Q6ZN18
Y	15	SER	-	expression tag	UNP Q6ZN18


- Molecule 12 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	S	
12	E	1	26	14	6	5	1	0




• Molecule 6: Histone H4



• Molecule 7: Histone H2A




• Molecule 7: Histone H2A



• Molecule 8: Histone H2B 1.1




• Molecule 8: Histone H2B 1.1






• Molecule 9: DNA (226-MER)



• Molecule 10: Histone-binding protein RBBP4



• Molecule 11: Zinc finger protein AEBP2



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	69000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.135	Depositor
Minimum map value	-0.069	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size (\AA)	505.88998, 505.88998, 505.88998	wwPDB
Map dimensions	330, 330, 330	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.533, 1.533, 1.533	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.25	0/3713	0.49	0/4992
2	E	0.30	0/4682	0.50	0/6308
3	G	0.26	0/3005	0.52	0/4070
4	I	0.38	0/930	0.60	0/1246
4	W	0.44	0/829	0.62	0/1111
5	H	0.65	0/4463	1.55	276/6888 (4.0%)
6	J	0.50	0/668	0.64	0/894
6	X	0.51	0/673	0.60	0/904
7	R	0.47	0/839	0.61	0/1132
7	U	0.44	0/835	0.62	0/1128
8	S	0.47	0/753	0.55	0/1014
8	V	0.49	0/748	0.57	0/1007
9	T	0.66	0/4391	1.59	276/6763 (4.1%)
10	O	0.24	0/3225	0.47	0/4394
11	Y	0.23	0/1070	0.52	0/1437
All	All	0.45	0/30824	0.98	552/43288 (1.3%)

There are no bond length outliers.

The worst 5 of 552 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	T	67	DC	OP1-P-O3'	10.30	127.86	105.20
9	T	46	DG	OP1-P-O3'	9.92	127.03	105.20
9	T	150	DC	OP1-P-O3'	9.89	126.96	105.20
5	H	104	DC	OP1-P-O3'	9.77	126.69	105.20
9	T	120	DC	OP1-P-O3'	9.68	126.50	105.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3633	0	3628	94	0
2	E	4580	0	4483	168	0
3	G	2930	0	2839	147	0
4	I	917	0	971	44	0
4	W	817	0	858	36	0
5	H	3972	0	2154	133	0
6	J	661	0	709	29	0
6	X	666	0	679	20	0
7	R	829	0	887	39	0
7	U	825	0	876	39	0
8	S	742	0	753	22	0
8	V	737	0	751	30	0
9	T	3919	0	2153	133	0
10	O	3139	0	2986	97	0
11	Y	1047	0	1102	22	0
12	E	26	0	19	5	0
All	All	29440	0	25848	873	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 873 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:191:ASN:HB3	3:G:211:LYS:HB3	1.50	0.92
3:G:175:ILE:HG13	3:G:182:CYS:HB2	1.54	0.90
7:U:84:GLN:HE21	7:U:88:ARG:HE	1.19	0.87
11:Y:243:TRP:HB2	11:Y:248:ILE:HG21	1.60	0.84
10:O:293:SER:HG	10:O:297:THR:HG1	1.28	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	423/619 (68%)	403 (95%)	20 (5%)	0	100	100
2	E	560/753 (74%)	511 (91%)	47 (8%)	2 (0%)	30	67
3	G	360/441 (82%)	329 (91%)	31 (9%)	0	100	100
4	I	112/136 (82%)	107 (96%)	5 (4%)	0	100	100
4	W	97/136 (71%)	92 (95%)	5 (5%)	0	100	100
6	J	81/106 (76%)	76 (94%)	5 (6%)	0	100	100
6	X	85/106 (80%)	78 (92%)	7 (8%)	0	100	100
7	R	106/133 (80%)	101 (95%)	5 (5%)	0	100	100
7	U	106/133 (80%)	103 (97%)	3 (3%)	0	100	100
8	S	94/123 (76%)	84 (89%)	10 (11%)	0	100	100
8	V	93/123 (76%)	85 (91%)	8 (9%)	0	100	100
10	O	391/425 (92%)	376 (96%)	15 (4%)	0	100	100
11	Y	124/303 (41%)	114 (92%)	9 (7%)	1 (1%)	16	53
All	All	2632/3537 (74%)	2459 (93%)	170 (6%)	3 (0%)	50	82

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	327	PRO
11	Y	187	HIS
2	E	492	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	411/575 (72%)	408 (99%)	3 (1%)	81	86
2	E	507/672 (75%)	507 (100%)	0	100	100
3	G	324/392 (83%)	324 (100%)	0	100	100
4	I	95/111 (86%)	95 (100%)	0	100	100
4	W	86/111 (78%)	86 (100%)	0	100	100
6	J	68/81 (84%)	68 (100%)	0	100	100
6	X	64/81 (79%)	64 (100%)	0	100	100
7	R	85/104 (82%)	85 (100%)	0	100	100
7	U	84/104 (81%)	84 (100%)	0	100	100
8	S	79/103 (77%)	79 (100%)	0	100	100
8	V	79/103 (77%)	79 (100%)	0	100	100
10	O	351/375 (94%)	350 (100%)	1 (0%)	91	92
11	Y	117/271 (43%)	116 (99%)	1 (1%)	75	83
All	All	2350/3083 (76%)	2345 (100%)	5 (0%)	91	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	124	ARG
1	D	129	ARG
1	D	672	MET
10	O	264	LYS
11	Y	180	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
4	W	125	GLN
10	O	48	GLN
10	O	403	GLN
3	G	146	ASN
2	E	560	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	SAH	E	801	-	23,28,28	1.27	3 (13%)	22,40,40	1.86	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	SAH	E	801	-	-	4/11/31/31	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	E	801	SAH	C2-N3	4.11	1.38	1.32
12	E	801	SAH	C2-N1	2.51	1.38	1.33
12	E	801	SAH	OXT-C	-2.32	1.23	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	E	801	SAH	N3-C2-N1	-6.18	120.28	128.67
12	E	801	SAH	C5'-SD-CG	-4.16	89.91	102.26
12	E	801	SAH	OXT-C-O	-2.61	118.17	124.08

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
12	E	801	SAH	C3'-C4'-C5'-SD
12	E	801	SAH	O4'-C4'-C5'-SD
12	E	801	SAH	C-CA-CB-CG
12	E	801	SAH	CA-CB-CG-SD

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	E	801	SAH	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-41141. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 165



Y Index: 165




Z Index: 165

6.2.2 Raw map



X Index: 165



Y Index: 165




Z Index: 165


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 175




Y Index: 207




Z Index: 148


6.3.2 Raw map



X Index: 174



Y Index: 207




Z Index: 148


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_41141_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 570 nm³; this corresponds to an approximate mass of 515 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.244 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.244 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.86	9.82	7.12

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.86 differs from the reported value 4.1 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-41141 and PDB model 8TAS. Per-residue inclusion information can be found in section 3 on page 8.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8410	 0.2590
D	 0.4870	 0.1280
E	 0.9630	 0.2310
G	 0.9910	 0.2000
H	 0.9900	 0.2930
I	 0.9920	 0.4530
J	 0.9780	 0.4400
O	 0.4800	 0.0920
R	 0.9910	 0.4540
S	 0.9960	 0.4590
T	 0.9940	 0.2910
U	 0.9880	 0.4640
V	 0.9890	 0.4610
W	 0.9920	 0.4520
X	 0.9920	 0.4630
Y	 0.1680	 0.1000

