



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 13, 2024 – 11:03 AM EDT

PDB ID : 2ZM5
Title : Crystal structure of tRNA modification enzyme MiaA in the complex with tRNA(Phe)
Authors : Sakai, J.; Yao, M.; Chimnaronk, S.; Tanaka, I.
Deposited on : 2008-04-11
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1004 (2.54-2.54)
Clashscore	180529	1055 (2.54-2.54)
Ramachandran outliers	177936	1048 (2.54-2.54)
Sidechain outliers	177891	1048 (2.54-2.54)
RSRZ outliers	164620	1004 (2.54-2.54)
RNA backbone	3690	1040 (2.80-2.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>2%</div> <div>62% 26% 8% .</div> </div>
1	B	316	<div> <div>3%</div> <div>43% 39% 13% . .</div> </div>
2	C	76	<div> <div>%</div> <div>39% 29% 16% 13% .</div> </div>
2	D	76	<div> <div>16%</div> <div>29% 38% 12% 12% 9%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA delta(2)-isopentenylpyrophosphate transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	Se	0	0	0
			2403	1523	428	443	2	7			
1	B	305	Total	C	N	O	S	Se	0	0	0
			2397	1518	427	443	2	7			

- Molecule 2 is a RNA chain called tRNA(Phe).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	74	Total	C	N	O	P	0	0	0
			1577	704	282	518	73			
2	D	69	Total	C	N	O	P	0	0	0
			1469	656	261	484	68			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	6	Total	Mg	0	0
			6	6		
3	D	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	53	Total	O	0	0
			53	53		
4	B	31	Total	O	0	0
			31	31		
4	C	57	Total	O	0	0
			57	57		
4	D	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA delta(2)-isopentenylpyrophosphate transferase




- Molecule 1: tRNA delta(2)-isopentenylpyrophosphate transferase





- Molecule 2: tRNA(Phe)





● Molecule 2: tRNA(Phc)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.20Å 89.40Å 150.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.55 20.00 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-2.55) 98.7 (20.00-2.55)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.54Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.240 , 0.270 0.240 , 0.271	Depositor DCC
R_{free} test set	2752 reflections (7.04%)	wwPDB-VP
Wilson B-factor (Å ²)	52.8	Xtriage
Anisotropy	0.702	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8015	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2447	0.68	0/3308
1	B	0.38	0/2441	0.70	1/3300 (0.0%)
2	C	0.68	1/1762 (0.1%)	1.40	21/2746 (0.8%)
2	D	0.54	0/1641	1.05	14/2557 (0.5%)
All	All	0.50	1/8291 (0.0%)	0.98	36/11911 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	1	2
2	D	0	4
All	All	1	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	9	A	C4'-C3'	-5.18	1.47	1.52

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	36	A	O5'-P-OP1	-23.35	82.68	110.70
2	C	35	A	OP2-P-O3'	-18.25	65.04	105.20
2	C	10	G	O5'-P-OP1	-16.58	90.78	105.70
2	C	36	A	O5'-P-OP2	-16.19	91.13	105.70
2	C	35	A	OP1-P-O3'	-16.08	69.81	105.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	64	A	C3'

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	20	U	Sidechain
2	C	37	A	Sidechain
2	D	17	C	Sidechain
2	D	22	G	Sidechain
2	D	34	G	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	0	2408	118	0
1	B	2397	0	2399	236	0
2	C	1577	0	800	41	0
2	D	1469	0	745	49	0
3	C	6	0	0	0	0
3	D	1	0	0	0	0
4	A	53	0	0	1	2
4	B	31	0	0	0	0
4	C	57	0	0	1	0
4	D	21	0	0	1	2
All	All	8015	0	6352	432	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:GLN:HE22	1:B:145:LEU:HD11	1.09	1.15
1:B:147:ARG:HH11	1:B:147:ARG:HB3	1.02	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:41:C:H5'	2:D:41:C:H6	1.13	1.08
1:B:37:GLU:HB3	1:B:95:ILE:HD13	1.38	1.05
1:B:92:MSE:HE1	1:B:115:LEU:HD11	1.39	1.03

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:329:HOH:O	4:D:86:HOH:O[1_565]	2.14	0.06
4:A:329:HOH:O	4:D:97:HOH:O[1_565]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/316 (96%)	294 (97%)	7 (2%)	3 (1%)	13	18
1	B	303/316 (96%)	284 (94%)	12 (4%)	7 (2%)	5	5
All	All	607/632 (96%)	578 (95%)	19 (3%)	10 (2%)	8	10

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	LYS
1	B	237	GLY
1	B	132	ARG
1	B	133	ILE
1	B	142	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/248 (100%)	210 (84%)	39 (16%)	2	1
1	B	249/248 (100%)	197 (79%)	52 (21%)	1	0
All	All	498/496 (100%)	407 (82%)	91 (18%)	1	1

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	129	VAL
1	B	168	LEU
1	B	132	ARG
1	B	140	GLN
1	B	186	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	220	GLN
1	B	309	GLN
1	B	302	GLN
1	B	67	HIS
1	B	216	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	73/76 (96%)	26 (35%)	16 (21%)
2	D	68/76 (89%)	22 (32%)	10 (14%)
All	All	141/152 (92%)	48 (34%)	26 (18%)

5 of 48 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	C
2	C	6	G
2	C	10	G
2	C	11	C
2	C	14	A

5 of 26 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	C	60	U
2	D	19	G
2	D	41	C
2	D	9	A
2	D	32	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/316 (94%)	-0.05	6 (2%) 64 69	29, 49, 70, 95	0
1	B	298/316 (94%)	0.20	10 (3%) 48 53	37, 57, 95, 98	0
2	C	74/76 (97%)	-0.21	1 (1%) 73 76	34, 49, 69, 91	0
2	D	69/76 (90%)	1.24	12 (17%) 5 6	46, 88, 98, 98	0
All	All	740/784 (94%)	0.15	29 (3%) 44 49	29, 53, 97, 98	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	44	G	4.3
1	B	311	VAL	3.8
1	B	168	LEU	3.5
1	B	150	GLN	3.5
2	D	18	G	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	C	81	1/1	0.82	0.24	35,35,35,35	0
3	MG	D	77	1/1	0.88	0.16	41,41,41,41	0
3	MG	C	78	1/1	0.89	0.27	33,33,33,33	0
3	MG	C	80	1/1	0.91	0.34	45,45,45,45	0
3	MG	C	82	1/1	0.93	0.19	49,49,49,49	0
3	MG	C	79	1/1	0.93	0.18	31,31,31,31	0
3	MG	C	77	1/1	0.94	0.15	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.