



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 09:29 PM EDT

PDB ID : 2O7E / pdb_00002o7e
Title : Tyrosine ammonia-lyase from *Rhodobacter sphaeroides* (His89Phe variant), bound to 2-aminoindan-2-phosphonic acid
Authors : Louie, G.V.; Bowman, M.E.; Moffitt, M.C.; Baiga, T.J.; Moore, B.S.; Noel, J.P.
Deposited on : 2006-12-11
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	<div> <div>10%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	B	521	<div> <div>7%</div> <div>84%</div> <div>13%</div> <div>..</div> </div>
1	C	521	<div> <div>7%</div> <div>82%</div> <div>16%</div> <div>..</div> </div>
1	D	521	<div> <div>8%</div> <div>84%</div> <div>13%</div> <div>..</div> </div>
1	E	521	<div> <div>7%</div> <div>84%</div> <div>13%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	521	 4%85%12% ..
1	G	521	 4%85%12% ..
1	H	521	 12%84%12% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 33671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative histidine ammonia-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	S	0	0	0
			3814	2366	729	705	14			
1	B	515	Total	C	N	O	S	0	0	0
			3814	2366	729	705	14			
1	C	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			
1	D	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			
1	E	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			
1	F	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			
1	G	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			
1	H	514	Total	C	N	O	S	0	0	0
			3807	2361	728	704	14			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	89	PHE	HIS	engineered mutation	UNP Q3IWB0
A	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
A	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
A	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
B	89	PHE	HIS	engineered mutation	UNP Q3IWB0
B	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
B	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
B	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
C	89	PHE	HIS	engineered mutation	UNP Q3IWB0
C	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
C	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
C	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
D	89	PHE	HIS	engineered mutation	UNP Q3IWB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
D	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
D	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
E	89	PHE	HIS	engineered mutation	UNP Q3IWB0
E	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
E	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
E	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
F	89	PHE	HIS	engineered mutation	UNP Q3IWB0
F	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
F	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
F	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
G	89	PHE	HIS	engineered mutation	UNP Q3IWB0
G	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
G	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
G	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
H	89	PHE	HIS	engineered mutation	UNP Q3IWB0
H	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
H	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
H	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0

- Molecule 2 is (2-AMINO-2,3-DIHYDRO-1H-INDEN-2-YL)PHOSPHONIC ACID (CCD ID: PMI) (formula: C₉H₁₂NO₃P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			14	9	1	3	1		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	C	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	D	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	E	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	F	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	G	1	Total	C	N	O	P	0	0
			14	9	1	3	1		
2	H	1	Total	C	N	O	P	0	0
			14	9	1	3	1		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	325	Total	O	0	0
			325	325		
3	B	373	Total	O	0	0
			373	373		
3	C	414	Total	O	0	0
			414	414		
3	D	355	Total	O	0	0
			355	355		
3	E	380	Total	O	0	0
			380	380		
3	F	479	Total	O	0	0
			479	479		
3	G	413	Total	O	0	0
			413	413		
3	H	350	Total	O	0	0
			350	350		

- Molecule 1: Putative histidine ammonia-lyase








- Molecule 1: Putative histidine ammonia-lyase




- Molecule 1: Putative histidine ammonia-lyase





- Molecule 1: Putative histidine ammonia-lyase







● Molecule 1: Putative histidine ammonia-lyase



● Molecule 1: Putative histidine ammonia-lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.57Å 154.84Å 164.53Å 90.00° 94.16° 90.00°	Depositor
Resolution (Å)	500.00 – 1.75 164.09 – 1.75	Depositor EDS
% Data completeness (in resolution range)	96.4 (500.00-1.75) 96.3 (164.09-1.75)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.75Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.189 , 0.210 0.187 , 0.207	Depositor DCC
R_{free} test set	21241 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	33671	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PMI, MDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	1/3863 (0.0%)	0.90	12/5254 (0.2%)
1	B	0.41	1/3863 (0.0%)	0.89	14/5254 (0.3%)
1	C	0.43	0/3855	0.90	18/5243 (0.3%)
1	D	0.41	1/3855 (0.0%)	0.90	17/5243 (0.3%)
1	E	0.42	1/3855 (0.0%)	0.89	11/5243 (0.2%)
1	F	0.45	1/3855 (0.0%)	0.91	13/5243 (0.2%)
1	G	0.43	0/3855	0.90	14/5243 (0.3%)
1	H	0.41	1/3855 (0.0%)	0.87	10/5243 (0.2%)
All	All	0.42	6/30856 (0.0%)	0.89	109/41966 (0.3%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	405	MET	SD-CE	-8.24	1.58	1.79
1	H	405	MET	SD-CE	-6.81	1.62	1.79
1	B	405	MET	SD-CE	-6.15	1.64	1.79
1	D	405	MET	SD-CE	-5.51	1.65	1.79
1	A	405	MET	SD-CE	-5.22	1.66	1.79

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	202	VAL	N-CA-C	8.64	121.60	112.96
1	F	278	VAL	N-CA-C	-8.24	98.02	108.89
1	C	202	VAL	N-CA-C	7.96	120.92	112.96
1	B	202	VAL	N-CA-C	7.85	120.81	112.96
1	F	252	ARG	CA-C-N	7.85	129.65	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3814	0	3857	62	0
1	B	3814	0	3858	59	0
1	C	3807	0	3850	66	0
1	D	3807	0	3849	56	0
1	E	3807	0	3850	55	0
1	F	3807	0	3850	61	0
1	G	3807	0	3849	54	0
1	H	3807	0	3850	72	0
2	A	14	0	9	1	0
2	B	14	0	9	3	0
2	C	14	0	9	3	0
2	D	14	0	9	1	0
2	E	14	0	9	3	0
2	F	14	0	9	2	0
2	G	14	0	9	2	0
2	H	14	0	9	3	0
3	A	325	0	0	9	0
3	B	373	0	0	7	0
3	C	414	0	0	12	0
3	D	355	0	0	4	0
3	E	380	0	0	5	0
3	F	479	0	0	11	0
3	G	413	0	0	8	0
3	H	350	0	0	9	0
All	All	33671	0	30885	430	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 430 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:83:GLN:HE22	1:H:196:ARG:HA	1.17	1.08
1:A:83:GLN:HE22	1:A:196:ARG:HA	1.21	1.02
1:B:83:GLN:HE22	1:B:196:ARG:HA	1.25	0.97

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:200:ALA:HB2	3:F:3012:HOH:O	1.64	0.97
1:G:83:GLN:HE22	1:G:196:ARG:HA	1.28	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/521 (98%)	498 (98%)	10 (2%)	2 (0%)	30	16
1	B	510/521 (98%)	502 (98%)	7 (1%)	1 (0%)	44	28
1	C	509/521 (98%)	502 (99%)	6 (1%)	1 (0%)	44	28
1	D	509/521 (98%)	501 (98%)	7 (1%)	1 (0%)	44	28
1	E	509/521 (98%)	500 (98%)	8 (2%)	1 (0%)	44	28
1	F	509/521 (98%)	501 (98%)	7 (1%)	1 (0%)	44	28
1	G	509/521 (98%)	503 (99%)	5 (1%)	1 (0%)	44	28
1	H	509/521 (98%)	497 (98%)	11 (2%)	1 (0%)	44	28
All	All	4074/4168 (98%)	4004 (98%)	61 (2%)	9 (0%)	44	28

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	253	PRO
1	E	253	PRO
1	A	8	LYS
1	A	253	PRO
1	H	253	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/389 (99%)	383 (100%)	1 (0%)	91	88
1	B	384/389 (99%)	383 (100%)	1 (0%)	91	88
1	C	383/389 (98%)	380 (99%)	3 (1%)	79	71
1	D	383/389 (98%)	381 (100%)	2 (0%)	86	82
1	E	383/389 (98%)	381 (100%)	2 (0%)	86	82
1	F	383/389 (98%)	382 (100%)	1 (0%)	91	88
1	G	383/389 (98%)	382 (100%)	1 (0%)	91	88
1	H	383/389 (98%)	380 (99%)	3 (1%)	79	71
All	All	3066/3112 (98%)	3052 (100%)	14 (0%)	86	82

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	202	VAL
1	E	443	THR
1	H	443	THR
1	H	54	ARG
1	H	202	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	164	GLN
1	H	508	GLN
1	G	203	ASN
1	H	78	ASN
1	D	83	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MDO	G	149	1,2	11,13,14	2.44	4 (36%)	15,18,20	2.51	7 (46%)
1	MDO	D	149	1,2	11,13,14	2.25	4 (36%)	15,18,20	2.39	6 (40%)
1	MDO	C	149	1,2	11,13,14	2.48	4 (36%)	15,18,20	2.39	7 (46%)
1	MDO	F	149	1,2	11,13,14	2.42	4 (36%)	15,18,20	2.38	6 (40%)
1	MDO	A	149	1,2	11,13,14	2.33	4 (36%)	15,18,20	2.58	7 (46%)
1	MDO	B	149	1,2	11,13,14	2.44	4 (36%)	15,18,20	2.48	7 (46%)
1	MDO	H	149	1,2	11,13,14	2.39	4 (36%)	15,18,20	2.43	5 (33%)
1	MDO	E	149	1,2	11,13,14	2.42	4 (36%)	15,18,20	2.47	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MDO	G	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	D	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	C	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	F	149	1,2	-	2/4/23/24	0/1/1/1
1	MDO	A	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	B	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	H	149	1,2	-	1/4/23/24	0/1/1/1
1	MDO	E	149	1,2	-	1/4/23/24	0/1/1/1

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	149	MDO	O2-C2	5.57	1.34	1.23
1	C	149	MDO	O2-C2	5.52	1.34	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	149	MDO	O2-C2	5.39	1.34	1.23
1	F	149	MDO	O2-C2	5.33	1.33	1.23
1	H	149	MDO	O2-C2	5.28	1.33	1.23

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	149	MDO	CA2-C2-N3	6.63	109.07	103.50
1	A	149	MDO	CA2-C2-N3	6.60	109.04	103.50
1	G	149	MDO	CA2-C2-N3	6.41	108.88	103.50
1	E	149	MDO	CA2-C2-N3	6.32	108.81	103.50
1	C	149	MDO	CA2-C2-N3	6.32	108.81	103.50

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	149	MDO	N2-C1-CA1-CB
1	B	149	MDO	N2-C1-CA1-CB
1	C	149	MDO	N2-C1-CA1-CB
1	E	149	MDO	N2-C1-CA1-CB
1	F	149	MDO	N2-C1-CA1-CB

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	149	MDO	1	0
1	F	149	MDO	1	0
1	B	149	MDO	1	0
1	H	149	MDO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PMI	F	701	1	12,15,15	2.16	4 (33%)	18,24,24	1.37	2 (11%)
2	PMI	H	701	1	12,15,15	2.10	4 (33%)	18,24,24	1.29	2 (11%)
2	PMI	C	701	1	12,15,15	2.10	3 (25%)	18,24,24	1.37	2 (11%)
2	PMI	E	701	1	12,15,15	2.06	3 (25%)	18,24,24	1.40	2 (11%)
2	PMI	A	701	1	12,15,15	2.14	4 (33%)	18,24,24	1.39	2 (11%)
2	PMI	B	701	1	12,15,15	2.13	4 (33%)	18,24,24	1.38	2 (11%)
2	PMI	D	701	1	12,15,15	2.13	4 (33%)	18,24,24	1.31	2 (11%)
2	PMI	G	701	1	12,15,15	2.07	4 (33%)	18,24,24	1.36	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PMI	F	701	1	-	0/6/19/19	0/2/2/2
2	PMI	H	701	1	-	0/6/19/19	0/2/2/2
2	PMI	C	701	1	-	0/6/19/19	0/2/2/2
2	PMI	E	701	1	-	0/6/19/19	0/2/2/2
2	PMI	A	701	1	-	0/6/19/19	0/2/2/2
2	PMI	B	701	1	-	0/6/19/19	0/2/2/2
2	PMI	D	701	1	-	0/6/19/19	0/2/2/2
2	PMI	G	701	1	-	0/6/19/19	0/2/2/2

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	PMI	C1'-C2'	5.67	1.49	1.39
2	B	701	PMI	C1'-C2'	5.63	1.49	1.39
2	F	701	PMI	C1'-C2'	5.58	1.49	1.39
2	C	701	PMI	C1'-C2'	5.48	1.49	1.39
2	E	701	PMI	C1'-C2'	5.47	1.49	1.39

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	PMI	C9'-C8'-C7'	4.71	110.52	104.06
2	B	701	PMI	C9'-C8'-C7'	4.66	110.47	104.06
2	E	701	PMI	C9'-C8'-C7'	4.66	110.45	104.06
2	F	701	PMI	C9'-C8'-C7'	4.58	110.35	104.06
2	G	701	PMI	C9'-C8'-C7'	4.56	110.33	104.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	701	PMI	2	0
2	H	701	PMI	3	0
2	C	701	PMI	3	0
2	E	701	PMI	3	0
2	A	701	PMI	1	0
2	B	701	PMI	3	0
2	D	701	PMI	1	0
2	G	701	PMI	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	514/521 (98%)	0.57	53 (10%) 13 15	12, 21, 41, 56	0
1	B	514/521 (98%)	0.25	39 (7%) 21 24	9, 18, 37, 52	0
1	C	513/521 (98%)	0.11	36 (7%) 24 27	9, 17, 33, 46	0
1	D	513/521 (98%)	0.37	41 (7%) 20 22	8, 19, 39, 52	0
1	E	513/521 (98%)	0.21	36 (7%) 24 27	9, 17, 36, 62	0
1	F	513/521 (98%)	-0.07	19 (3%) 45 52	7, 14, 29, 41	0
1	G	513/521 (98%)	0.02	23 (4%) 39 43	7, 16, 31, 46	0
1	H	513/521 (98%)	0.42	60 (11%) 10 12	8, 19, 40, 55	0
All	All	4106/4168 (98%)	0.24	307 (7%) 22 24	7, 18, 37, 62	0

The worst 5 of 307 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	523	VAL	7.9
1	E	523	VAL	7.8
1	D	523	VAL	6.9
1	B	277	HIS	6.6
1	F	523	VAL	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MDO	H	149	13/14	0.91	0.10	16,19,26,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	MDO	B	149	13/14	0.93	0.08	16,19,23,26	0
1	MDO	F	149	13/14	0.95	0.08	12,15,20,21	0
1	MDO	G	149	13/14	0.95	0.07	13,14,17,19	0
1	MDO	A	149	13/14	0.95	0.06	15,18,21,21	0
1	MDO	C	149	13/14	0.96	0.05	13,14,18,19	0
1	MDO	D	149	13/14	0.96	0.07	18,20,22,24	0
1	MDO	E	149	13/14	0.96	0.06	12,14,18,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PMI	B	701	14/14	0.90	0.12	26,29,31,31	0
2	PMI	H	701	14/14	0.91	0.12	25,31,32,33	0
2	PMI	E	701	14/14	0.92	0.11	20,24,26,27	0
2	PMI	D	701	14/14	0.93	0.10	26,27,29,32	0
2	PMI	A	701	14/14	0.93	0.10	23,26,28,29	0
2	PMI	F	701	14/14	0.93	0.11	21,25,27,28	0
2	PMI	C	701	14/14	0.93	0.10	19,22,23,23	0
2	PMI	G	701	14/14	0.95	0.09	21,22,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.